NOAA Technical Report NMFS SSRF-675

Proceedings of the International Billfish Symposium Kailua-Kona, Hawaii, 9-12 August 1972 Part 3. Species Synopses

RICHARD S. SHOMURA and FRANCIS WILLIAMS (Editors)

SEATTLE, WA
June 1975

noaa NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION / National Marine Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Special Scientific Report—Fisheries Series

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The Special Scientific Report—Fisheries series was established in 1949. The series carries reports on scientific investigations that document long-term continuing programs of NMFS, or intensive scientific reports on studies of restricted scope. The reports may deal with applied fishery problems. The series is also used as a medium for the publication of bibliographies of a specialized scientific nature.

NOAA Technical Reports NMFS SSRF are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained (unless otherwise noted) from D83, Technical Information Division, Environmental Science Information Center, NOAA, Washington, D.C. 20235. Recent SSRF's are:

619. Macrozooplankton and small nekton in the coastal waters off Vancouver Island (Canada) and Washington, spring and fall of 1963. By Donald S. Day, January 1971, iii + 94 pp., 19 figs., 13 tables.

620. The Trade Wind Zone Oceanography Pilot Study. Part IX: The sea-level wind field and wind stress values, July 1963 to June 1965. By Gunter R. Seckel. June 1970, iii + 66 pp., 5 figs.

621. Predation by sculpins on fall chinook salmon, *Oncorhynchus tshawytscha*, fry of hatchery origin. By Benjamin G. Patten. February 1971, iii + 14 pp., 6 figs., 9 tables.

622. Number and lengths, by season, of fishes caught with an otter trawl near Woods Hole, Massachusetts, September 1961 to December 1962. By F. E. Lux and F. E. Nichy. February 1971, iii + 15 pp., 3 figs., 19 tables.

623. Apparent abundance, distribution, and migrations of albacore, *Thunnus alalunga*, on the North Pacific longline grounds. By Brian J. Rothschild and Marian Y. Y. Yong. September 1970, v + 37 pp., 19 figs., 5 tables.

624. Influence of mechanical processing on the quality and yield of bay scallop meats. By N. B. Webb and F. B. Thomas. April 1971, iii + 11 pp., 9 figs., 3 tables.

625. Distribution of salmon and related oceanographic features in the North Pacific Ocean, spring 1968. By Robert R. French, Richard G. Bakkala, Masanao Osako, and Jun Ito. March 1971, iii + 22 pp., 19 figs., 3 tables.

626. Commercial fishery and biology of the freshwater shrimp, *Macrobrachium*, in the Lower St. Paul River, Liberia, 1952-53. By George C. Miller. February 1971, iii + 13 pp., 8 figs., 7 tables.

627. Calico scallops of the Southeastern United States, 1959-69. By Robert Cummins, Jr. June 1971, iii + 22 pp., 23 figs., 3 tables.

628. Fur Seal Investigations, 1969. By NMFS, Marine Mammal Biological Laboratory. August 1971, 82 pp., 20 figs., 44 tables, 23 appendix A tables, 10 appendix B tables.

629. Analysis of the operations of seven Hawaiian skipjack tuna fishing vessels, June-August 1967. By Richard N. Uchida and Ray F. Sumida. March 1971, v + 25 pp., 14 figs., 21 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

630. Blue crab meat. I. Preservation by freezing. July 1971, iii + 13 pp., 5 figs., 2 tables. II. Effect of chemical treatments on acceptability. By Jurgen H. Strasser, Jean S. Lennon, and Frederick J. King. July 1971, iii + 12 pp., 1 fig., 9 tables.

631. Occurrence of thiaminase in some common aquatic animals of the United States and Canada. By R. A. Greig and R. H. Gnaedinger. July 1971, iii + 7 pp., 2 tables.

632. An annotated bibliography of attempts to rear the larvae of marine fishes in the laboratory. By Robert C. May. August 1971, iii + 24 pp., 1 appendix I table, 1 appendix II table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

633. Blueing of processed crab meat. II. Identification of some factors involved in the blue discoloration of canned crab meat *Callinectes sapidus*. By Melvin E. Waters. May 1971, iii + 7 pp., 1 fig., 3 tables.

634. Age composition, weight, length, and sex of herring, *Clupea pallasii*, used for reduction in Alaska, 1929-66. By Gerald M. Reid. July 1971, iii + 25 pp., 4 figs., 18 tables.

635. A bibliography of the blackfin tuna, *Thunnus atlanticus* (Lesson). By Grant Beardsley and David C. Simmons. August 1971, 10 pp. For sale by the Superintendent Documents, U.S. Government Printing Office, Washington, D.C. 20402.

636. Oil pollution on Wake Island from the tanker *R. C. Stoner*. By Reginald Gooding. May 1971, iii + 12 pp., 8 figs., 2 tables. For sale by the Superintendent Documents, U.S. Government Printing Office, Washington, D.C. 20402.

637. Occurrence of larval, juvenile, and mature crabs in the vicinity of Beaufort Inle North Carolina. By Donnie L. Dudley and Mayo H. Judy. August 1971, iii + 10 pp., 1 fi 5 tables. For sale by the Superintendent of Documents, U.S. Government Printing Offi Washington, D.C. 20402.

638. Length-weight relations of haddock from commercial landings in New Englan 1931-55. By Bradford E. Brown and Richard C. Hennemuth. August 1971, v + 13 pp., figs., 6 tables, 10 appendix A tables. For sale by the Superintendent of Documents, U. Government Printing Office, Washington, D.C. 20402.

639. A hydrographic survey of the Galveston Bay system, Texas 1963-66. By E. J. Pulle W. L. Trent, and G. B. Adams. October 1971, v + 13 pp., 15 figs., 12 tables. For sale by t Superintendent of Documents, U.S. Government Printing Office, Washington, D. 20402.

640. Annotated bibliography on the fishing industry and biology of the blue cra *Callinectes sapidus*. By Marlin E. Tagatz and Ann Bowman Hall. August 1971, 94 pp. F sale by the Superintendent of Documents, U.S. Government Printing Office, Washingto D.C. 20402.

641. Use of threadfin shad, *Dorosoma petenense*, as live bait during experimental po and-line fishing for skipjack tuna, *Katsuwonus pelamis*, in Hawaii. By Robert T. Iversen. August 1971, iii + 10 pp., 3 figs., 7 tables. For sale by the Superintendent Documents, U.S. Government Printing Office, Washington, D.C. 20402.

642. Atlantic menhaden *Brevoortia tyrannus* resource and fishery—analysis of decli By Kenneth A. Henry. August 1971, v + 32 pp., 40 figs., 5 appendix figs., 3 tables, appendix tables. For sale by the Superintendent of Documents, U.S. Government Printi Office, Washington, D.C. 20402.

643. Surface winds of the southeastern tropical Atlantic Ocean. By John M. Steigner a Merton C. Ingham. October 1971, iii + 20 pp., 17 figs. For sale by the Superintendent Documents, U.S. Government Printing Office, Washington, D.C. 20402.

644. Inhibition of flesh browning and skin color fading in frozen fillets of yellowe snapper (*Lutzanus vivanus*). By Harold C. Thompson, Jr., and Mary H. Thompso February 1972, iii + 6 pp., 3 tables. For sale by the Superintendent of Documents, U Government Printing Office, Washington, D.C. 20402.

645. Traveling screen for removal of debris from rivers. By Daniel W. Bates, Ernest Murphey, and Martin G. Beam. October 1971, iii + 6 pp., 6 figs., 1 table. For sale by Superintendent of Documents, U.S. Government Printing Office, Washington, D. 20402.

646. Dissolved nitrogen concentrations in the Columbia and Snake Rivers in 1970 a their effect on chinook salmon and steelhead trout. By Wesley J. Ebel. August 1971, iii + pp., 2 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Governme Printing Office, Washington, D.C. 20402.

647. Revised annotated list of parasites from sea mammals caught off the west coast North America. By L. Margolis and M. D. Dailey. March 1972, iii + 23 pp. For sale by Superintendent of Documents, U.S. Government Printing Office, Washington, D. 20402.

Continued on inside back cover.

NOAA Technical Report NMFS SSRF-675

Proceedings of the International Billfish Symposium Kailua-Kona, Hawaii, 9-12 August 1972 Part 3. Species Synopses

RICHARD S. SHOMURA and FRANCIS WILLIAMS (Editors)

SEATTLE, WA
June 1975

UNITED STATES
DEPARTMENT OF COMMERCE
Rogers C. B. Morton, Secretary

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION
Robert M. White, Administrator

National Marine
Fisheries Service
Robert W. Schoning, Director

For sale by the Superintendent of Documents, U.S. Government
Printing Office, Washington, D.C. 20402

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Page

RIVAS, L. R. Synopsis of biological data on blue marlin, *Makaira nigricans* Lacépède, 1802 1
NAKAMURA, IZUMI. Synopsis of the biology of the black marlin, *Makaira indica* (Cuvier), 1831 17
ROBINS, C. RICHARD. Synopsis of biological data on the longbill spearfish, *Tetrapturus pfluegeri* Robins and de Sylva 28
KIKAWA, SHOJI. Synopsis on the biology of the shortbill spearfish, *Tetrapturus angustirostris* Tanaka, 1914 in the Indo-Pacific areas 39
MATHER, F. J., III, H. L. CLARK, and J. M. MASON, JR. Synopsis of the biology of the white marlin, *Tetrapturus albidus* Poey (1861) 55
BEARDSLEY, G. L., JR., N. R. MERRETT, and W. J. RICHARDS. Synopsis of the biology of the sailfish, *Istiophorus platypterus* (Shaw and Nodder, 1971) 95
DE SYLVA, DONALD P. Synopsis of biological data on the Mediterranean spearfish, *Tetrapturus belone* Rafinesque 121
UEYANAGI, SHOJI, and PAUL G. WARES. Synopsis of biological data on striped marlin, *Tetrapturus audax* (Philippi), 1887 132

Digitized by the Internet Archive
in 2025

https://archive.org/details/noaa-technical-report-nmfs-ssrf_1975-06_675-part-3

Synopsis of Biological Data on Blue Marlin, *Makaira nigricans* Lacépède, 1802

L. R. RIVAS[1]

IDENTITY

1.1 Nomenclature

1.11 Valid name

Makaira nigricans Lacépède, 1802.

1.12 Objective synonymy

Tetrapturus herschelli Gray, 1838.
Tetrapturus amplus Poey, 1860.
Tetrapturus mazara Jordan and Snyder, 1901.
Makaira bermudae Mowbray, 1931.
Eumakaira nigra Hirasaka and Nakamura, 1947.
Makaira perezi de Buen, 1950.
Istiompax howardi Whitley, 1954.

1.2 Taxonomy

1.21 Affinities

Suprageneric

Phylum Chordata
Subphylum Vertebrata
Superclass Gnathostomata
Class Teleostomi
Subclass Actinopterygii
Order Perciformes
Suborder Xiphioidei
Superfamily Xiphioidae
Family Istiophoridae

Generic

Genus *Makaira* Lacépède, 1802.

Makaira Lacépède, 1802 (type species *Makaira nigricans* Lacépède, 1802).

Statement of generic concept: According to my interpretation, the black marlin, *indica,* should be placed in the genus *Makaira* in addition to *nigricans* (see 1.22 Taxonomic status). Therefore, the genus *Makaira* comprises only two species.

Subjective generic synonyms:

Tetrapturus (not of Rafinesque, 1810) Cuvier *in* Cuvier and Valenciennes, 1831 (type species *Tetrapturus indicus* Cuvier, 1832).
Machaera Cuvier, 1832 (type species *Machaera velifera* Cuvier, 1832 by monotypy).
Macaria Nardo, 1833 (amended spelling of *Makaira).*
Istiompax Whitley, 1931 (type species *Istiompax australis* Whitley, 1931, nomen nudum).
Eumakaira Hirasaka and Nakamura, 1947 (type species *Eumakaira nigra* Hirasaka and Nakamura, 1947).
Orthocraeros Smith, 1956 (type species *Makaira bermudae* Mowbray, 1931).

Diagnosis: Makaira is distinguished from all the other genera of the family Istiophoridae (*Istiophorus, Tetrapturus)* by the following characters:

Anterior height of first dorsal fin less than depth of body at origin of first dorsal fin. Second predorsal length 75 to 79% of body length. Precaudal vertebrae 11, caudal 13. Flesh pale. Maximum weight more than 682 kg (1,500 lb), often weighing more than 182 kg (400 lb), and usually more than 114 kg (250 lb).

Specific

Identity of type specimen: A sketch of a 365 kg (803 lb) *Makaira nigricans,* blue marlin, sent to Lacépède (see Morrow, 1959b).

Type locality: Ile de Ré, Bay of Biscay, France.

Diagnosis: Makaira nigricans is distinguished from *M. indica* by the following characters: pectoral joint flexible, the fin folding flat against the sides of body without breaking joint. Median ridge in isthmus present. Anterior height of first dorsal fin about equal to, or longer than distance between origin of first dorsal fin and insertion of pectoral fin. Lateral line not visi-

[1]Southeast Fisheries Center, National Marine Fisheries Service, Miami, Fla.; present address: Panama City Laboratory, Southeast Fisheries Center, National Marine Fisheries Service, NOAA, Panama City, FL 32401.

ble externally, complex, forming a reticulate pattern covering sides of body (visible when large patch of skin is removed and dried). Dorsal spines 39 to 46, usually 41 to 43. Anal spines 14 to 17, usually 15 or 16. Pectoral rays 20 to 22, usually 21 or 22.

Material examined: Specimens studied and allocated to this species (as *Makaira ampla*) by me (Rivas, 1956) were examined in the field and discarded because of their large size. Records of measurements, counts, sex, gonad condition, and other characters for these and other specimens are in my files.

Subjective synonymy: See above under objective synonymy.

1.22 Taxonomic status

I have always held that *Makaira nigricans* constitutes a single pantropical species occurring in the Atlantic, Pacific, and Indian oceans (Rivas, 1956, as *M. ampla*). The same view was held by Royce (1957), Briggs (1960), Robins and de Sylva (1960), Jones and Silas (1964), and Morrow (1964). Recently, however, Nakamura, Iwai, and Matsubara (1968) have indicated that two forms are involved which they interpret as closely related but separate species. They consider *M. nigricans* as restricted to the Atlantic and *M. mazara* as occurring in the Indo-Pacific. The only evidence given by these authors is a slight difference in the lateral line pattern; however, it should be noted that the specimens they compared were of different sizes. I disagree with Nakamura et al. (1968) that two species are involved and maintain that there is only one pantropical species.

1.23 Subspecies

As a corrollary to the above discussion it may be indicated that several authors have considered the Atlantic and Indo-Pacific forms as subspecies. This has been discussed by Nakamura et al. (1968). At the present state of knowledge, however, no subspecific division should be considered.

1.24 Standard common names and vernacular names.

Standard common names: Blue marlin (English-speaking countries), Aguja de Casta (Cuba), Marlin azul (Mexico, Central America, Venezuela), Pieto (Madeira), Kurokajiki (Japan), lan fu yii (China), taketonga (New Zealand).

Vernacular names: Cuban black marlin (United States), Castero (Cuba), Katsuokui, Katokui, Kuroka, Kurokawa, Shiroka, Shirokajiki, Gemba (Japan), Njiachi (Okinawa).

1.3 Morphology

1.31 External morphology

For description of spawn, larvae, and adolesce see 3.17, 3.22, 3.23.

Individual and ontogenetic variation: Data a comments on individual and ontogenetic variat (allometric growth) have been presented by Ri (1956), Royce (1957), and Nakamura et al. (19 Much more information is needed.

Geographic variation: As yet, information is available to establish the extent of geographic va tion (see also comments under 1.22 and 1.23 abo This also applies to the morphological definitio subpopulations.

1.32 Cytomorphology

Nothing is known about chromosome number a other aspects of the cytomorphology of this speci

1.33 Protein specificity

No information on comparative serology available for this species.

2 DISTRIBUTION

2.1 Total Area

Geographic distribution: The geographic a seasonal distributions are fairly well known as t result of data published by Ueyanagi et al. (1970) a Mather, Jones, and Beardsley (1972) for the Atlan Ocean and by Howard and Ueyanagi (1965) a Nakamura et al. (1968) for the Pacific Ocean.

In the Atlantic, the latitudinal range of blue mar varies seasonally and extends from about lat. 45°N about lat. 35°S. Monthly distribution of catches Japanese longliners shows two main seasonal conce trations. One occurs from January through April, the southwestern Atlantic, between lat. 5° and 30 and the other from June through October, in t northwestern Atlantic, between lat. 10° and 35° May, November, and December appear to be tra sitional months. The species is less abundant in t eastern Atlantic where it occurs mostly off Afri between lat. 25°N and 25°S.

In the Pacific, the latitudinal range of the bl marlin is also seasonal and extends from about la 48°N to about lat. 48°S. As in the Atlantic, month distribution of catches by Japanese longliners sho two main seasonal concentrations. One center of co centration occurs from December through March, the western and central South Pacific, between lat. and 26°S, and the other from May through Octob

he western and central North Pacific, between lat. nd 24°N. In April and November the fish tend to centrate in the equatorial Pacific between lat. N and 10°S. As in the Atlantic, the blue marlin in Pacific Ocean is less abundant towards the East. he extreme eastern Pacific, off the continental st, the fish usually occurs between lat. 23°N and . Kume and Joseph (1969b) indicate that blue lin occur in heaviest concentrations at about lat. S and west of long. 110°W, north of lat. 13°N; ever, their relative abundance is quite low. Kume Joseph (1969a) record longline catches along the uvian coast as far south as lat. 12°S.

the Indian Ocean, the blue marlin is known to oc- around Ceylon, Mauritius, and off the east coast frica (Morrow, 1959a, 1964). Off the east coast of ca, between the equator and lat. 13°S, the blue lin is apparently abundant during the southeast nsoon period, April to October (Merrett, 1971).

2.2 Differential Distribution

2.21 Spawn, larvae, and juveniles

as of occurrence and seasonal variations: Three ae were reported by Gehringer (1956) from the tern Atlantic off Georgia (lat. 30°57′N, long. 37′W). These were later identified as blue marlin ae by Ueyanagi and Yabe (1959). Two postlarvae e recorded from Jamaica by Caldwell (1962). hmeyer and Bullis (1968) recorded four larvae and tlarvae from the western Atlantic (two off Cat Cay, amas, one at lat. 32°06′N, long. 72°00′W and one about 40 miles northest of Fort Pierce, Fla.). tlett and Haedrich (1968) recorded 85 larvae from Brazil between Cabo de São Roque and lat. 26°S. eyanagi and Yabe (1959) recorded about 400 lar- from the western Pacific, and Ueyanagi (1964) orted that 1,015 larvae had been collected in the ific. Ueyanagi et al. (1970) have given the seasonal ribution of larval blue marlin for the Atlantic.

2.22 Adults

as of occurrence and seasonal variations: In most as within its range the occurrence of adult male lin appears to be seasonal and in some regions re is also a seasonal variation by sex.

he seasonal variation and relative abundance of e marlin in the northern Gulf of Mexico have been ussed by Nakamura and Rivas (1972). Off Puerto o the largest numbers of blue marlin have been ght during August, September, and October, and smallest catches have been made in December dman, 1962). In Hawaii, blue marlin catches are highest in summer and lowest in winter (Royce, 7; Strasburg, 1970).

ery large blue marlin, probably females, are orted to occur off the south coast of Jamaica in the mer while similar large fish are reported off the north coast in the winter (de Sylva, 1963). Nine blue marlin specimens taken in late December and early January from Jamaican waters were examined by me; all were males. Off Puerto Rico, males and females occur in about equal numbers in July and August; larger males tend to appear in May and there is a sudden increase of small males in September (Erdman, 1968).

In the eastern Pacific, extreme differences in sex ratios between certain regions suggest strongly the possibility that blue marlin segregate into distinct areal groups according to sex (Kume and Joseph, 1969b).

2.3 Determinants of Distribution Changes

Effects of ecological determinants: The distribution of the blue marlin is mostly confined to the tropics within the 24°C surface isotherms of both hemispheres, and is noted to shift northward and southward according to season. In the western Atlantic off the northeastern United States, blue marlin have been taken in waters with surface temperatures ranging from 23.9° to 28.3°C (Squire, 1962). Blue marlin catches were recorded at 47 longline stations occupied during the past 22 yr by National Marine Fisheries Service (NMFS) vessels in the Caribbean Sea, Gulf of Mexico, and off the southeastern United States; surface temperatures at these stations ranged from 21.7° (February) to 30.5°C (July).

In certain regions water masses and currents apparently affect the horizontal and seasonal distribution of the blue marlin. For example, in the northern Gulf of Mexico the occurrence of blue marlin seems to be controlled by the so-called "Loop Current." This current, an extension of the Caribbean Current, flows northward into the Gulf through the Yucatan Channel. Its northward extent is seasonal and reaches its maximum in the summer.

Water color also affects the occurrence of the blue marlin, at least in the northern Gulf of Mexico, where the fish shows preference for blue water (Nakamura and Rivas, 1972).

There is indication of a periodic fluctuation in abundance which is probably due to ecological interaction with other species of marlin. For example, within the fishing season (April to November) in the northern Gulf of Mexico, anglers tend to catch more blue marlin when the white marlin catch is lowest and vice versa (Nakamura and Rivas, 1972). A similar phenomenon was reported by Strasburg (1970) for the blue and striped marlins around Hawaii. Strasburg suggests that perhaps each responds to some environmental factor, such as temperature or food, in a way which excludes the other.

2.4 Hybridization

There are no records of hybridization for the blue marlin.

3 BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

The species is heterosexual. No occurrence of hermaphrodism or intersexuality has been observed. Males and females are indistinguishable externally, but the females attain a much larger size than the males.

For the central Pacific, Royce (1957) indicated that females have been taken at weights greater than 727 kg (1,600 lb) whereas the largest male weighed only 99 kg (218 lb). Around Hawaii most of the males weigh between 45 and 91 kg (100 and 200 lb), whereas most of the females weigh between 91 and 227 kg (200 and 500 lb) (Strasburg, 1970). In Puerto Rico, the largest male examined by Erdman (1968) weighed 135 kg (297 lb) and the largest female 370 kg (814 lb). Data from the Japanese longline fishery show that males seldom exceed 116 kg (255 lb) and that most males weigh from 39 to 80 kg (85 to 175 lb). De Sylva (1963) recorded a 140-kg (308-lb) male from Bimini, Bahamas, and Strasburg (1970) recorded a 133-kg (293-lb) male from Hawaii. Of 30 blue marlin examined by me in the northwestern Atlantic, 18 were males weighing 46 to 94 kg (101 to 207 lb) and 12 were females weighing 59 to 277 kg (130 to 609 lb). Usually, a blue marlin heavier than 136 kg (300 lb) is a female.

According to de Sylva (1963) protandry, the condition whereby a fish begins its life as a male and subsequently changes into a female, has been considered for the blue marlin.

3.12 Maturity

The age at which sexual maturity is reached cannot be calculated because methods of age determination have not been developed. As already indicated, males are smaller than females and size has to be considered. The smallest sexually mature male reported by Erdman (1968) from Puerto Rico weighed 35 kg (76 lb) and the smallest ripe male reported from Jamaica by de Sylva (1963) weighed 44 kg (97 lb). Of 26 males from the central Pacific examined by Royce (1957) the smallest specimen found with milt in the central duct of the testes weighed 42 kg (93 lb). The smallest sexually mature male reported for the western Indian Ocean by Merrett (1971) weighed 41 kg (90 lb). The smallest sexually mature female reported by Erdman (1968) from Puerto Rico weighed 61 kg (135 lb) and the smallest mature female reported by Merrett (1971) from the western Indian Ocean weighed 47 kg (103 lb). Of 1,152 females studied by Kume and Joseph (1969b) from the eastern Pacific, the smallest specimens found in a condition believed to be near spawning were about 200 cm long (eye to fork; no weight given). The indications are, therefore, that males reach sexual maturity at a weight of 35 to 44 kg (76 to 97 lb) and females at 47 to 61 kg (103 to lb).

3.13 Mating

Probably polygamous.

3.14 Fertilization

Evidence indicates that fertilization is externa

3.15 Gonads

The relation of gonad weight to body weight determined by Erdman (1968) for 14 males and females from Puerto Rico. Males weighed from 4 123 kg (103 to 270 lb) and their testes from 55 g to kg (0.12 to 5.30 lb). Females weighed from 33 to 29 (72 to 643 lb) and their ovaries from 9.1 g to 6.5 (0.02 to 14.2 lb). The weight of the testes, as the cent of body weight, ranged from 0.10 to 2.30% in males and from 0.03 to 3.60% in the females. Acc ing to Erdman less than 0.10% for the testes ovaries corresponds to sexually immature individu In the males, 0.10% to less than 0.70% correspond the developing stage with few to many spermatoc and few or no sperm. Prespawning stage correspo to a range of 0.70% to less than 2.30% with many s matocytes and many sperm. At the ripe stage relative weight of the testes decreases to a rang 0.60% to less than 1.60% but with few or no s matocytes and many sperm. At the postspawn stage, 0.10% to less than 0.20%, there are no s matocytes but many sperm. In the females, 0.10 less than 1.0% corresponds to the developing st with oocytes and no ova or a few developing ones, no easily visible eggs. The subripe stage correspo to a range of 1.00% to less than 3.60% with advan oocytes and pale-yellow eggs. At the ripe stage weight of the ovaries decreases to less than 1.50 body weight, but is still 1.00% or more, with advan oocytes, transparent eggs, and some postovula follicles. At the postspawning stage, 0.50% to less t 1.00%, there are some atretic bodies, many em follicles, and degenerating oocytes.

Krumholz (1958) gives a range of 0.80 to 1.9 with an average of 1.43%, for the weight of teste relation to body weight in blue marlin from Bahamas.

The relationship between ovary weight and b length for 1,152 blue marlin from the eastern Pa was given by Kume and Joseph (1969b). O weights ranged from a few grams to 14 kg (31 lb) lengths (eye to fork) from about 140 to about 338 In the majority of specimens the ovaries weighed than 1 kg (2.2 lb) and probably were not in a spawn condition.

The length of the gonads, as a percentage of b length, can be used as an index of sexual maturit

e testes and ovaries. Of 12 females studied by me e range was 7.6 to 13.2% in five specimens; their aries were in a resting condition. In the other seven h, the range was 16.7 to 20.8% and their ovaries ere developing but not yet ripe. In five males udied, the range was 22.0 to 32.4% and their testes ere all ripe with much milt running in the ducts. No ales in the resting or developing stage were available r comparison.

There is no information available on fecundity.

3.16 Spawning

Mather et al. (1972) believe that the two widely parated concentrations of blue marlin in the estern Atlantic represent separate spawning pop- ations. They indicate that the evidence suggests at the blue marlin in the North Atlantic spawn ainly from July through September and those in the outh Atlantic spawn in February and March. These thors further state that it is unlikely that a single opulation of blue marlin spawns at two widely parate locations at different times of the year.

According to Erdman (1968) the occurrence of ripe stes from May to November indicates a protracted awning season for the blue marlin off Puerto Rico. e found females with well-formed eggs from late ay to September and a female with flowing ripe eggs September. Erdman assumes that July and August e the peak spawning months because during that riod the sex ratio is nearly equal.

De Sylva (1963) suggested May and June as the awning season for the blue marlin found in waters f Florida and Bahamas. He also stated that mature h taken off Cape Hatteras in June appear to have en recent spawners and that females taken in ptember and October from Jamaican and Puerto ican waters have long since spawned.

For the Pacific Ocean, information on spawning of e blue marlin was summarized by Howard and eyanagi (1965) and Strasburg (1970). From the oc- rrence of larvae, gonad condition, and sex ratio, awning is assumed to take place between about lat. °N and 10°S throughout the year. During the mmer season, however, spawning is assumed to ke place in the broader latitudinal area bounded by t. 30°N and 30°S. Males with freely flowing milt ve been captured in the central Pacific from bruary to October, and May to July has been garded as the spawning season in the Philippine ea.

Kume and Joseph (1969b) indicated that in the uthwestern portions of the eastern Pacific Ocean, ue marlin spawn primarily during the southern mmer.

Among the blue marlin specimens examined by yce (1957) from the central Pacific no ripe females ere found, but a number of males had freely flowing ilt in the gonads from February through October.

Off Puerto Rico, the annual average male:female sex ratio was 4:1 based on 328 specimens examined by Erdman (1968). As already indicated, the 47 specimens examined by him in July and August show a more nearly equal ratio of 25:22. Every September, there is a sudden increase in the catch of males and the ratio changes to 4.5:1.

Of 39 specimens examined by de Sylva (1963) in Jamaica during early October, 37 were males for a ratio of 18.5:1.

Nakamura and Rivas (1972) recorded sex ratios for the northern Gulf of Mexico during the sport fishing season (June through October) as follows. Off the mouth of the Mississippi River (South Pass) the male:female sex ratio was 1:5.6 in 1967, 1:7.7 in 1968, 1:4.8 in 1969, 1:8.0 in 1970, and 1:33 in 1971. Off northwest Florida the male:female sex ratio was 1:2.5 (Destin) and 1:2 (Panama City).

In the central Pacific, according to data presented by Royce (1957), the mean annual male:female sex ratio is 1.2:1.

3.17 Spawn

Subripe ova are opaque, white to yellow, and 0.3 to 0.5 mm in diameter. Transparent, spherical eggs flowing out of a ripe ovary measure 1 mm in diameter (Erdman, 1968).

3.2 Preadult Phase

3.21 Embryonic phase

There is no information on the embryonic phase.

3.22 Larval phase

Various workers have contributed to the still very incomplete knowledge of the larval and juvenile stages of the blue marlin. Their findings are summarized below.

Gehringer (1956) described three unidentified larvae 11.3, 21.0, and 45 mm long from the Atlantic. These were later identified as blue marlin larvae by Ueyanagi and Yabe (1959). These authors also described eight specimens, 2.8 to 23.2 mm in length, selected from about 400 larval blue marlin collected in the western Pacific. Caldwell (1962) described two postlarval blue marlin measuring 201.4 and 206 mm in length from Jamaica. Ueyanagi (1964) reported that as of that date 1,015 larval blue marlin, 3 to 33 mm (mostly under 7 mm), had been collected in the Pacific by the Nankai Regional Fisheries Research Laboratory. Eschmeyer and Bullis (1968) described three larval (33.5, 35.3, and 51.5 mm) and one postlarval (194.1 mm) blue marlin from the western Atlantic. Bartlett and Haedrich (1968) reported on 85 larval blue marlin, 4.9 to 32.0 mm long from the southwestern Atlantic.

In summary, larval stages are fairly well known

from about 3 to 52 mm. There is a gap between about 52 and 194 mm and another, at the postlarval and juvenile stages, between 206 and 846 mm. The latter, described by de Sylva (1958) is the smallest young stage known. Ueyanagi (1963) discussed methods of identification for larval blue marlin based on Indo-Pacific material.

3.23 Adolescent phase

As already discussed above, under 3.12, males weighing less than 35 kg (76 lb) and females weighing less than 47 kg (103 lb) have not reached sexual maturity. Individuals in this category, therefore, may be considered to be in the young stage or adolescent phase provided they are past the juvenile stage. Two specimens of young blue marlin, 846 and 1,320 mm in length (tip of lower jaw to fork) were described by de Sylva (1958) from the Bahamas and southeast Florida, respectively. The smaller specimen weighed 2.3 kg (5 lb) and the other 13.9 kg (30.5 lb).

3.3 Adult Phase (Mature Fish)

3.31 Longevity

Estimates of blue marlin age may be obtained from analyses of modal progressions in length and weight frequency. Also, the sex must be known since males are much smaller than females. No definite information on age is, as yet, available for the blue marlin.

3.32 Hardiness

As already discussed under 2.22, blue marlin have been taken at surface temperatures as high as 30.5°C (July) and as low as 21.7°C (February).

Individuals captured, tagged, and liberated appear to suffer no ill effects since a few have been recaptured after several months of freedom.

3.33 Competitors

There is no definite information on this subject but there are indications that in the Atlantic, the white marlin, *Tetrapturus albidus*, might compete with the blue marlin for food. In the Indo-Pacific, the striped marlin, *T. audax*, and the black marlin, *Makaira indica*, might compete with the blue marlin for food. To a lesser extent the sailfish, *Istiophorus platypterus*, and the spearfishes, *T. belone, T. pfluegeri, T. angustirostris*, and *T. georgii*, might also compete with the blue marlin for food.

3.34 Predators

My own observations agree with the well-known fact that sharks frequently attack hooked blue marlin especially if the fish is tired after a long fight. It is not known whether sharks will attack a free-swimming, healthy, blue marlin, but I have seen a mako shark attack, kill, and eat a free-swimming broadbill swo fish.

3.35 Parasites, diseases, injuries, and abnormalities

Parasites and diseases: On several occasions, I ha observed ectoparasitic calgoid copepods on the he of fresh-caught blue marlin. Cressey and Lachr (1970) reported that the marlinsucker, *Remora teochis*, occurs on the body and in the gill cavity of t blue marlin where they feed on ectoparasi copepods. Their examination of marlinsuck stomachs revealed the presence of *Caligus* a *Pennella* on blue marlin. Ectoparasitic copepods not seem to affect the physiology, behavior, or fo value of the blue marlin. The copepods are apparen kept in check by the marlinsucker who maintains symbiotic relationship with the blue marlin and a as a "cleaner."

The trematode *Capsala poeyi* from the skin of t blue marlin has been reported from Cuba by Viguer (1935), from the Gulf of Mexico by Manter (1954) a Koratha (1955), and from the Pacific by Iversen a Hoven (1958).

Stomach ulcers were found in 10 of 114 blue mar examined by Iversen and Kelley (1974) in Hawa These ulcers were noncancerous and morphologica similar to gastric ulcers found in many mammals, i cluding marine mammals. Iversen and Kelley belie that endoparasites or mechanical injury to t stomach lining, from sharply pointed food items, a the most likely cause.

Injuries and abnormalities: Some of the blue mar specimens examined by me have had some scars malformed fins or bills. On several occasions the tip the bill has been broken off. From time to time sp fishermen have told me of seeing blue marlin witho a bill, but I have not personally seen specimens w this abnormality. Moore (1950) reported a 248- (545-lb) blue marlin from Hawaii without a spe The appearance of the specimen indicated that t spear had been lost by injury and there was no indi tion that the loss was at all recent. Moore also me tioned that the specimen was equal in condition normal blue marlin and that apparently the spear not necessary for natural and adequate feeding. also remarked that according to the operators of t Honolulu market spearfishes without a spear h been observed before but that such occurrences w very rare.

Erdman (1957) recorded a blue marlin with the b missing and healed at the point of rupture. The upp jaw was shorter than the lower but the fish had actu ly more food in its stomach than four other bl marlin captured the same day.

Broken bills in marlin are apparently caused by t pugnacity of these fishes, according to Smith (195 He states that many floating bales of rubber w

ken tips of marlin bills in them were found in the zambique Channel and believes that marlin must berately charge floating or submerged objects sibly to secure food or from plain aggressive-s.

blue marlin with a bent bill was reported by hinnikov (1970) from off West Africa.

3.4 Nutrition and Growth

3.41 Feeding

ccording to data from the northern Gulf of Mexico sented by Nakamura and Rivas (1972) blue marlin to a surface trolled bait more often in the morning ween the hours of 1000 and 1100 and least often ween 1200 and 1300 (central standard time). There greement in this for both South Pass, La., and the thwest Florida areas. The South Pass data also w that, in the afternoon, there is another well-rked peak between 1500 and 1600 but the thwest Florida data show a steady decrease from 0 on.

Many sport fishermen have told me that they have d without success to raise marlin during the night surface trolling. I have tried, also without success. rom the information discussed above, it may be tatively concluded that blue marlin do not feed at surface at night and that there is a marked feeding iod in the morning between 1000 and 1100.

De Sylva (1963) stated that during the 1963 naica International Fishing Tournament more rlin were raised in the afternoon between 1400 and 0 than at any other time of day. His results, vever, were given in terms of the number of fish ed during 2-h intervals as a percentage of the total nber of fish raised. No data on the effort expended given.

have heard many discussions among sport fishing des and anglers about whether or not the blue and er marlins use their bill in capturing prey. These cussions, however, refer mostly to prey in the form rtificial or natural dead baits trolled at the surface uniform speed, usually 4 to 8 knots. In my opin-based on many years of observation, marlins do use their bill when taking a trolled bait. The ral thrusts of the bill, which appear to be aimed at ing the bait, are apparently the result of rapid nges in direction of movement of the fish and/or effect of its swimming motions when the bill and d are partly out of the water.

s to the use of the bill when the fish is pursuing -swimming prey, there is, again, difference of nion but the general concensus is that the bill is generally used to stun the prey.

vchinnikov (1970) discussed the use of the bill by lins and concluded that it is not of importance in capture of food. As pointed out by Ovchinnikov as discussed above under 3.35 *Injuries and abnormalities*, marlins without bills or with broken or malformed bills are as healthy as the normal fish.

The various incidents of fish, boats, and various other floating objects impaled by marlin bills cited in the literature are, in my opinion, accidents resulting from feeding. It is well known that small fish congregate under such floating objects and that a marlin, in attempting to catch a fish too close to the floating object, may accidently impale it with its bill.

Tinsley (1964) gives a detailed discussion on the use of the bill in the sailfish which could well apply to the blue and other marlins.

3.42 Food

The literature shows that the types of food eaten by the blue marlin vary somewhat with the region where they occur. It is also indicated that the blue marlin feeds at or near the surface and in deep water, and near shore as well as out in the open sea.

Stomachs of blue marlin contained mostly squid in the Philippine Sea (Nakamura, 1942), tunalike fishes in New Zealand (Baker, 1966) and the central Pacific (Royce, 1957), and dolphin (*Coryphaena*) and tunalike fishes, especially frigate mackerel (*Auxis*), in the Bahamas (Krumholz and de Sylva, 1958). On a volumetric basis, tunalike fishes constitute more than 85% of the diet in Hawaii (Strasburg, 1970). In Puerto Rico tunalike fishes, in both number and volume, were the chief food of the blue marlin examined by Erdman (1962). Frigate mackerel was the most frequent individual item, and whenever it occurred in abundance blue marlin fishing was better than average. De Sylva (1963) indicates that only a relatively few types of organisms are eaten by blue marlin in Jamaica, the tunalike fishes being the most important. In the northern Gulf of Mexico fishes, especially dolphin and scombrids, were the most important food items found in the stomachs of blue marlin (Nakamura and Rivas, 1972). In the tropical western Atlantic, Ovchinnikov (1970) indicates that the blue marlin feeds mostly on fishes and cephalopods.

The size range of the organisms eaten by the blue marlin is relatively large. Krumholz and de Sylva (1958) indicated that the overall range in length for fishes was from about 20 to 102 cm (about 8 to about 40 inches). One of the octopods was about 15 cm (about 6 inches) whereas the largest one was about 61 cm (about 24 inches) long. Erdman (1962) recorded from Puerto Rico a 135-kg (279-lb) blue marlin which had eaten a postlarval surgeonfish 38 mm long. He also mentioned a squid weighing up to 11 kg (23 lb) taken from the stomach of a blue marlin. Strasburg (1969) mentioned that a blue marlin was caught in Hawaii with a 29-kg (63-lb) bigeye tuna, *Thunnus obesus*, in its stomach. The marlin including the tuna weighed 340 kg (748 lb). Ovchinnikov (1970) reported from the Gulf of Guinea a 290-kg (638-lb) blue marlin

which had swallowed a bigeye tuna weighing about 50 kg (about 110 lb).

The blue marlin is known to feed at and near the surface, but there are indications that it may also feed in relatively deep water. Off Puerto Rico, deep-sea fishes, such as *Pseudoscopelus*, were found in blue marlin stomachs (Erdman, 1962). The deep-dwelling squirrelfish, *Holocentrus lacteoguttatus*, was found in Hawaiian blue marlin stomachs by Strasburg (1970).

3.43 Growth rate

There is very little information on growth rate although estimates could be obtained from analyses of modal progressions in length and weight frequencies. Royce (1957) presented data on weight frequency for central Pacific blue marlin but no modal progression can be ascertained. Kume and Joseph (1969b) presented and discussed data on the size composition of blue marlin in the eastern Pacific, but they could not estimate growth from progression of modal groups.

3.44 Metabolism

No information available.

3.5 Behavior

Feeding behavior is discussed above under 3.41. There is no information on reproductive behavior.

3.51 Migrations and local movements

Ovchinnikov (1970) mentioned that, in the western Atlantic, distribution by months indicate that the blue marlin shows a tendency to seasonal migrations. Mather et al. (1972) are of the opinion that longline catches have produced no evidence that the blue marlin moves between the northwestern and southwestern Atlantic. These authors believe that there may be two populations in the western Atlantic or that there may be a single population which is unavailable to the fishery while the fish are migrating between the areas.

In the northwestern Atlantic, 561 blue marlin have been tagged since 1955 and only 4 have been recaptured, all near their respective release points (Mather, 1971). Although these tag returns are inconclusive, it does indicate that meaningful information can be obtained if sufficient numbers of fish are tagged.

For the central Pacific, Royce (1957) stated that north of the equatorial area the seasonal occurrence of blue marlin suggests a northward summer movement followed by a return south in late autumn. Anraku and Yabuta (1959) considered blue marlin in the Pacific to be a single intermingling unit which moves to the southeastern Pacific during the southern summer and returns to the northwestern Pacific during the northern summer. In the southeastern Pacific, Suda and Schaefer (1965) postulated a "strong acti migration" of blue marlin in the vicinity of lat. 20 long. 120°W. Howard and Ueyanagi (1965) indicat that the blue marlin migrates between the North a South Pacific towards the southeast and northwe respectively. Fish taking part in this seasonal migr tion are generally limited in size, 140 to 180 cm; a fi larger fish over 200 cm also take part in the migrati This suggests differential migration by sex, migrati activity being greater in males. Howard and Ueyan consider that migration between the North and Sou Pacific indicates shift of habitat in accordance w spawning and seasonal change of sea conditions.

In the Pacific Ocean, 170 blue marlin have be tagged from 1963 through 1970. There have been recoveries to date.

Sonic tags have already proved useful in studies migration and local movements. A blue marlin w tagged on 14 July in Hawaii 3.1 miles west of Keauh (Yuen, Dizon, and Uchiyama, 1974). The tag was serted at 0935 and the fish was tracked by the NM RV *Charles H. Gilbert* until 0800 the next morning temperature sensitive sonic tag was selected in orc to also obtain information on the depth of the fis During the tracking period, the fish moved to about miles north of the point of release on an erratic cou between the 183- and 549-m (100- and 300-fatho isobaths. Calculated speeds of the fish ranged fr 0.6 to 4.4 knots with an average of 1.6. Swimmi depth varied from the surface to about 73 m (fathoms) but it was mostly in the upper 37 m (abo 20 fathoms).

3.52 Schooling

According to published information, mostly in t sport fishing literature, the blue marlin has nev been observed occurring in schools. Verbal repo from anglers and sport fishing guides and my o observations confirm this. Occasionally, two or thr fish will rise simultaneously to baits trolled from single boat which indicates a small aggregation b not what would be termed a school. Ovchinnik (1970) states that, unlike the sailfish, the blue mar does not form "accumulations" in coastal waters. also reports that, in the open ocean, blue marlin rar gather in schools but are usually found "scatter singly."

Data on composition of blue marlin stocks by si and sex are given by various workers for the Atlan and the Indo-Pacific.

Royce (1957) recorded weight frequencies by mon for 4,712 central Pacific blue marlin but the sex w not recorded. According to Strasburg (1970), Royc data show that the weight distribution was wide January when 45- to 136-kg (100- to 300-lb) fi dominated the catch. By April, heavier fish 136 to 2 kg (300 to 500 lb), presumably females, were mo common. By June, smaller fish, presumably mal

numerous. These small fish dominated the mer catch, reached a peak in September, and slowly declined. By December, conditions were lar to January.

rdman (1968) recorded weight frequencies for 263 es and 65 females taken in waters off Puerto Rico. data show that both males and females reach a of abundance in September.

ume and Joseph (1969b) recorded the length- uency distribution for 3,595 blue marlin from the ern Pacific. The length ranged from 100 to 340 cm to fork) but most of the specimens fell between and 250 cm. The frequency curve was bimodal; dominant mode occurred at about 200 cm.

eyanagi et al. (1970) recorded the length composi- of Atlantic blue marlin by sex, season, and area. ir data show that in the North Atlantic the lominant length for males was about 150 cm (eye rk) during May to October and about 170 cm dur- November to April. The predominant length for ales was about 210 cm during May to October and ut 210 to 220 cm during November to April. In the atorial Atlantic the predominant length for males about 160 to 190 cm throughout the year, and for ales about 230 to 240 cm during May to October about 140 cm from November to April. In the th Atlantic the predominant length for males was ut 240 cm from May to October and about 190 cm h November to April. The predominant length for ales was about 210 to 290 cm from May to October from November to April there were two modes at ut 130 and 250 cm. Ueyanagi et al. (1970) data also v that males were usually more numerous than ales in all areas and seasons. Of the 1,209 imens studied by them 785 were males and 424 ales.

vchinnikov (1970) gave the size composition by th and weight of 80 specimens of blue marlin from Atlantic but the material was not segregated by He states that the most common length ranges n 180 to 220 cm and the most common weight from o 80 kg (110 to 176 lb), but his table shows a mark- node at 190 to 200 cm and at 20 to 30 kg (44 to 66 Obviously, the weights are erroneous in the table. ength and weight frequencies of total longline hes of blue marlin from the equatorial western In- Ocean were presented by Merrett (1971). Males ged from 150 to 200 cm in length (tip of lower jaw rk) and from 43 to 114 kg (95 to 250 lb) in weight. ales ranged from 235 to 270 cm and from 191 to kg (420 to 595 lb).

dditional data on composition of stocks by size sex is given under 3.12 and 3.16. As already in- ted, there is no information on age composition.

3.53 Responses to stimuli

ironmental stimuli: As already discussed under and 3.22, the blue marlin normally occur in waters warmer than 24°C, but it has been found at surface temperatures as high as 30.5°C and as low as 21.7°C.

As to responses to light, Ovchinnikov (1970) stated that according to recent investigations, the epiphysis and contiguous regions of the diencephalon, have a very high light sensitivity in fishes. He also stated that ". . . in all billfishes the glandular structure of the diencephalon, the analogue of the epiphysis, is powerfully developed . . . ." In the blue marlin, as well as in the other istiophorids, the tissues overlying the diencephalon are transluscent and admit light into the brain cavity. The structures involved are similar to the pineal apparatus of tunas described by me (Rivas, 1953). As indicated by Ovchinnikow, it is assumed that the pineal apparatus of the blue marlin reacts to light and may serve for orientation during horizontal and vertical movements.

There is no information available on the responses of the blue marlin to mechanical and chemical stimuli.

Artificial stimuli: In the northern Gulf of Mexico various natural and artificial baits are used in the sport fishery for billfishes (Nakamura and Rivas, 1972). According to these authors, blue marlin prefer striped mullet, *Mugil cephalus*, over ballyhoo, *Hemiramphus brasiliensis*, and bonito strip. These are the three baits most frequently used in the northern Gulf where blue marlin have also been taken on artificial lures (Kona head, rubber squid, etc.). In taking a dead bait or artificial lure trolled at the surface it is not known whether blue marlin are responding to taste, smell, color, size, shape, or action. It is well known that, in the longline fishery, blue marlin are taken on various types of dead bait.

Electrical: No information available.

4 POPULATION

4.1 Structure

4.11 Sex ratio

See 3.16 and 3.52.

4.12 Age composition

No information available.

4.13 Size composition

See 3.12 for size at maturity and 3.52 for length and weight composition.

The largest blue marlin from the Atlantic Ocean officially recorded by the International Game Fish Association weighed 384 kg (845 lb) and measured 399 cm (157 inches) in total length with a girth of 180 cm

(71 inches). It was taken off St. Thomas, Virgin Islands.

According to a personal communication from an angler, a large blue marlin was taken in the northern Gulf of Mexico, off northwest Florida, several years ago. The stated total length was 427 cm (168 inches) and the girth was 203 cm (80 inches). The fish was not weighed because scales were not available but according to the measurements it must have weighed in the neighborhood of 455 kg (1,000 lb).

Cuban commercial fishermen operating off Havana have told me that, on several occasions, they have taken "casteros," blue marlin weighing over 455 kg (1,000 lb). These fish are taken well below the surface with "palangres," a gear similar to the longline but with fewer hooks.

The largest blue marlin from the Pacific Ocean officially recorded by the International Game Fish Association weighed 524 kg (1,153 lb) and measured 447 cm (176 inches) in total length with a girth of 185 cm (73 inches). It was taken off Guam. The former record weighed 505 kg (1,110 lb) and measured 419 cm (165 inches) in total length with a girth of 202 cm (79.5 inches). This fish was taken in the Indian Ocean off Mauritius.

Although not officially accepted as a world record because three anglers participated in the catch, the largest blue marlin known to date was caught in the Pacific Ocean off Waikiki Beach in the Hawaiian Islands. The fish weighed 820 kg (1,805 lb).

The length-weight relationships given in Figures 1 and 2 are based on 58 males and 104 females from the western Atlantic. A separate curve is given for each sex. Each curve was eye-fitted by joining the points of mean length and weight for the upper, middle, a lower thirds of the total range. The length w measured from the tip of the lower jaw to the fork the tail (Rivas, 1956). The curves show that, on t average, males from about 190 to 220 cm in leng weigh less than females in the same length ran Within this length range the mean weight of males 77 kg (169 lb) and that of females 87 kg (191 lb). lengths of less than about 190 cm and more th about 220 cm, the length-weight relationship is abo the same for males and females.

Length-weight data for 7 juveniles, 90 males, and females, mostly from Puerto Rico, were given tabular form by Erdman (1968). Merrett (197 presented length-weight relationships for about specimens of blue marlin from the equatorial weste Indian Ocean. Strasburg (1969) gave a length-weig curve for blue marlin from Hawaii covering a range about 76 to 495 cm (30 to 195 inches; fork length fr tip of bill) and about 750 kg (20 to 1,650 lb).

4.2 Abundance and Density (of Population)

4.21 Average abundance

There is no information available.

4.22 Changes in abundance

See 2.3 and 3.33.

4.23 Average density

There is no information available.



Figure 1.—The length-weight relationship of male blue marlin in the western Atlantic Ocean.

4.24 Changes in density

Jeyanagi et al. (1970) reported on the annual nges in longline catch and fishing effort for the antic Ocean during 1958 through 1966. Until 1962, he early stages of exploitation, the landings of blue rlin increased with the expansion of the fishery. er 1962, however, the catch per unit of effort show- a definite decrease. The relative abundance of blue rlin in 1965 is about one-fourth the level of 1962. e relative abundance of blue marlin began to rease in the Atlantic Ocean after the fishing effort ended over virtually the entire distributional area he species and after the annual landings exceeded out 80,000 fish.

Based on longline fishing data presented by Merrett 71) for the western Indian Ocean, the relative undance of the blue marlin decreased during the iod covered by the study. The catch rate decreased m 0.204 fish per hundred hooks in 1964 to 0.020 fish 1966.

For the blue marlin sport fishery in the northern lf of Mexico, Nakamura and Rivas (1972) used umber-of-fish-raised-per-hour-of-trolling" as an in- x of relative abundance. Their study only covered the 1971 fishing season (May through October) and the relative abundance, by weekly periods, varied from 0.012 to 0.130 fish-per-hour-of-trolling without any marked peaks.

See 2.1 and 2.3 for additional information on seasonal variation of stocks.

4.3 Natality and Recruitment

4.31 Reproduction rates

No information available.

4.32 Factors affecting reproduction

No information available.

4.33 Recruitment

No information available.

4.4 Mortality and Morbidity

4.41 Mortality rates

No information available.



Figure 2.—The length-weight relationship of female blue marlin in the western Atlantic Ocean.

4.42 Factors causing or affecting mortality

No information available.

4.43 Factors affecting morbidity

No information available.

4.44 Relation of morbidity to mortality rates

No information available.

4.5 Dynamics of Population (as a Whole)

There is not enough information to permit construction of mathematical models for prediction of catch characteristics.

4.6 The population in the Community and the Ecosystem

The species composition of the community is that of the pelagic tropical and subtropical habitat. Several species of billfishes and tunas are sympatric and syntopic with the blue marlin and probably compete with it for food and space. Among those attaining a size comparable to that of young and adult blue marlin are the black marlin;, striped marlin; white marlin; longbill spearfish, *Tetrapturus pfluegeri*; shortbill spearfish, *T. angustirostris*; sailfish; bluefin tuna, *Thunnus thynnus*; bigeye tuna, *T. obesus*; and yellowfin tuna, *T. albacares*. In the Atlantic, the blue marlin occurs with white marlin, longbill spearfish, sailfish, bluefin tuna, bigeye tuna, and yellowfin tuna. In the Pacific and Indian oceans, the blue marlin occurs with black marlin, striped marlin, shortbill spearfish, sailfish, bluefin tuna, bigeye tuna, and yellowfin tuna.

Because the blue marlin is a large predator it is considered a climax feeder after reaching the adult stage.

See 2.22 and 3.32 for information on physical features of the biotope and changes in environmental factors.

5 EXPLOITATION

5.1 Fishing Equipment

5.11 Gear

In the commercial fishery the gear in current use is the conventional Japanese longline which is described amply in the literature (Nakamura, 1938; Bullis 1955; Captiva 1955). Off Havana, Cuba blue marlin are taken commercially with palangres, a type of gear similar to the longline but with only a few hooks.

In the sport fishery, blue marlin are taken by conventional rod and reel methods. Natural or artificial baits are trolled at the surface at speeds varying from 4 to 8 knots. Usually two to four lines are fis simultaneously. Echo sounders or fish detectors not used, but bathythermographs are used to loc the depth of the thermocline. Longline gear is usu fished above the thermocline.

In the sport fishery, artificial hookless wobbl lines called "teasers" are trolled close behind boat. Presumably, the action of these teasers attra marlin to the bait. It is also the opinion of m anglers and fishing guides that the size of the boat degree of vibration of the motors are instrumenta attracting the marlin. Nakamura and Rivas (19 analyzed the effect of boat size in the northern Gul Mexico but could not reach any definite conclusio

5.12 Boats

Japanese-type longliners are used in the comm cial fishery with certain modifications in size equipment according to locality. The Cuban palan referred to above is tended from small boats powe by sail and/or motors (inboard or outboard).

5.2 Fishing Areas

5.21 General geographical distribution

The geographic distribution of commercial fish areas corresponds to the geographic distribution of species as discussed in 2.1. Sport fishing areas limited to only a few countries.

In the United States, sportsmen fish for blue ma off southern Texas, Louisiana, Alaban northwestern and southeastern Florida, No Carolina, southern California, and Hawaii. Act sport fishing is also conducted off Puerto Rico and Virgin Islands.

In the Bahama Islands, an active sport fishery blue marlin is conducted off Bimini, Cat Cay, Sal, Grand Bahama, and Walkers Cay. Many b marlin are caught in the Tongue of the Ocean off east coast of Andros, the largest island of the gro

In Mexico, most of the blue marlin sport fishery curs along the Pacific coast. Acapulco, Mazatlán, the southern tip of Baja California are the most act areas; however, blue marlin are not caught in la numbers in these areas.

In Central and South America most of the sp fishing for blue marlin takes place along the Pac coast. There is an active sport fishery off Piñas B Panama and another off Ecuador. Sport fishing blue marlin in Venezuela is conducted mostly off Guaira.

In the central Pacific, blue marlin are commo taken around the Hawaiian Islands and Tahiti.

In Africa, blue marlin have been caught sportsmen off Senegal and the Ivory Coast. A more tive sport fishery, however, is conducted along coast of South Africa.

In the Indian Ocean, blue marlin are taken

glers fishing off Mauritius. Cairns, on the east coast Australia, is rapidly becoming a very active sport ning center for blue marlin as well as other species marlin. An active sport fishery for blue marlin has sted off northern New Zealand for several years.

5.22 Geographic ranges

According to data published by Howard and yanagi (1965) and Ueyanagi et al. (1970) the com-rcial longline fishery for blue marlin takes place in high seas as well as near the coast. The distance m the coast of commercial operations is limited by eanographic conditions (temperature, depth, etc.) ecting the occurrence of the fish or by restrictions posed by the various countries. In the sport fishery, range of the fishery is limited by the range of the ats used by the anglers as well as oceanographic nditions. For example, off Bimini the habitat orable to the blue marlin is found only 3 or 4 miles m shore. Off Louisiana and northwest Florida, wever, blue marlin are usually found not less than miles from shore.

The areas of greatest abundance have already been cussed in 2.1 and 2.3.

Commercial longline fishing for blue marlin in the antic Ocean developed rapidly from 1958 to 1962 eyanagi et al., 1970). After that year commercial ning operations covered practically the entire dis-butional area of the blue marlin in the Atlantic.

The sport fishery for blue marlin has grown steadily ce its inception early in this century; growth has en particularly rapid since the end of World War II. me of the sport fishing areas developed as a result of mmercial or exploratory fishing operations. Off uth Pass, La., the sport fishery for marlins and as started in the midfifties following longline ex-ratory work by the Bureau of Commercial Fisheries w National Marine Fisheries Service) (Bullis, 55; Captiva, 1955).

5.23 Depth ranges

No information available.

5.24 Conditions of the grounds

No information available.

5.3 Fishing Seasons

5.31 General pattern of seasons

The fishing seasons vary according to the seasonal vements of the fish as discussed in 2.1 and 3.51.

5.32 Dates of beginning, peak, and end of season

In the commercial fishery the longline fleets can low the seasonal movements of the fish; however, this is not true for the sport fishery. For example, in the northern Gulf of Mexico the sport fishing season begins in April or May, reaches a peak in July and August, and ends in October or November (Nakamura and Rivas, 1972). On the other hand, in tropical areas such as Puerto Rico and Jamaica, sport fishing for the blue marlin is conducted throughout the year.

5.33 Variation in date or duration of season

As discussed above in 5.32, the off-season for the sport fishery in the northern Gulf of Mexico is from November to April. During that period adverse weather conditions prevent sport fishing boats from venturing out. Furthermore, low water temperatures render the habitat unsuitable to the blue marlin. The same applies to the sport fishery in areas with a similar climate.

5.4 Fishing Operations and Results

5.41 Effort and intensity

In the longline fishery the unit of effort is the number-of-fish-per-hundred-hooks. In the sport fishery the same unit of effort may be used, but Nakamura and Rivas (1972) have suggested that number-of-fish-raised-per-hour-of-trolling may be a better index of relative abundance.

Ueyanagi et al. (1970) have given the monthly change of the mean hook-rate by area for the blue marlin in the Atlantic Ocean during 1956 through 1965.

In the commercial longline fishery the causes of variation in fishing effort are the result of changes in economic factors and fluctuations in stock abundance. In the sport fishery, weekends, national holidays, and traditional summer vacations are causes of variation in fishing intensity. In general, fishing intensity is highest on weekends. In certain areas, such as the northern Gulf of Mexico, sport fishing intensity drops markedly during the Labor Day weekend. Presumably, anglers select that particular holiday to spend time with their families before the children go back to school. However, in areas such as southern California this is the period when the peak fishing effort for billfish occurs.

5.42 Selectivity

The longline is selective for larger fish and in the sport fishery the baits are rigged to catch the larger fish. Anglers and fishing guides believe that the larger the bait and the hook, within limits, the larger the blue marlin that the bait will attract. This is the rule only in certain areas. Off South Pass, La., anglers troll mostly large baits with a corresponding large hook. They specialize almost exclusively in blue marlin fishing and usually catch more of them (number-of-fish-per-hour-of-trolling) as compared to northwest

Florida where smaller baits are trolled (Nakamura and Rivas, 1972). At present there is no way of telling whether the better catches off South Pass result from larger baits, better fishermen, or greater abundance of blue marlin.

5.43 Catches

Ueyanagi et al. (1970) have given the annual longline catches (1958-66) of blue marlin for the Atlantic Ocean on the basis of numbers of fish caught rather than total weights. Their graph shows the estimated catches of blue marlin by year is as follows:

Year	*Estimated catch (number)*
1958	8,900
1959	21,500
1960	26,100
1961	41,200
1962	110,400
1963	95,300
1964	82,900
1965	44,000
1966	20,200

Annual longline catches (1962-70) of blue marlin were recently published by Gottschalk (1972) for the entire Atlantic and the eastern Pacific. In the Atlantic, the catches are expressed in the number of fish caught:

Year	*Fish caught (number)*
1962	111,000
1963	96,000
1964	84,000
1965	45,000
1966	22,000
1967	11,000
1968	9,000
1969	14,000
1970	11,000

In the eastern Pacific, there were:

Year	*Fish caught (number)*
1962	37,000
1963	76,000
1964	46,000
1965	26,000
1966	22,000
1967	22,000
1968	28,000
1969	34,000
1970	20,000

Data collected from the sport fishery of the northern Gulf of Mexico during the 1971 season (May to October) show that 84 blue marlin weighing a total of 8,995 kg (19,788 lb) were caught. Of these, 20 were, males weighing 28 to 127 kg (62 to 280 lb) and 64 were females weighing 21 to 224 kg (46 to 492 lb). Average weight of males was 71 kg (156 lb) and that of females was 118 kg (260 lb). The average weight of all 84 fish, males and females combined, was 107 kg (236 lb). T fish were caught out of South Pass, La., and P sacola, Destin, and Panama City, Fla.

6 PROTECTION AND MANAGEMENT

6.1 Regulatory (Legislative) Measures

6.11 Limitation or reduction of total cat

At present, there are no limitations or reductions any kind applying to the blue marlin fishery. A stea decline in the catch, however, since 1962 (see 5 above) has called attention to the need for regulat measures. Analysis of the annual catches of b marlin in the Atlantic from 1958 to 1962 shows t pattern of a virgin fishery when first opened to ploitation. Instead of leveling off, however, the cat has steadily declined since 1962, despite increased fort, from 111,000 fish to 11,000 in 1970. I have no formation on whether or not the average size of t blue marlin has also declined.

6.12 Protection of portions of population

There are no closed areas or seasons and there a no limitations on gear. There are, howeve limitations on the use of blue marlin for food in t United States according to the degree of mercury co tamination of the flesh. Shomura and Craig (197 reported on the total mercury levels found in t white muscle tissue of 37 blue marlin caught Hawaiian waters. Thirty of the 37 fish had mercu levels that exceeded the 0.5 ppm. level used as guideline by the U.S. Food and Drug Adminstrati as fit for human consumption. The mercury lev ranged from 0.7 to 7.86 ppm. in fish weighing betwe 43.5 and 410.9 kg (96 and 906 lb, respectively).

7 POND FISH CULTURE

Not applicable.

LITERATURE CITED

ANRAKU, N., and Y. YABUTA.
1959. Seasonal migration of black marlin. [In Jap., En synop.] Rep. Nankai Reg. Fish. Res. Lab. 10:63-71.

BAKER, A. N.
1966. Food of marlins from New Zealand waters. Cop 1966:818-822.

BARTLETT, M. R., and R. L. HAEDRICH.
1968. Neuston nets and South Atlantic larval blue mar (*Makaira nigricans*). Copeia 1968:496-474.

BRIGGS, J. C.
1960. Fishes of worldwide (circumtropical) distributi Copeia 1960:171-180.

LLIS, H. R., JR.
1955. Preliminary report on exploratory long-line fishing for tuna in the Gulf of Mexico and the Caribbean Sea. Part I - Exploratory fishing by the *Oregon*. Commer. Fish. Rev. 17(10):1-15.
LDWELL, D. K.
1962. Postlarvae of the blue marlin, *Makaira nigricans*, from off Jamaica. Los Ang. Cty. Mus., Contrib. Sci. 53, 11 p.
PTIVA, F. J.
1955. Preliminary report on exploratory long-line fishing for tuna in the Gulf of Mexico and the Caribbean Sea. Part II - Long-line gear used in yellowfin tuna exploration. Commer. Fish. Rev. 17(10):16-20.
ESSEY, R. F., and E. A. LACHNER.
1970. The parasitic copepod diet and life history of diskfishes (Echeneidae). Copeia 1970:310-318.
VIER, G. L. C. F. D.
1832. Sur le poisson appele *Machaera*. Nouv. Annu. Mus. Nat. Hist. Natur. (Paris) 1:43-49.
VIER, G., and A. VALENCIENNES.
1831. Histoire naturelle des poissons, Vol 8. Chez F. G. Levrault, Paris, 509 p.
BUEN, F.
1950. Contribucion a la ictiologia. 3. La familia Istiophoridae y descripción de una especie uruguaya (*Makaira perezi* de Buen). Publ. Cient. Serv. Oceanogr., Montevideo 5:163-178.
SYLVA, D. P.
1958. Juvenile blue marlin, *Makaira ampla* (Poey), from Miami, Florida, and West End, Bahamas. Bull. Am. Mus. Nat. Hist. 114:412-415.
1963. Preliminary report on the blue marlin sport fishery of Port Antonio, Jamaica. Univ. Miami, Inst. Mar. Sci., 10 p.
DMAN, D. S.
1957. Marlin fishing in Puerto Rican waters. Proc. Second Int. Game Fish Conf., 13 p. Univ. Miami Mar. Lab., Miami.
1962. The sport fishery for blue marlin off Puerto Rico. Trans. Am. Fish. Soc. 91:225-227.
1968. Spawning cycle, sex ratio, and weights of blue marlin off Puerto Rico and the Virgin Islands. Trans. Am. Fish. Soc. 97:131-137.
CHMEYER, W. N., and H. R. BULLIS, JR.
1968. Four advanced larval specimens of the blue marlin, *Makaira nigricans*, from the western Atlantic Ocean. Copeia 1968:414-417.
HRINGER, J. W.
1956. Observations on the development of the Atlantic sailfish *Istiophorus americanus* (Cuvier), with notes on an unidentified species of istiophorid. U.S. Fish Wildl. Serv., Fish. Bull. 57:139-171.
TTSCHALK, J. S.
1972. Longlines and billfish. Mar. Fish. Rev. 34(11-12): 20-25.
AY, J. E.
1838. Description of a new species of *Tetrapturus* from the Cape of Good Hope. Ann. Mag. Nat. Hist., Ser. 1, 1:313.
ASAKA, K., and H. NAKAMURA.
1947. On the Formosan spearfishes. Bull. Oceanogr. Inst. Taiwan 3:9-24.
WARD, J. K., and S. UEYANAGI.
1965. Distribution and relative abundance of billfishes (*Istiophoridae*) of the Pacific Ocean. Univ. Miami Inst. Mar. Sci., Stud. Trop. Oceanogr. 2, 134 p.
IVERSEN, E. S., and E. E. HOVEN.
1958. Some trematodes of fishes from the central equatorial Pacific. Pac. Sci. 12:131-134.
IVERSEN, R. T. B., and R. B. KELLEY.
1974. Occurrence, morphology and parasitism of gastric ulcers in blue marlin, *Makaira nigricans*, and black marlin, *Makaira indica*, from Hawaii. *In* R. S. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972, part 2. Review and Contributed Papers, p. 149-153. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-675.
JONES, S., and E. G. SILAS.
1964. A systematic review of the scombroid fishes of India. Proc. Symp. Scombroid Fishes, Part I, p. 1-105. Mar. Biol. Assoc. India, Mandapam Camp.
JORDAN, D. S., and J. O. SNYDER.
1901. Descriptions of nine new species of fishes contained in museums of Japan. J. Coll. Sci., Imp. Univ. Tokyo 15:301-311.
KORATHA, K. J.
1955. Studies on the monogenetic trematodes of the Texas coast. I. Results of a survey of marine fishes at Port Aransas, with a review of Monogenea reported from the Gulf of Mexico and notes on euryhalinity, host-specificity, and relationship of the Remora and the Cobia. Univ. Tex., Publ. Inst. Mar. Sci. 4:233-249.
KRUMHOLZ, L. A.
1958. Relative weights of some viscera in the Atlantic marlins. Bull. Am. Mus. Nat. Hist. 114:402-405.
KRUMHOLZ, L. A., and D. P. DE SYLVA
1958. Some foods of marlins near Bimini, Bahamas. Bull. Am. Mus. Nat. Hist. 114:406-411.
KUME, S., and J. JOSEPH
1969a. The Japanese longline fishery for tunas and billfishes in the eastern Pacific Ocean east of 130°W, 1964-1966. [In Engl. and Span.] Bull. Inter-Am. Trop. Tuna Comm. 13:277-418.
1969b. Size composition and sexual maturity of billfish caught by the Japanese longline fishery in the Pacific Ocean east of 130°W. [In Engl., Jap. summ.] Bull. Far Seas Fish. Res. Lab. (Shimizu) 2:115-162.
LACÉPÈDE, B. G. E.
1802. Histoire naturelle des poissons. 4:689-697.
MANTER, H. W.
1954. Trematoda of the Gulf of Mexico. U.S. Fish Wildl. Serv., Fish. Bull. 55:335-350.
MATHER, F. J., III.
1971. Summary of results of billfish tagging in the Atlantic 1954-1970. Coop. Game Fish Tagging Program, Woods Hole Oceanogr. Inst.
MATHER, F. J., III, A. C. JONES, and G. L. BEARDSLEY, JR.
1972. Migration and distribution of white marlin and blue marlin in the Atlantic Ocean. Fish. Bull., U.S. 70:283-298.
MERRETT, N. R.
1971. Aspects of the biology of billfish (Istiophoridae) from the equatorial western Indian Ocean. J. Zool. (Lond.) 163:351-395.
MOORE, H. L.
1950. The occurrence of a black marlin, *Tetrapterus mazara*, without spear. Pac. Sci. 4:164.
MORROW, J. E., JR.
1959a. Distribution of the blue marlin and black marlin in the Indo-Pacific. Bull. Mar. Sci. Gulf Caribb. 9:321-323.

1959b. On *Makaira nigricans* of Lacépède. Postilla 39, 12 p.
1964. Marlins, sailfish and spearfish in the Indian Ocean. Proc. Symp. Scombroid Fishes, Part 1, p. 429-440. Mar. Biol. Assoc. India, Mandapam Camp.

MOWBRAY, L. L.
1931. A Bermudian spearfish, family Istiophoridae. *Makaira bermudae* n. sp. Mowbray. Fauna Bermudensis 1 [p. 7-8].

NAKAMURA, E. L., and L. R. RIVAS.
1972. Big game fishing in the northeastern Gulf of Mexico during 1971. Natl. Mar. Fish. Serv., Panama City, Fla., 20 p.

NAKAMURA, H.
1938. Report of an investigation of the spearfishes of Formosan waters. [In Jap.] Reports of the Taiwan Government-General Fishery Experiment Station, No. 10. (Translated by W. G. Van Campen, 1955, U.S. Dep. Int., Fish Wildl. Serv., Spec. Sci. Rep. Fish. 153, 46 p.)
1942. [On the ecology of the istiophorid fishes of Taiwan waters.] [In Jap.] Sci. Fish. Assoc., Fac. Agr. Tokyo Imp. Univ., Suisan Gakkwai Ho. 9:45-51.

NAKAMURA, I., I. IWAI, and K. MATSUBARA.
1968. A review of the sailfish, spearfish, marlin and swordfish of the world. [In Jap.] Kyoto Univ., Misaki Mar. Biol. Inst., Spec. Rep. 4, 95 p.

NARDO, G. D.
1833. De Skeponopodo novo piscium genere et de Guebucu marcgravii specii, illi cognata. Isis (Oken) 26:415-419.

OVCHINNIKOV, V. V.
1970. Mech-ryba i parunikovye (Atlanticheskii okean. Ekologiya i funktsional'naya morfologiya) (Swordfishes and billfishes in the Atlantic Ocean. Ecology and functional morphology). Atl. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr., Kaliningrad, 106 p. (Translated by H. Mills, Israel Program Sci. Transl., 1971, 77 p.; available U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA, as TT 71-50011.)

POEY, F.
1860. Poissons de Cuba, espèces nouvelles. Mem. Hist. Nat. Isla de Cuba 2:115-356.

RAFINESQUE, C. S.
1810. Caratteri di alcuni nuovi generi e nuove specie di animali e piante della Sicilia, con varie osservazioni sopri i medesimi. Palermo, 105 p.

RIVAS, L. R.
1953. The pineal apparatus of tunas and related scombrid fishes as a possible light receptor controlling phototactic movements. Bull. Mar. Sci. Gulf Caribb. 3:168-180.
1956. The occurrence and taxonomic relationships of the blue marlin (*Makaira ampla* Poey) in the Pacific Ocean. Bull. Mar. Sci. Gulf Caribb. 6:59-74.

ROBINS, C. R., and D. P. DE SYLVA.
1960. Description and relationship of the longbill spearfish, *Tetrapturus belone*, based on Western North Atlantic specimens. Bull. Mar. Sci. Gulf Caribb. 10:383-413.

ROYCE, W. F.
1957. Observations on the spearfishes of the central Pacific. U.S. Fish Wildl. Serv., Fish. Bull. 57:497-554.

SHOMURA, R. S., and W. L. CRAIG.
1974. Mercury in several species of billfishes taken Hawaii and southern California. *In* R. S. Shomura F. Williams (editors), Proceedings of the Internatic Billfish Symposium, Kailua-Kona, Hawaii, 9-12 Aug 1972, Part 2. Review and Contributed Papers, p. 160- U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-6

SMITH, J. L. B.
1956. Pugnacity of marlins and swordfish. Nature (Lor 178:1065.

SQUIRE, J. L., JR.
1962. Marlin and swordfish in oceanic waters of the west North Atlantic. Copeia 1962:216-219.

STRASBURG, D. W.
1969. Billfishes of the central Pacific Ocean. U.S. F Wildl. Serv., Circ. 311, 11 p.
1970. A report on the billfishes of the central Pacific Oce Bull. Mar. Sci. 20:575-604.

SUDA, A., and M. B. SCHAEFER.
1965. General review of the Japanese tuna long-line fish in the eastern tropical Pacific Ocean 1956-1962. B Inter-Am. Trop. Tuna Comm. 9:307-462.

TINSLEY, R.
1964. The sailfish—swashbuckler of the open seas. Un Fla. Press, Gainesville, 216 p.

UEYANAGI, S.
1963. Methods for identification and discrimination of t larvae of five istiophorid species distributing in the Ind Pacific. [In Jap., Engl. synop.] Rep. Nankai Reg. Fis Res. Lab. 17:137-150.
1964. Description and distribution of larvae of five ist phorid species in the Indo-Pacific. Proc. Symp. Sco broid Fishes, Part 1, p. 499-528. Mar. Biol. Assoc. Ind Mandapam Camp.

UEYANAGI, S., S. KIKAWA, M. UTO, and Y. NISHIKAWA
1970. Distribution, spawning, and relative abundance billfishes in the Atlantic Ocean. [In Jap., Engl. synop Bull. Far Seas Fish. Res. Lab. (Shimizu) 3:15-55.

UEYANAGI, S., and H. YABE.
1959. Larvae of the black marlin (*Eumakaira nigra* Nak mura). [In Jap., Engl. summ.] Rep. Nankai Reg. Fis Res. Lab. 10:151-169. (English transl. by W. G. V Campen, 1961, 20 p.)

VIGUERAS, I. P.
1935. *Tristomum poeyi*, n. sp. (Tremotoda), parasito *Makaira ampla* Poey (Pisces). Mem. Soc. Cubana His Nat. 9:43-44.

WHITLEY, G. P.
1931. New names for Australian fishes. Aust. Zool. 6(310-334.
1954. More new fish names and records. Aust. Zo 12(1):57-62.

YUEN, H. S. H., A. E. DIZON, and J. H. UCHIYAMA.
1974. Notes on the tracking of the Pacific blue marli *Makaira nigricans*. *In* R. S. Shomura and F. Willia (editors), Proceedings of the International Billfish Sy posium, Kailua-Kona, Hawaii, 9-12 August 1972, Part Review and Contributed Papers. 265-268. U.S. De Commer., NOAA Tech. Rep. NMFS SSRF-675.

Synopsis of the Biology of the Black Marlin, *Makaira indica* (Cuvier), 1831

IZUMI NAKAMURA[1]

IDENTITY

1.1 Nomenclature

1.11 Valid name

Makaira indica (Cuvier) 1831.

Original combination: *Tetrapturus indicus* Cuvier 1831. *In* Cuvier and Valenciennes, Hist. Nat. Poiss. Paris 8:286-287. (Original locality: Sumatra.)

1.12 Objective synonymy

Tetrapturus indicus Cuvier 1831. *In* Cuvier and Valenciennes, Hist. Nat. Poiss. Paris. 8:286-287.
Makaira indica Jordan and Evermann 1926. Calif. Acad. Sci., Occas. Pap. 12:67.
Makaira indicus (partim) Deraniyagala 1933. Spolia Zeylan. 18:55-56.
Istiompax indicus Morrow 1959. Copeia 1959:347-349.
Makaira (Istiompax) indica Robins and de Sylva 1960. Bull. Mar. Sci. Gulf Caribb. 10:406.

1.2 Taxonomy

1.21 Affinities

Suprageneric

Phylum Chordata
Subphylum Vertebrata
Superclass Gnathostomata
Class Osteichthyes
Subclass Actinopterygii
Order Perciformes
Suborder Xiphioidei
Family Istiophoridae

Generic

Genus *Makaira* Lacépède.

Makaira Lacépède 1803, Hist. Nat. Poiss. 4:688. (Type-species: *Makaira nigricans* Lacépède 1803.)

This genus is distinguished by the height of first dorsal fin being less than body depth, short pelvic fin with poorly developed membrane, body slightly compressed laterally, posterodorsal part of the head remarkably convex, skull stout and broad, neural and haemal spines showing a high trapezoid when viewed laterally, lateral apophysis of vertebrae well developed, vertebral count of 11 + 13 = 24 (Nakamura, Iwai, and Matsubara, 1968).

Specific

Types of this species might be unknown. This species was described from a drawing of a specimen 274 cm (9 feet) long and weighing 440 kg (200 pounds), from Sumatra.

The substance of the original description of this species is as follows:

> Spear long, five times in body length including caudal fin, dorsal lobe 1⅓ to 1 in depth of body, ventrals one-half of depth; dorsal fin low posteriorly, the 4th or 5th middle spines highest and nearly equal to body depth.

This species is distinguished clearly from other members of *Makaira* by its rigid pectoral fin and a second dorsal fin situated a little forward of the insertion of the second anal fin (Nakamura et al., 1968).

Subjective synonymy

Tetrapturus australis Macleay 1854. Illus. Sydney News 1(23):179.
Histiophorus brevirostris Playfair 1866. *In* Playfair and Günther, The fishes of Zanzibar, p. 53 and 145, London.
Histiophorus gladius Ramsay 1881 (*non* Bloch 1793). Proc. Linn. Soc. N.S.W. 5:295-297.
Makaira marlina Jordan and Hill 1926. *In* Jordan and Evermann. Calif. Acad. Sci., Occas. Pap. 12:59-60, pl. 17.
Istiompax australis Whitley 1931. Rec. Aust. Mus. 18(4):147-150.
Makaira nigricans tahitiensis Nichols and LaMonte 1935. Am. Mus. Novit. 807:1-2.
Makaira nigricans marlina Nichols and LaMonte 1935. Nat. Hist., N.Y. 36:328.
Makaira ampla marlina Nichols and LaMonte 1941. Ichthyol. Contrib. Int. Game Fish Assoc. 1(1):8, fig. 1.

[1] Fisheries Research Station, Kyoto University, Maizuru, Kyoto Japan.

Makaira ampla tahitiensis Nichols and LaMonte 1941. Ichthyol. Contrib. Int. Game Fish Assoc. 1(1):8, fig. 3.
Tetrapturus brevirostris Smith 1946. Ann. Mag. Nat. Hist. 11:793-807.
Marlina marlina Hirasaka and Nakamura 1947. Bull. Oceanogr. Inst. Taiwan 3:15, pl. 3, fig. 1.
Istiompax dombraini Whitley 1954. Aust. Zool. 12 (1):60.
Makaira mazara (partim) LaMonte 1955 (*non* Jordan and Snyder 1901). Bull. Am. Mus. Nat. Hist. 107: 336.
Makaira mazara tahitiensis LaMonte 1955. Bull. Am. Mus. Nat. Hist. 107:342.
Makaira xantholineata Deraniyagala 1956. Spolia Zeylan. 28(1):23-24.
Istiompax marlina Royce 1957. U.S. Fish Wildl. Serv., Fish. Bull. 57:524-528, fig. 2d, 3a.
Istiompax brevirostris Morrow 1958. Bull. Mar. Sci. Gulf Caribb. 8:358.
Makaira australis Marshall 1964. Fishes of the Great Barrier Reef and coastal waters of Queensland, p. 349-350, pl. 47, Narberth.

The following key to the species of *Makaira* is based on Nakamura et al. (1968):

a. Pectoral fin folds back against the side of the body. Lateral line not single line.
b. Lateral line system simple loops in shape - - - - *Makaira mazara* Jordan and Snyder, Indo-Pacific blue marlin.
bb. Lateral line system reticulate - - - - *Makaira nigricans* Lacépède, Atlantic blue marlin.
aa. Pectoral fin extends stiffly away from the body, cannot be folded back against the side of the body. Lateral line obscure but single line - - - - *Makaira indica* (Cuvier), black marlin, (Fig. 1).

1.22 Taxonomic status

This species was placed in genus *Istiompax* by several authors, e.g., Whitley, 1931a, 1931b, 1948, 1955; Royce, 1957; Morrow, 1959a, 1959b, 1959c, but many other authors have placed it in genus *Makaira*. Species concept of *M. indica* is still one of morphospecies.

1.23 Subspecies

Some authors proposed a subspecies status for t species, e.g., Nichols and LaMonte, 1935a, 193 LaMonte and Marcy, 1941; Rosa, 1950; Morrow, 19 1957, but this has not been generally accepted due the lack of adequate evidence. Further studies geographical morphological forms or subspecies a highly needed.

1.24 Standard common names and v nacular names

In current scientific literature "black marlin" is variably used as the common name for *Makaira dica.*[2] The following vernacular names are used fishermen:

Location	*Common name*
Sri Lanka (general)	Marlin, Ahin Koppara
Sri Lanka (Tamil)	Kopparan, Kopparaikulla
Sri Lanka (Sinhala)	Kapparava, Makara, Sapparava
Indonesia	Joo Hoo
U.S.A.	Pacific black marlin, giant black marl silver marlin
New Zealand, Australia, New Guinea	Black marlin
Formosa	Kȳau-shit-á
Tokyo, Japan	Shirokawa
Kochi, Japan	Genba
Okinawa, Japan	Shiruachi
Various parts of Japan	Shirokajiki, Shirokawakajiki

[2]"Black marlin" as the common name for *Makaira indica* was n universally accepted until the mid-1960's. Prior to this agreeme some authors, principally scientists from Japan, called this spec "white marlin" (see for example Ueyanagi, 1964). To add to t confusion, much of the earlier Japanese literature used "bla marlin" to refer to the species currently known as "blue marlin *Makaira nigricans*. The fact that the scientific names have go through a similar period of changes has not helped t nomenclature situation any.

Figure 1.—Black marlin, *Makaira indica* (Cuvier).

1.3 Morphology

1.31 External and internal morphology

escription of adult (after Nakamura et al. 1968):

First dorsal fin rays 38-42, second dorsal fin rays 6- first anal fin rays 13-14, second dorsal fin rays 6-7, ectoral fin rays 19-20, pelvic fin 1, 2.

Body elongated (greatest depth of body[3] about 4.8-2 in body length), slightly compressed laterally idth of body at origin of pectorals about 11.0-13.4 in dy length). Snout long (bill length about 0.95 in ead length) with a cross section nearly circular. cales distributed densely on body surface; scale ends inted and long.

Rasplike teeth on jaws and palatine. Lateral line scure but consists of a single line. Head large (head ngth about 4.0-4.2 in body length). Eye moderate in ze. Posterodorsal part of the head between preor-tal and origin of the first dorsal fin strongly convex. audal fin large and strongly falcate. A pair of keels tuated a little posteriorly on caudal peduncle. Pec-ral fin situated low on body, long (length of pectoral n about 1.1-1.3 in head) and its tip pointed. Pectoral n extends stiffly away from the body, cannot be fold-l back against the side of the body. First dorsal fin arts above supraposterior margin of the opercle and nds just before the origin of the second dorsal fin; nterior portion lower than the body height and adually decreasing in height posteriorly. The tip of e anterior part of the first dorsal fin pointed. First nal fin large and triangular in shape, its tip pointed. econd dorsal fin nearly equal to second anal fin in ze and shape, origin opposite each other, or the rmer a little farther forward than the latter. Pelvic n shorter than pectoral fin.

Membrane of first dorsal fin dark blue, other fin embranes brownish black. No marks or blotches on e body. Dorsal part of the body blackish dark blue, ntral part of body silvery white. Color of the body des after death, so this species is called "Shiroka-i" (= white marlin) in Japan. While alive, the color body is tinged with black, and hence this species is lled "black marlin" in English-speaking countries.

Nasal rosette composed of about 50 laminae. Gonad mmetrical. Anus situated just anterior of the origin first anal fin. Skull broad and strong; the preorbital rt elongated and its postorbital part shortened. entral part of the vomer and anteroventral part of e parasphenoid flat. Temporal ridge parallel with erotic ridge. Haemal and neural spines of the cen-al part of vertebrae high trapezoid in shape. Lateral ophysis well developed but not so much as in *akaira mazara* and *M. nigricans*. Vertebrae 11 + 13 24.

[3]Measurements were carried out by the method of Rivas (1956).

2. DISTRIBUTION

2.1 Total Area

Black marlin are distributed widely in the Indian and Pacific oceans; very few catches have been recorded from the Atlantic Ocean (Fig. 2). Only stray black marlin seem to invade the Atlantic Ocean by way of the Cape of Good Hope; possibly breeding stocks do not exist in the Atlantic Ocean. Good commercial fishing grounds exist in the East China Sea, around Formosa, in the waters off northwest Australia, the Arafura Sea, the Sulu Sea, the Celebes Sea, and the Coral Sea. Good sport fishing grounds for black marlin are found in Piñas Bay of Panama and off Ecuador (Fig. 3).

Sea surface temperatures at or near the areas where the black marlin are caught range from about 15° to 30°C (Howard and Ueyanagi, 1965).

2.2 Differential Distribution

2.21 Spawn, larvae, and juveniles

Very few larvae have been obtained from the tropical western north Pacific, tropical Indian Ocean, and tropical water off northwestern Australia and the Coral Sea (Ueyanagi, 1964). It has been surmised that the spawning grounds are in the northwestern part of the Coral Sea (Ueyanagi, 1960; Ueyanagi and Yabe, 1960) and in the vicinity of Hainan Island, South China Sea (Nakamura, 1941, 1942). But spawn is not identified.

2.22 Adults

This species is distributed widely in warmer parts of the Pacific Ocean and the Indian Ocean and densely in coastal waters. In tropical open seas areas, distribution is very scattered but continuous, whereas in temperate open seas, there is almost no occurrence of this species (Howard and Ueyanagi, 1965). In tropical open seas areas, there is a tendency for this species to occur in the vicinity of islands (Nakamura, 1953; Royce, 1957; Koto, Furukawa, and Kodama, 1959; Ueyanagi, 1963).

2.3 Determinants of Distribution Changes

In the East China Sea, black marlin fishing grounds, which are shallower than those of the other billfishes, are found in the areas where the Kuroshio and Tsushima currents are mixed with the waters of the Yellow Sea (Koto et al., 1959). Optimum temperatures for harpoon fishing of black marlin westward of Uotsurijima in the East China Sea, are 23° to 25°C between October and April (Morita, 1952).

2.4 Hybridization

No data.



Figure 2.—Distribution and relative abundance of black marlin based on the data from the Japanese longline fisheries in 1969 (chiefly), 1968, 1967, and 1964.

BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

The sexes of the black marlin are separate. The males and females cannot be distinguished by external characters. In the waters around Formosa, the sex ratio is 53/414 male throughout a size range of 20 to 00 kg in body weight (Nakamura, 1944a). Females become larger than males (Nakamura, 1944b).

3.12 Maturity

Age at which sexual maturity is reached is not known.

3.13 Mating

The mating of the black marlin has not been observed.

3.14 Fertilization

External.

3.15 Gonad

Black marlin are densely distributed in the northwestern part of the Coral Sea between October and December. Almost all fishes caught in this area have well-developed gonads (Table 1); thus these fishes are believed to be from spawning schools in this area. The skewed sex ratios may also be indicative of this possibility (Ueyanagi, 1960).

Merrett (1970) studied histological gonad development in billfishes including this species from the Indian Ocean.

3.16 Spawning

There is very little information relating to spawning grounds and spawning seasons of the black marlin. Nakamura (1941, 1942, 1944b) surmised that spawning occurs in the vicinity of Hainan Island and the South China Sea in May or June. Ueyanagi (1960) also presumed spawning to occur in the northwestern part of the Coral Sea between October and December.

The sex ratio of the black marlin varies with area and season (Table 2). Variations in sex ratios seem to be related to spawning (Nakamura, 1942).

3.17 Spawn

The egg of the black marlin seems to be pelagic and nonadhesive, but details are not known.

3.2 Preadult Phase

3.21 Embryonic phase

No data.

3.22 Larvae phase

The embryological development of the black marlin is not known. Some larvae have been obtained by research vessels. Morphological features of these larvae have been described by Ueyanagi (1964).

The back of the larvae is slightly concave where the



gure 3.—Distribution and fisheries of black marlin. Shaded areas show good fishing grounds for commercial longline fisheries. orizontal barred area shows harpoon fishing ground. Dotted areas show sport fishing grounds. Solid line indicates northern and uthern limits of distribution of black marlin based on data from the longline catches. Dotted line shows movement of black marlin om the Indian Ocean into the Atlantic Ocean.

Table 1.—Data on the gonad weight of *Marlina marlina* (Jordan and Hill) in the northwestern area of the Coral during November 1956 (*No. 1 Satsuma-maru*) (Ueyanagi, 1960).

Body length (cm)	Testis weight (g)					Ovary weight (g)					
	less than 500	500-1,000	1,000-2,000	2,000-4,000	more than 4,000	less than 500	500-1,000	1,000-2,000	2,000-5,000	5,000-10,000	more than 10,00
131-140	1			1							
141-160	5	18	25								
161-180	2	38	84	22					1		
181-200		11	66	58	6						
201-220	1	4	13	95	30					2	
221-240				24	31				2	5	
241-260				2	3				1	12	9
261-280				1					1	7	11
281-300									3	2	8
301-320										3	7

head joins the trunk. This species can be identified also by the short snout, specific shape of the pectoral fin and vertebral formulae (11 + 13). Because of similar short snouts, the head profile of the black marlin larvae resembles that of blue marlin; however, the two species can be discriminated since the blue marlin has eye orbits with protruding anterior edges. In small specimens under 6 mm the snout is also somewhat longer than in the blue marlin. The shape of the pectoral fin is unique in larvae over 3.5 mm. In specimens 3.5 mm in length, the pectoral fin stands out rigidly from the lateral side of the body and ca not be folded against the body without breaking t joint. In specimens of about 5 mm not only does th characteristic become more distinct, but also the fin turned slightly counterclockwise, because of the d tortion of the base of the fin. Even in specimens of mm in length, the transition of the fin to the adu condition is not yet complete, but it has progressed f enough to show a relationship with the adult form. addition, the shape of the dorsal fin seems to different in this species (Fig. 4).

Table 2.—The sex ratio of the black marlin in various areas.

Date	Localities	Female		Male		References
		No.	%	No.	%	
Dec. 1937	Suao	44	85	8	15	Nakamura, 1942
July 1939	Hainan Island	4	25	12	75	Nakamura, 1942
Dec. 1939	Suao	44	92	5	8	Nakamura, 1941
Mar. 1940	Kaohsiung	52	87	8	13	Nakamura, 1942
Nov. 1940	Suao	8	80	2	20	Nakamura, 1942
Jan. 1941	Suao	16	89	2	11	Nakamura, 1942
May 1941	Kaohsiung	11	71	3	29	Nakamura, 1942
Jan. 1942	Suao	15	94	1	6	Koto and Kodama, 1962
Nov. 1942	Suao	22	85	4	15	Koto and Kodama, 1962
Nov. 1942	Kaohsiung	22	82	5	18	Koto and Kodama, 1962
Dec. 1942	Suao	60	82	14	18	Koto and Kodama, 1962
Jan. 1943	Suao	18	75	6	25	Koto and Kodama, 1962
Feb. 1943	Kaohsiung	16	94	1	6	Koto and Kodama, 1962
Mar. 1943	Kaohsiung	9	90	1	10	Koto and Kodama, 1962
Apr. 1943	Kaohsiung	16	94	1	6	Koto and Kodama, 1962
May 1943	Kaohsiung	15	100	0	0	Koto and Kodama, 1962
Dec. 1943	Suao	78	86	13	14	Koto and Kodama, 1962
Feb. 1944	Suao	83	85	15	15	Koto and Kodama, 1962
Feb. 1944	Kaohsiung	24	89	3	11	Koto and Kodama, 1962
Mar. 1944	Kaohsiung	32	100	0	0	Koto and Kodama, 1962
1942-1944	Kaohsiung	107	95	6	5	Nakamura, 1944a
1942-1944	Suao	254	84	47	16	Nakamura, 1944a
Nov. 1956	Northwestern area of the Coral Sea	541	88	74	12	Ueyanagi, 1960
1952-1956	Eastern Sea of Formosa	361	88	51	12	Koto and Kodama, 1962
Sep. 1964-Dec. 1967	Equatorial western Indian Ocean	7	88	1	12	Merrett, 1971

3.23 Adolescent phase

No data on the stages between 23 and ca. 1,300 mm body length.

3.3 Adult Phase

3.31 Longevity

No data.

3.32 Hardiness

No data.

3.33 Competitors

Generally, large sharks are thought to be competitors or predators and tunas and small marlins are thought to be competitors or victims for large marlins (*Makaira* spp.) and swordfish (Parin, 1968).

3.34 Predators

Pelagic carnivorous fishes such as sharks, scombroids, carangoids, and other istiophorids are predators of larval and juvenile stages of the black marlin. There are no true predators on the adult fish.

3.35 Parasites, diseases, injuries, and abnormality.

No data except parasites on which some descriptions are in the papers of Silas (1967) and Silas and Ummerkutty (1967).



Figure 4.—Larvae of the black marlin. A. 3.6 mm in body length. B. Approximately 5.2 mm in body length. C. 23 mm in body length. (From Ueyanagi, 1960.)

3.4 Nutrition and Growth

3.41 Feeding

Very little is known about the feeding behavior of the black marlin. During fishing operations off Hana, Maui, a large black marlin (referred to as silver marlin in the publication; see footnote 2) was caught on a longline at a depth of about 60 fathoms. The fish had an abnormally distended abdomen and, upon gutting, a bigeye tuna, *Thunnus obesus* Lowe, was found in the stomach. This tuna, which weighed about 160 pounds (72.7 kg), had been caught on the longline first, for the hook was still set in the tuna's mouth when the specimen was removed from the stomach of the marlin. The marlin evidently had found the hooked tuna an easy prey and swallowed it head first (June 1951). Black marlin have often been observed to swallow adult skipjack tuna, *Euthynnus pelamis* (Linnaeus), or bigeye tuna, head first (Togo, pers. commun.[4]).

3.42 Food

Tanoue (1953) studied the food of the black marlin near Mangole Island and Timor Island. The most important food items of the black marlin from Mangole Island were various fishes such as *Decapterus, Auxis, Gempylus serpens* Cuvier, *Pseudoscopelus,* Tetraodontidae, Atherinidae, and Paralepididae and decapod molluscs. Principal food items of black marlin from Timor Island were fishes such as Syngnathidae, *Auxis* and *Oxyporhamphus,* and megalop larvae of Crustacea, Isopoda, and decapod molluscs.

In the East China Sea, most important food items of black marlin are mackerels, *Scomber* spp. and horse mackerels, *Trachurus* spp. (Morita, 1960).

Watanabe (1960) studied the food of the black marlin in the Pacific equatorial waters. Fishes of the families Scombridae, Gempylidae, Coryphaenidae, Xiphiidae, and Carangidae and decapod molluscs were the most important food of the black marlin. Fishes of the families Sternoptychidae, Paralepididae, Alepisauridae, Chiasmodontidae, Chaetodontidae, Balistidae, Ostraciontidae, and Tetraodontidae, and octopods and macrura crustaceans were of lesser importance.

3.43 Growth Rate

Koto and Kodama (1962) studied the black marlin of the East China Sea, and the annual growth in length was estimated as follows: 1) in the 150- to 200-mm group (the minimum size class in the commercial catches) the annual growth is 50 cm; 2) in the 200- to 230-cm group, which is 1 yr older than the abo group, it is 30 cm; 3) in the 230- to 250-cm group, i 20 cm; and 4) above the 250-cm group, it is difficult estimate the annual growth, as seasonal shifts of t modes are not discernible. Black marlin in the E China Sea grow most rapidly during the period fr early summer to late autumn.

3.44 Metabolism

No studies.

3.5 Behavior

3.51 Migration and local movement

In the East China Sea, the schools of the bla marlin migrate northward during spring and summ and migrate southward during autumn and win (Koto et al., 1959). In the Japan Sea some bla marlin seem to migrate northward with the core of t Tsushima Current during summer and migra southward against the Tsushima Current during la summer and early autumn (Nakamura, unpu data).

In the tropical western Indian Ocean, the bla marlin is more abundant during the northeast mo soon period than the southeast monsoon (Merre 1971).

3.52 Schooling

There is little data on the schooling behavior of t black marlin. When the monsoonal winds chan suddenly in direction and intensity around the wate of Taiwan, schools of black marlin appear at the su face. The harpoon fishery becomes active in th period (Nakamura, 1938).

3.53 Responses to stimuli

No data.

4. POPULATION

- 4.1 Structure
- 4.2 Abundance and Density of Population
- 4.3 Natality and Recruitment
- 4.4 Mortality and Morbidity
- 4.5 Dynamics of Population as a Whole
- 4.6 The Population in the Community and t Ecosystem

There are a few fragmentary observations on the topics under item 4, but there are no specific studie

[4]Togo, S., 1972, personal communication on feeding habits of the black marlin. Soroku Togo is an expert technician on the research vessel of the Fisheries Agency of Japan.

EXPLOITATION

5.1 Fishing Equipment

5.11 Gears

Tuna longlines are employed to catch black marlin most fishing grounds where this species occurs; eas include the East China Sea, the Sulu Sea, the elebes Sea, and the Coral Sea. Harpoon fishing is rried out in the fishing grounds of Taiwan and uthern Japan. Sport fishing using trolling gear tches many black marlin in Piñas Bay, Panama and f Ecuador.[5] Sometimes this species is caught by set t used to catch yellowtail in the Japan Sea.

5.12 Boats

The black marlin is mostly caught by ordinary tuna ngliners. The vessels of the longline fleet vary conderably in size; the largest are about 1,900 tons with veral small catcher boats stored on each side of the ck. Longline vessels are constructed of wood or eel; those of more than 100 gross tons are usually of eel. Most of the longliners are 250 to 350 gross tons; this size they appear to be the most economical and ficient to operate (Yoshida, 1966).

Small wooden boats are employed for harpoon shing in Taiwan and southern Japan. The wooden oat is about 10 m long, and is powered by a small gine.

Ordinary trolling motorboats are employed for trollg in the sport fisheries of Panama, northeastern ustralia, and Ecuador. In these areas black marlin e caught more frequently than in other sport fishing ounds for tunas and billfishes.

5.2 Fishing Areas

5.21 General geographical distribution

The black marlin is distributed widely in the opical and temperate waters of the Indo-Pacific, but ore restricted in the Atlantic (Fig. 2). The species is ken commonly by tuna longline as a by-catch. The ecies is densely distributed in the vicinity of islands id in coastal waters. The most important commeral fishing grounds are in the East China Sea, the aters around Formosa, the waters around northwest ustralia and the Coral Sea (Figs. 2, 3).

5.22 Geographical ranges

Spawning stocks of this fish are believed to be conned to the Indian and Pacific oceans. Specimens ray occasionally into the Atlantic Ocean (Fig. 2). om commercial fisheries data the northern limit is out lat. 40°N in the northern Pacific Ocean and about lat. 45°S in the southern Pacific Ocean and the Indian Ocean. In the southern area off Tasmania, the limit extends to lat. 50°S (Fig. 2).

5.23 Depth ranges

The commercial longline fishery is generally conducted in waters over 100 m deep. In the equatorial western Indian Ocean, peak catch rates of black marlin have been reported to occur at the 201- to 500-fathom depth range (Merrett, 1971). The swimming layer of this species is usually considered to be shallower than those of other species of billfishes.

5.24 Condition of the grounds

Most areas exploited at present are coastal water, but the grounds are various.

5.3 Fishing Seasons

In the East China Sea from lat. 30° to 34°N the fishing season begins in July and ends in January (peak in August, September, and October); from lat. 25° to 30°N the fishing season occurs from May to July and October to April with peaks in June and July, and October and January (Koto et al., 1959).

The fishing season begins in October and ends in April (peak in November and December) near Uotsurijima in the East China Sea (Morita, 1952). The fishing season around Formosa extends throughout the year; the peak occurs between October and December in the South China Sea, and in February and March off eastern Formosa (Nakamura, 1953).

5.4 Fishing Operations and Results

5.41 Effort and intensity

In all of the areas, there is insufficient data on effort and intensity of commercial fishing for this species. However, it is known that most of the effort and intensity occurs in the East China Sea, the waters around Formosa, the waters around northwest Australia, the Coral Sea, the Arafura Sea, the Sulu Sea, and the Celebes Sea.

5.42 Selectivity

No data.

5.43 Catches

Catches by Japanese tuna longliners for 1962 to 1970 are given in Table 3.

6. PROTECTION AND MANAGEMENT

6.1 Regulatory Measures

6.2 Control or Alteration of Physical Features of the Environment.

[5] Recently Cairns, Queensland of Australia has become famous for olling of black marlin.

6.3 Control or Alteration of Chemical Features of the Environment

6.4 Control or Alteration of Biological Features of the Environment

6.5 Artificial Stocking

No data on above mentioned items.

7. POND FISH CULTURE

No data.

LITERATURE CITED

FISHERIES AGENCY OF JAPAN, RESEARCH DIVISION.
1972. Annual report of effort and catch statistics by area on Japanese tuna longline fishery, 1970. 326 p.

HOWARD, J. K., and S. UEYANAGI.
1965. Distribution and relative abundance of billfishes (Istiophoridae) of the Pacific Ocean. Univ. Miami Inst. Mar. Sci., Stud. Trop. Oceanogr. 2, 134 p.

JUNE, F. C.
1951. Note on the feeding habits of the giant white marlin of the Pacific. Pac. Sci. 5:287.

KOTO, T., I. FURUKAWA, and K. KODAMA.
1959. Studies on the tuna long line fishery in the E China Sea. IV. Ecological studies on the so-called wh marlin, *Marlina marlina* (Jordan & Hill). [In Jap., En abstr.] Rep. Nankai Reg. Fish. Res. Lab. 11:108-129.

KOTO, T., and K. KODAMA.
1962. Some considerations on the growth of marlins, usi size-frequencies in commercial catches. II. Attempts estimate the growth of so-called white marlin, *Marli marlina* (J. and H.). [In Jap., Engl. abstr.] Rep. Na kai Reg. Fish. Res. Lab. 15:109-126.

LAMONTE, F., and D. E. MARCY.
1941. Swordfish, sailfish, marlin and spearfish. Ichthy Contrib. Int. Game Fish Assoc. 1(2):1-24.

MERRETT, N. R.
1970. Gonad development in billfish (Istiophoridae) fro the Indian Ocean. J. Zool. (Lond.) 160:355-370.
1971. Aspects of the biology of billfish (Istiophoridae) fro the equatorial western Indian Ocean. J. Zool. (Lond 163:351-395.

MORITA, T.
1952. On the relation between the yield of marlin and t water temperature around the Uotsurizima. [In Jap Engl. summ.] Kagoshima Univ., Mem. Fac. Fish. 2(1 15-19.
1960. Fishing-ground constitutional studies on the whit marlin, *Marlina marlina* JORDAN & HILL, over the Ea China Sea—I. On the fishing-ground constitution occu ring with the seasonal change of the environment. [Jap., Engl. abstr.] Bull. Jap. Soc. Sci. Fish. 26:887-89

Table 3.—Annual estimates by major fishing grounds, of fishing effort in numbers of hooks and catch numbers of black marlin, 1962-70 (Anonymous, 1972).

Year	All major fishing grounds combined		Whole Pacific Ocean		North Pacific Ocean region		South Pacific Ocean region	
	Hooks (× 1,000)	Number of fishes (× 1,000)	Hooks	No.	Hooks	No.	Hooks	No.
1962	412,571	85	290,051	33	156,140	5	109,183	26
1963	449,427	62	337,113	32	154,036	4	130,976	23
1964	437,010	62	283,140	27	139,893	6	81,248	16
1965	466,204	69	288,771	38	152,942	12	92,226	23
1966	445,011	69	301,617	35	149,691	7	104,382	24
1967	463,461	58	305,999	22	192,960	6	70,615	12
1968	435,665	64	286,853	17	165,372	3	71,040	10
1969	436,815	58	306,024	22	168,750	5	70,114	13
1970	402,290	44	282,531	18	162,674	4	67,735	12

Year	Eastern Pacific Ocean region		Indian Ocean region		Atlantic Ocean region	
	Hooks	No.	Hooks	No.	Hooks	No.
1962	24,727	2	68,416	49	54,104	3
1963	52,101	4	57,309	30	55,004	1
1964	62,000	4	68,872	36	84,998	0
1965	43,603	3	79,852	31	97,581	0
1966	47,544	4	89,580	33	53,814	0
1967	42,425	4	126,307	36	31,154	0
1968	50,441	4	118,565	47	30,247	0
1969	67,159	4	101,115	36	29,676	0
1970	52,122	3	78,180	26	41,580	0

ORROW, J. E.
1954. Fishes from East Africa, with new records and descriptions of two new species. Ann. Mag. Nat. Hist. Ser. 12, 7:797-820.
1957. Shore and pelagic fishes from Peru, with new records and the description of a new species of Sphoeroides. *Makaira*. Bull. Bingham Oceanogr. Collect., Yale Univ. 16(2):5-55.
1959a. On *Makaira nigricans* of Lacépède. Yale Peabody Mus. Nat. Hist. 39:1-12.
1959b. Distribution of the blue marlin and black marlin in the Indo-Pacific. Bull. Mar. Sci. Gulf Caribb. 9(3):321-323.
1959c. *Istiompax indicus* (Cuvier) 1831, a prior name for the black marlin. Copeia 1959:347-349.

AKAMURA, H.
1938. Report of an investigation of the spearfishes of Formosan waters. [In Jap.] Reports of the Taiwan Government-General Fishery Experiment Station, No. 10. (Translated by W. G. Van Campen, 1955, U.S. Dep. Int., Fish Wildl. Serv., Spec. Sci. Rep. Fish. 153, 46 p.)
1941. On the habits of the Istiophorid fishes, particularly the spawning of the black marlin, *Makaira mazara*. [In Jap.] Zool. Mag. 53:17-21.
1942. Habits of fishes of the family Istiophoridae in Formosan waters. [In Jap.] Suisan Gakkwai Ho. 9:45-51.
1944a. Studies on the fishes of the family Istiophoridae from the Formosan waters. 7. Sexual differences in the size of fish. [In Jap.] Trans. Nat. Hist. Soc. Taiwan 34 (250):242-246.
1944b. Studies on the fishes of the family Istiophoridae from the Formosan waters. 8. Seasonal variations in size. b. White marlin. [In Jap.] Trans. Nat. Hist. Soc. Taiwan 34(251):293-297.
1953. A review on the fishing ground of tuna longline based on the past data. [In Jap.] Rep. Nankai Reg. Fish. Res. Lab. 1:1-144.

AKAMURA, I., T. IWAI, and K. MATSUBARA.
1968. A review of the sailfish, spearfish, marlin and swordfish of the world. [In Jap.] Kyoto Univ., Misaki Mar. Biol. Inst., Spec. Rep. 4, 95 p.

ICHOLS, J. T., and F. R. LaMONTE.
1935a. The Tahitian black marlin, or silver marlin swordfish. Am. Mus. Novit. 807, 2 p.
1935b. How many marlins are there? Nat. Hist. 36:327-330.

ARIN, N. V.
1968. Ikhtiofauna okeanskoi epipelagiali (Ichthyofauna of the epipelagic zone). Akad. Nauk SSSR, Tr. Inst. Okeanol., Moskva. (Translated by M. Raveh, 1970, Isreal Program Sci. Transl., 206 p.)

IVAS, L. R.
1956. Definitions and methods of measuring and counting in the billfishes (Istiophoridae, Xiphiidae). Bull. Mar. Sci. Gulf Caribb. 6:18-27.

ROSA, H., JR.
1950. Scientific and common names applied to tunas, mackerels and spearfish of the world with notes on their geographic distribution. FAO (Food Agric. Organ. U.N.), Wash., D.C., 235 p.

ROYCE, W. F.
1957. Observations on the spearfishes of the central Pacific. U.S. Fish Wildl. Serv., Fish. Bull. 57:497-554.

SILAS, E. G.
1967. Parasites of scombroid fishes. Part I. Monogenetic trematodes, digenetic trematodes, and cestodes. Proc. Symp. Scombroid Fishes, Part 3, p. 799-875. Mar. Biol. Assoc. India, Mandapam Camp.

SILAS, E. G., and A. N. P. UMMERKUTTY.
1967. Parasites of scombroid fishes. Part II. Parasitic copepoda. Proc. Symp. Scombroid Fishes, Part 3, p. 876-993. Mar. Biol. Assoc. India, Mandapam Camp.

TANOUE, T.
1953. An experiment on the tuna fishery by long-line in the sea off Mangole and Timor Islands. [In Jap., Engl. summ.] Mem. Fac. Fish., Kagoshima 3(1):1-34.

UEYANAGI, S.
1960. On the larvae and the spawning areas of the Shirokajiki, *Marlina marlina* (Jordan & Hill). [In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. Lab. 12:85-96.
1963. A study of the relationships of the Indo-Pacific istiophorids. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 17:151-165.
1964. Description and distribution of larvae of five istiophorid species in the Indo-Pacific. Proc. Symp. Scombroid Fishes, Part 1, p. 499-528. Mar. Biol. Assoc. India, Mandapam Camp.

UEYANAGI, S., and H. YABE.
1960. On the larva possibly referable to *Marlina marlina* (Jordan & Hill). Rec. Oceanogr. Works Jap., New Ser. 5(2):167-173.

WATANABE, H.
1960. Regional differences in food composition of the tunas and marlins from several oceanic areas. [In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. Lab. 12:75-84.

WHITLEY, G. P.
1931a. New names for Australian fishes. Aust. Zool. 6(4): 310-334.
1931b. Studies in ichthyology. No. 5. Rec. Aust. Mus. 18(4):138-160.
1948. A list of the fishes of Western Australia. West. Aust. Fish. Dept. Bull. 2, 35 p.
1955. The Australian Museum's marlins. Aust. Mus. Mag. 11:292-297.

YOSHIDA, H. O.
1966. Tuna fishing vessels, gear, and techniques in the Pacific Ocean. *In* T. A. Manar (editor), Proc. Gov. Conf. Cent. Pac. Fish. Resour., p. 67-89. State of Hawaii.

Synopsis of Biological Data on the Longbill Spearfish *Tetrapturus pfluegeri* Robins and de Sylva[1]

C. RICHARD ROBINS[2]

1 IDENTITY

1.1 Nomenclature

1.11 Valid name

The valid scientific name of this taxon is *Tetrapturus pfluegeri* Robins and de Sylva (1963:86-89).

1.12 Objective synonymy

All previous references to the species were incorrectly or incompletely attributed, mostly to the Mediterranean spearfish, *Tetrapturus belone* Rafinesque. In describing *pfluegeri,* Robins and de Sylva (1963:86, 89-90) gave complete synonymies for both species. There are no earlier invalid names nor any junior synonyms.

1.2 Taxonomy

1.21 Affinities

Tetrapturus pfluegeri has been assigned by all authors to *Tetrapturus* Rafinesque (1810) as employed by Robins and de Sylva (1960:402-406). This genus is one of three living genera recognized by those authors in the family Istiophoridae which in turn comprises with the Xiphiidae in suborder Xiphioidei of the order Perciformes.

A synonymy of *Tetrapturus* was given by Robins and de Sylva (1960:385). This is repeated below:

Tetrapturus Rafinesque

Tetrapturus Rafinesque, 1810: 54-55 (*T. belone,* type species by monotypy).

Skeponopodus Nardo, 1832: 99 (*nomen nudum);* 1833: 415-419 (*S. typus,* [= *T. belone* Rafinesque], type species by virtue of the name *typus*).

Tetrapterurus Bonaparte, 1841: 19 (emended spelling).

Tetrapterus Agassiz, 1843: 7, 89-92, table E (emended spelling).

Tetraplurus Vérany, 1847: 492-494 (misprint for *Tetrapturus*?).

Scheponopodus Canestrini, 1872: 112 (emended spelling).

Tetraperus Radcliffe, 1926: 112 (misprint for *Tetrapturus*).

Marlina Grey, 1928: 47 (*Tetrapturus mitsukurii* Jordan and der, [= *T. audax* Philippi], type species by monotypy; t of *Marlina* at the generic level is probably a slip).

Kajikia Hirasaka and Nakamura, 1947: 13-14 (*Kajikia sana,* [= *Tetrapturus audax* Philippi], type species by typy).

Pseudohistiophorus de Buen, 1950: 171 (*Tetrapturus illing* Jordan and Evermann [= *T. angustirostris* Tanaka], typ cies by original designation).

Lamontella Smith, 1956: 32 (*Tetrapturus albida* [sic] Poey species by original designation and monotypy).

Tetrapturus pfluegeri Robins and de Sylva

Figures 1 and 2

Type specimen.—The following designations made by Robins and de Sylva, 1963:88. The speci numbers in parentheses will enable the reader to tain the morphometric data for each specimen reference to Robins and de Sylva (1960: Tables 1963: Tables 1, 2). Also meristic data for the holo were indicated by an asterisk (*) in Table 5 of the paper.



Figure 1.—*Tetrapturus pfluegeri,* juvenile, 368-mm body le taken off Fort Lauderdale, Fla. Drawing by D. P. de S (after Robins and de Sylva, 1963:Fig. 2).

Figure 2.—*Tetrapturus pfluegeri,* adult, 1,482-mm body le 24 lb, taken off Miami, Fla. (spec. 4, see Robins and de Sy 1960:392-393 for further data).

[1]Contribution No. 1709, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, FL 33149.

[2]Maytag Professor of Ichthyology, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, FL 33149.

Table 1.—Fin-ray counts of the longbill spearfish, *Tetrapturus pfluegeri.*[1]

Dorsal spines									Second dorsal rays		
45	46	47	48	49	50	51	52	53	5	6	7
1	4	6	7	11*	4	3	1	1	2	23	6*

Anal spines								Second anal rays				Pectoral rays (left fin)			
11	12	13	14	15	16	17	18	5	6	7	8	17	18	19	20
1	2	5	9	7	4*	-	1	1	11	19*	1	1	11	13	5*

[1]Data from Robins and de Sylva (1963: Table 5, supplemented by counts from specimens reported on here).

*Indicates count of holotype of *T. pfluegeri.*

Holotype.—USNM 202818, formerly UMML 2231 pec. 5), an adult, 1,530 mm, caught by R. E. Maytag f San Juan, Puerto Rico, 1 October 1957, photo.

Paratypes.—UMML 5955, an adult, 1,790 mm, ught by M. A. Madden on 1 June 1958, east of iami Beach, Fla., photo (spec. 15). UMML 2230, a venile, 914 mm, caught off Palm Beach, Fla., 23 larch 1958, photo (spec. 7). UMML 5956, an adult, 655 mm, caught 1.5 miles off Hillsboro Light, late ecember 1957, or early January 1958 (spec. 8). MML 3484 (skeleton only), 1,482 mm, caught off iami, Fla., 14 August 1957, by S. Bergman, photo pec. 4). Spec. 21a, 1,740 mm, female, lat. 12°38′N, ng. 69°17′W, MV *Oregon* station 2764, 10 April 1960 pecimen at the American Museum of Natural istory). Spec. 22, 1,620 mm, female, lat. 12°36′N, ng. 63°40′W, MV *Oregon* station 2768, 14 April 960 (specimen at the American Museum of Natural istory). UMML 11095, juvenile, 368 mm (spec. 31), e data listed above and Figure 1.

ype locality.—Off San Juan, Puerto Rico.

iagnosis.—The following is repeated from Robins nd de Sylva (1963:86).

> . . . First dorsal elements 45-53 (usually 46-50), second dorsal elements 6-7 (usually 6), first anal elements 12-16 (usually 14), second anal elements 6-7 (usually 7), pectoral rays 18-20 (usually 19); second anal elements usually one more than second dorsal elements. Vertebrae 24 (12 precaudal, 12 caudal). Anus far anterior to anal-fin origin, distance 8.4-11 per cent body length (tip of lower jaw to fork of tail) and usually greater than the anterior height of first anal fin. First dorsal fin unspotted and with high profile, especially in juveniles, its 25th element varying from 17 per cent of body length (at 914-mm body length) to 7.7 per cent of body length (at 1,790 mm body length). Pectoral fin becoming proportionally longer with growth of fish (12-13 per cent of body length at 914 mm to 20-22 per cent at 1,460 mm or larger). Dorsal profile straight from origin of first dorsal-fin to a point in front of eyes. Bill long, the distance from the tip of the upper jaw to the anterior margin of the eye 27-19 per cent of body length, the lower figures characterizing the larger specimens (bill growth is negatively allometric).

Material examined.—In addition to material examined and reported on by Robins and de Sylva (1960, 1963), the writer has studied the following specimens.

Spec. 37a, 1,808 mm body length, 51 lb (23.1 kg), off Ft. Lauderdale, Fla., 19 June 1964, T. Treadway.

Spec. 38, 1,803 mm, 56.6 lb (25.7 kg), off Bakers Haulover, Dade County, Fla., 10 May 1964, M. Lojinger.

Spec. 39, 1,320 mm, 14.5 lb (6.6 kg), off Miami, Fla., 25 July 1964, G. H. Ludins.

Spec. 40, 1,617 mm, 38.8 lb (17.6 kg), off Miami Beach, Fla., March 1965.

Spec. 41, 1,616 mm, 44.5 lb (20.2 kg), off Ft. Lauderdale, Fla., 19 November 1965, Nancy Brodno.

Spec. 42, 853 mm, ca. 2 lb (0.9 kg), off Islamorada, Fla., 26 January 1966, Paul Wright.

Spec. 43, 871 mm, in surf at Miami Beach, Fla., 22 March 1965, Bud Raulston.

Spec. 44, 1,530 mm, 27.5 lb (12.5 kg), off Ft. Lauderdale, Fla., J. H. Van Ness.

Spec. 45, 1,875 mm, 60 lb (27.2 kg), off Islamorada, Fla., 15 July 1968.

Spec. 46, 1,880 mm, 80 lb (36.3 kg), off Chub Cay, Bahamas, 15 April 1971, Edwin Jay Gould.

Spec. 47, ca. 600 mm, *Oregon II,* stn. 108, lat. 12°53′N, long. 70°35′W, 23 February 1973.

During this same period other specimens from south Florida were seen and identified by the writer or de Sylva but were not measured, usually because the lower jaw was broken precluding an accurate measure of body length (= tip of lower jaw to fork of tail).

Data on specimens 37 through 47 are provided in Tables 2 and 3. In combination with previously published data this information should provide a basis for morphometric comparison with samples from the South Atlantic and elsewhere in the range for which such information is not yet available.

Synonymy.

Tetrapturus belone: LaMonte, 1955:326 (in part; reference to Florida only).—Hoese, 1958:341 (com-

Table 2.—Morphometric data for 11 specimens of *Tetrapturus pfluegeri* from the Western North Atlantic. Measurements (in millimeters) are as defined by Rivas (1956) unless otherwise indicated. Numbers in parentheses refer to the numbered definitions of R see Robins and de Sylva, 1960:384-385, for explanation of abbreviations.

Specimen number	47	42	43	39	44	41	40	38	37a	45
Body length (1)	600±	853	871	1,320	1,530	1,616	1,617	1,803	1,808	1,875
First predorsal length (3)	--	181	176	280	325	317	337	383	378	378
Second predorsal length (4)	--	699	728	1,095	1,240	1,308	1,337	1,497	1,482	1,538
Prepectoral length (5)	--	219	212	322	370	374	405	443	440	448
Prepelvic length (6)	--	205	222	345	395	410	423	457	468	485
First preanal length (7)	--	528	528	845	910	995	990	1,130	1,089	1,150
Second preanal length (8)	--	688	720	1,075	1,225	1,290	--	1,482	1,438	1,505
Orig. D_1 to orig. P_1 (9)	55	64	--	124	142	--	--	183	196	--
Orig. D_1 to orig. P_2 (10)	72	95	--	160	188	--	--	234	246	--
Orig. D_2 to orig. A_2 (11)	42	--	--	94	115	--	--	130	143	146
Tip mandible to anus	--	451	451	705	780	846	855	934	913	972
Orig. P_2 to nape (13)	61	117	--	173	208	--	--	250	244	--
Greatest body depth (14)	62	78	--	144	182	--	--	230	227	--
Depth at orig. D_1 (15)	58	76	87	144	182	209	207	228	227	--
Depth at orig. A_1 (16)	55	65	74	119	156	195	187	198	185	--
Least depth c. p. (17)	17	24	25	40	50	55	54	57	55	59
Width at P_1 base (18)	25	31	37	65	91	93	90	111	115	--
Width at A_1 orig. (19)	26	32	39	70	87	119	105	123	107	--
Width at A_2 orig. (20)	20	28	--	63	75	--	--	95	83	--
Width c.p. (in front of keels)	12	14	18	28	35	49	40	38	42	49
Length upper keel (22)	16	20	23	38	44	49	40	51	56	64
Length lower keel (23)	15	19	25	41	44	47	40	54	54	58
Head length (24)	--	213	207	364	--	367	381	433	427	433
Snout length (25)	--	106	102	189	183	173	192	210	221	--
Bill length (26)	159±	234	235+	302	343	344	374	480	391	413
Maxillary length (28)	--	131	132	209	235	218	246	270	272	270
Orbit diameter (29)	17	22	26	36	42	40	41	47	53	45
Depth of bill (33)	5.4	8.2	--	10	11	--	--	10	12	14
Width of bill (34)	6.6	9.9	--	15	17	--	--	17	19	22
Height D_1 (39)	108	172	152	213	292	243	273	255	265	268
Length 25th D_1 Spine (40)	121	143	130	--	--	146	147	--	144	130
Height D_2 (41)	21	29	28	53	56	94	72	72	61	63
Height A_1 (42)	50	82	83	129	184	143	188	150	162	158
Height A_2 (43)	16	21	26	40	52	51	55	51	55	57
Length P_1 (44)	58	118	114	249	345	356	378	360	371	403
Length P_2 (45)	140	--	160	295±	342	354	--	--	367	403
Length last D_2 ray	23	37	40	69	92	63	103	104	93	106
Length last A_2 ray	24	36	42	67	78	91	94	104	92	95
Orig. D_1 + Orig. D_2	381	546	562	830	930	973	998	1,139	1,133	1,170
Anus to orig. A_1	55	77	80	115	136	154	139	179	173	176
Weight (lb)	1.1	2	--	14.5	27.5	44.5	38.8	56.5	51	60

piled; Texas).—Migdalski, 1958:70 (Puerto Rico and Florida; lower figure opposite p. 100 except for fish at upper right).—Robins and de Sylva, 1960 (in part; diagnosis, description, and data on *belone* in Tables 1-3, and Figs. 1, 2, 3b, 4 entirely referable to *pfluegeri*; those items in synonymy which are based solely on Mediterranean specimens or accounts and discussion based on photographs of Mediterranean specimens are not referable to *pfluegeri*).—Tortonese, 1962:8-9 (in part; information referring to Florida only).—Cavaliere, 1962:171 (in part; reference to western Atlantic material reported by Robins and de Sylva, 1960).—Peronaci, 1966 (in part, Atlantic references only; *T. pfluegeri* regarded to be synonym of *T. belone*).

Young *Makaira:* LaMonte, 1955:346-347 (fish take "spring of 1939 half-way between Bimini, Baha and Miami, Florida").

Spearfish, Swann, 1957:15, 28 (photograph of sp men captured off Port Aransas, Texas; subseque ly reported on by Springer and Hoese [1958] *Tetrapterus belone* [?] and listed by Hoese [19 as *Tetrapturus belone*).

Tetrapterus belone: Robins, 1958:16 (in part, Flo only).

Tetrapterus belone (?): Springer and Hoese, 1958: 346 (Port Aransas, Texas; measurements, cou coloration).

ble 3.—Morphometric data for 11 specimens of *Tetrapturus pfluegeri* from the Western North Atlantic expressed in percentage of dy length. Measurements are as defined by Rivas (1956) unless otherwise indicated. Numbers in parentheses refer to the numbered finitions of Rivas; see Robins and de Sylva, 1960:384-385, for explanation of abbreviations.

ecimen number	47	42	43	39	44	41	40	38	37a	45	46
dy length (1)	[1]600±	853	871	1,320	1,530	1,616	1,617	1,803	1,808	1,875	1,880
rst predorsal length (3)	--	21	20	21	22	20	21	21	21	20	21
cond predorsal length (4)	--	82	84	83	81	81	83	83	82	82	81
epectoral length (5)	--	26	24	24	24	23	25	25	24	24	24
epelvic length (6)	--	24	25	26	26	25	26	25	26	26	26
rst preanal length (7)	--	62	61	65	60	62	61	63	60	61	62
cond preanal length (8)	--	81	83	81	80	80	--	82	80	80	81
ig. D_1 to orig. P_1 (9)	9.2	7.6	--	9.4	9.3	--	--	10	11	--	11
ig. D_1 to orig. P_2 (10)	12	11	--	12	12	--	--	13	14	--	15
ig. D_2 to orig. A_2 (11)	6.9	--	--	7.2	7.5	--	--	7.2	7.9	7.8	8.4
p mandible to anus	--	53	52	53	51	52	53	52	50	52	54
ig. P_2 to nape (13)	10	14	--	13	14	--	--	14	14	--	--
eatest body depth (14)	10	9.1	--	11	12	--	--	13	13	--	--
epth at orig. D_1 (15)	10	8.9	10	11	12	13	13	13	13	--	13
epth at orig. A_1 (16)	9	7.6	7.4	9.1	10	12	12	11	10	--	13
ast depth c.p. (17)	2.9	2.8	2.5	3.0	3.3	3.4	3.3	3.2	3.0	3.1	3.6
idth at P_1 base (18)	4.1	3.6	4.2	4.9	5.9	5.8	5.6	6.1	6.4	--	6.2
idth at A_1 orig. (19)	4.3	3.8	4.5	5.3	5.7	7.4	6.5	6.8	5.9	--	7.6
idth at A_2 orig. (20)	3.4	3.3	--	4.7	4.9	--	--	5.3	4.6	--	--
idth c.p. (in front of keels)	3.8	1.6	2.1	2.1	2.3	3.2	2.5	2.1	2.3	2.6	2.5
ngth upper keel (22)	2.7	2.3	2.6	2.9	2.8	3.2	2.5	2.8	3.1	3.4	2.6
ngth lower keel (23)	2.3	2.2	2.9	3.1	2.8	2.9	2.5	3.0	3.0	3.1	2.8
ead length (24)	24	25	23	28	--	23	24	24	24	23	23
out length (25)	--	12	12	14	12	11	12	12	12	--	--
ill length (26)	26	27	27	23	22	21	23	27	22	22	21
axillary length (28)	--	15	15	16	15	14	15	15	15	14	14
rbit diameter (29)	2.8	2.6	3.0	2.7	2.7	2.5	2.5	2.6	2.9	2.4	2.5
epth of bill (33)	1.1	0.9	--	0.7	0.7	--	--	0.6	0.7	0.7	0.7
idth of bill (34)	--	1.1	--	1.1	1.1	--	--	0.9	1.0	1.2	1.2
eight D_1 (39)	18	20	17	16	19	15	17	14	15	14	14
ngth 25th D_1 Spine (40)	20	17	15	--	--	9.0	9.1	--	8.0	6.9	5.7
eight D_2 (41)	3.5	3.4	3.2	4.0	3.7	5.8	4.4	4.0	3.4	3.4	--
eight A_1 (42)	8.3	9.6	9.5	9.8	12	14	12	8.3	9.0	8.4	9.4
eight A_2 (43)	2.8	2.5	3.0	3.0	3.4	3.2	3.4	2.9	3.0	3.0	2.6
ngth P_1 (44)	10	14	13	19	22	22	23	20	20	22	20
ngth P_2 (45)	23	--	18	22±	22	22	--	--	20	22	--
ngth last D_2 ray	3.8	4.3	4.6	5.2	6.0	3.9	6.4	5.8	5.1	5.6	--
ngth last A_2 ray	4.0	4.2	4.8	5.1	5.1	5.6	5.8	5.8	5.1	5.1	4.9
rig. D_1 + orig. D_2	63	64	64	63	61	60	62	63	63	62	63
nus to orig. A_1	10	9.0	9.2	8.7	8.9	9.5	8.6	9.9	9.6	9.4	9.7

[1]Mandible broken, percentages approximate.

etrapturus beloni [sic]: Briggs, 1958:287 (listed from Florida; distribution, habitat).

etrapterus . . . . . Robins, 1958:17 (southeastern Florida).—Erdman and Roman, 1959:figure on p. 117 (Puerto Rico).

etrapturus sp. Migdalski, 1958:70-71, lower figure opposite p. 100, except for fish in upper right (Florida and Texas).

etrapturus pfluegeri Robins and de Sylva, 1963:86-89, Tables 1, 2, Figs. 1, 2 (original description; distinguished from *T. belone*).—De Sylva, 1963:125, 130 (juvenile compared to other Atlantic species). Anon., 1963:53 (review of original account of species); 1964a:40 (one reported caught during operations of MV *Delaware* off New England); 1964b:31 (one reported caught during operations of MV *Delaware* between North Carolina and Georges Bank).—Ueyanagi and Watanabe, 1965 (differences in vertebral column between various istiophorids including *T. pfluegeri*).—Nakamura, Iwai, and Matsubara, 1968 (review of history and morphometry; new data provided on Atlantic distributions of adults and larvae and on various aspects of anatomy).—Ovchinnikov, 1970 (recorded from Caribbean Sea; various notes on eye size, swimming capability).—Ueyanagi et al., 1970:21-23, Figs. 7, 9-11 (geographical and seasonal distribution of adults and larvae, notes on maturity).—Anon., 1972:40 (photograph of 22-lb (10.0-kg) specimen from off Dade County,

Fla.—Penrith and Talbot, 1973 (morphometry of two specimens taken northwest of Capetown in the southeastern Atlantic).—Wise and Davis, 1973:20 (comments on meaning of Japanese longline data regarding this species).

A key to all istiophorid species was presented by Robins and de Sylva (1960) based on all species known at that time. Since then *Tetrapturus pfluegeri* was described and *T. georgei* resurrected. De Sylva (1974) in his summary of information for *Tetrapturus belone* has presented a revised and inclusive key to all species of *Tetrapturus.*

1.22 Taxonomic status

The taxonomic status of *T. pfluegeri* is clear insofar as the juveniles and adults are easily distinguished from all other billfishes. Its distinctive morphology establishes its species status. Only Peronaci (1966) among recent writers has failed to perceive its distinctiveness. Virtually all morphometric data available on *pfluegeri* are from the western Atlantic and especially Florida. There is no basis for commentary on geographic variation except that manuscript data on two specimens from South Africa from Penrith and Talbot (1973), kindly provided by M. J. Penrith, show perfect agreement with Florida specimens of comparable size from the opposite end of the range of the species. This plus experience with related species strongly suggests that we should not expect noteworthy geographic variation in the species. This does not mean that less obvious but measurable differences might not exist.

1.23 Subspecies

No subspecies are recognized.

1.24 Standard common names and vernacular names

The standard name for the species (Bailey et al., 1970) is longbill spearfish, the name originally attributed to it by its describers. It is known by anglers that recognize it by this name or simply as spearfish. The Japanese name is Kuchinaga-furai (see Howard and Ueyanagi, 1963) or Kuchinagufurai (Wise and Davis, 1973:2). The Soviet literature terms it malyi kop'enosets. Other names are those loosely applied to any of the billfishes.

1.3 Morphology

Fin rays.—T. pfluegeri has a high dorsal spine count 45-53, usually 48-51. Care must be taken to count the last one or two which may be imbedded in the skin anterior to the second dorsal fin in large adults. No other Atlantic istiophorid has such a high count. There are usually 6 (rarely 7) second dorsal and usually 7 (occasionally 6 and rarely 8) second anal rays. There are 12-18 (usually 13-10) elements in the first anal fin. Pectoral rays vary from 17 to 20, but are m frequently 19. Variation in fin ray counts may be s by reference to Table 1, which is based on data fi Robins and de Sylva (1963), supplemented with d reported for the first time in Tables 2 and 3.

Vertebrae.—As with all species of *Tetrapturus* th are always 24 vertebrae divided equally betw precaudal and caudal, the division sharply defi (see Nakamura et al., 1968, Fig. 7). The verteb column of *pfluegeri* is illustrated by Ueyanagi Watanabe (1965:pl. 3a, b).

Fins.—The spinous dorsal fin is high throughout small juveniles tapering sharply downward at the several elements. A slight dip forms behind anterior lobe in larger juveniles, but the posterior p of the fin never exceeds the height of the anterior l as it does for example in *Istiophorus.* In adults height of the fin fades away from the anterior lobe is higher than in *T. albidus* and notably higher than species of *Makaira* nearly to its end. Apparently posterior section of the fin ceases to grow at a bo length less than 900 mm. This can easily be seen Robins and de Sylva (1960:Fig. 2, lowermost sectio The pectoral fin is positively allometric being sh (12-14% of body length) in juveniles and long (20-2 of body length) in adults. The transition is sharp a specimens of about 1,000- and 1,300-mm body leng could easily be thought to be separate species on t basis.

The first anal fin is low, decidedly less than body height at that point and pointed.

Bill.—The bill, as with all species of *Tetrapturus,* proportionally longest in juveniles, shorter in adul The name longbill spearfish was used to contrast t species from its closest allies, the Mediterrane spearfish, *T. belone,* and the shortbill spearfish, *angustirostris,* both of which are short snouted. T bill in *T. pfluegeri* is shorter than that of the wh marlin, *T. albidus,* or the sailfish, *Istiopho platypterus.*

Dorsal profile.—The profile is straight from the ba of the bill to the origin of the dorsal fin; this spec thus lacks the humped nape of *T. albidus.*

Color.—The color is dark bluish black above, silv on the sides, and whitish below; the membrane of spinous dorsal fin is bright blue without dark spo Vertical barring is never prominent though pale b are usually visible when a specimen is freshly caug Juveniles have a dark blotch from the base of the 1 to the 17th dorsal spines.

There has been no published report cytomorphology or protein specificity and o scattered information on internal morpholo Nakamura et al. (1968) report generally on the int

l anatomy of billfishes using various species in-
uding *pfluegeri* to illustrate their points. The gonads
e assymetrical (Y-shaped) in *pfluegeri*.
Tetrapturus pfluegeri was described in detail by
obins and de Sylva (1960) under the name *T. belone*
d later (1963) diagnosed and distinguished from
lone and other species. Nakamura et al. 1968:62-63
so summarize its morphology.

DISTRIBUTION

2.1 Total Area

Tetrapturus pfluegeri was originally reported
nder the name *T. belone*) in the western Atlantic
om Maryland to Venezuela including the entire Gulf
Mexico and Antillean region (Robins and de Sylva,
60:Fig. 4). In formally describing the species,
obins and de Sylva (1963:88) extended the range to
ew Jersey. It was then reported from off New
ngland (Anon., 1964a) and from somewhere between
orth Carolina and Georges Bank (Anon., 1964b).
apanese longlines have taken it repeatedly and in the
rth central and South Atlantic from southeastern
razil to South Africa (see especially Ueyanagi et al.,
70:Fig. 7). Most recently it has been recorded from
outh Africa (125 miles northwest of Cape Colum-
ne) by Penrith and Talbot (1973).
Some of the Japanese data may be based on the
undscale spearfish, *T. georgei,* only recently diag-
sed (Robins, 1974), but it is clear that *T. pfluegeri*
wide ranging in the Atlantic Ocean and perhaps is
ost common in the mid- or South Atlantic. Other re-
nt references in sporting magazines are all from
thin the described range.
When this species was described and brought to the
tention of anglers in the Florida region, I was confi-
nt that the recognition would result in more being
ought to the taxidermy shops. Captains and guides
recognize the fish but catches remain few. The only
fference is that most now coming to the shops are
rrectly identified. I can only conclude that relative
other istiophorid species in the area, *T. pfluegeri* is
ly rare in waters of the Bahamas and the United
ates. Whether it will prove more common in the
id- or South Atlantic is unknown. Hook rates are
ghest (Ueyanagi et al., 1970:Fig. 7) in these regions
d in the Caribbean, but such data cannot be assess-
without knowing the actual numbers of hooks set
d the total numbers of the various fishes caught.
ghest hook rates could be experienced with the
vest hooks out and a small catch.
In summary, *T. pfluegeri* ranges widely through the
lantic from about lat. 40°N to 34°S and from New
ngland, Texas, and southeastern Brazil to between
ng. 20° and 25°W and the Cape Verde Islands in the
rth Atlantic and to South Africa in the South
lantic. It is not yet known to occur in African waters
rth of South Africa.

2.2 Differential Distribution

In data provided by Ueyanagi et al., 1970 (Figs. 7, 9-11) it is apparent that the size distribution is similar for the 11 geographic areas analyzed (see their Fig. 1) with the possible exception of areas C and F. Here samples are small and these two areas, the most eastern in the North Atlantic, are those where misidentifications of *T. georgei* are most likely to have occurred. Similarly there is no noteworthy difference in sex ratios although males may be more common in the north and west. Mature individuals were taken only in the January-March and April-June quadrants with the exception again of the area around the Cape Verde Islands and of the Caribbean where some mature individuals are recorded in October-December. In most instances these data are based on small samples. Mature individuals were not recorded north of lat. 20°N in the western Atlantic nor south of lat. 30°S. Surprisingly, maturation occurs at the same time rather than at the same season both in the northern and southern hemisphere. More than any other factor, this is suggestive of homogeneity of stock.

2.4 Hybridization

There is no morphological indication that *T. pfluegeri* hybridizes with any of its relatives. Robins (1974) considered and rejected the possibility that *T. georgei* as defined by him could be based on hybrids between *T. pfluegeri* and *T. albidus.*

3 BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

Sexes are separate in the longbill spearfish. There is no demonstrated sexual dimorphism in the morphometry of *T. pfluegeri,* nor is there any evidence of hermaphroditism.

3.12 Maturity

There has been no published study on age and growth nor age at maturity. Data from Japanese longline catches suggest that spawning occurs in late winter. This is based on their notes on seasonal occurrence of mature individuals. Looking at all available data, spawning no doubt occurs between late November and early May with a peak perhaps in late winter. To provide some answers to the question of age and growth the writer has assumed that spawning occurs in mid-February (15 Feb.) and has then grouped length and weight data from all specimens examined by him by month from that point. This was done twice, once by using the length data alone, the

second time using weight data alone. In only one instance was a spot placed differently. The results are seen in Figure 3 and indicate the presence of at most four year classes (0, I, II, III). One-year-old fish (beginning of year class I) can be expected to be about 800 to 1,000 mm in body length and 4 to 6 pounds. Two-year-old fish are about 1,600 to 1,770 mm in body length and 38 to 41 pounds. Three-year-old fish are not much longer, about 1,800 mm body length, but are heavier, 60 to 80 pounds. First spawning would occur at the end of the second year and few fish apparently survive beyond a second spawning a year later, assuming that the species spawns but once per year. A short life history is not surprising and this fit of the



Figure 3.—Age and growth analysis of *Tetrapturus pfluegeri* basd on data from the sport fishing catch in the western Atlantic. (Year classes 0, I, II, and III and months of the year are indicated on the ordinate. The assumed spawning time is indicated by the horizontal line at mid-February. See text for explanation.)

data seems most parsimonious. That is, attempt fit the data to five instead of four year classes on basis of length or weight result in greater inconsist cies in the other parameter than those exhibite Figure 3. The variation seen is remarkably small c sidering that a point-in-time spawning was assu when in fact spawning no doubt occurs over sev months.

The writer also has available length data from Pflueger taxidermy shop records; this information provided by Donald P. de Sylva. Lengths are in 2-i increments and are measured from the rear margi the eye to the fork of the tail. These data gathe from 1958 to 1967 include some of the same f analyzed by the writer, especially prior to 1964. T results are similar to those above and show 4 groupings. Because of the overlap in specimens a different means of measurement these data are further treated.

3.13 Mating

Whether spawning is in pairs or larger groups is known.

3.14 Fertilization

Nothing is known of the reproductive behavior of fecundity. Fertilization surely is external a females probably spawn but once during the year. T eggs have not been described.

3.15 Gonads

The ovaries are unequally developed, that of the l side being approximately twice that of the right. T right meets the left at an angle about half way ba the result is that the ovary appears Y-shaped wh first seen, the stem of the Y pointing forward. specimen 45 (see material examined above) the l ovary was 475 mm, the right 230 mm; in specimen these measurements are 458 and 242 mm. On t other hand the left ovary in specimen 35 was 193 m the right 212 as determined by W. P. Davis. The tes are similarly asymmetrical, again the left side be longer. In specimen 27 the two measured 65 and mm and in specimen 14 (see Fig. 4) the two are 2 and 110 mm. A study of changes in gonad size a shape with maturation is needed. In those specim with asymmetrical gonads the junction point of "Y" is almost directly above the anus. Merr (1970:357, pl. 16) notes that of all istiophorids amined by him in the Indian Ocean, only *angustirostris* Tanaka has asymmetrical gonads a that the left is the longer, the junction being at level of the urogenital operture.

3.2 Preadult Phase

De Sylva and Ueyanagi, on the occasion of International Billfish Symposium at Kona, Aug 1972, showed slides of larvae which they identified



ure 4.—Testes of *Tetrapturus pfluegeri*, 1,577-mm body gth, 34 lb, taken off Fort Lauderdale, Fla., 25 September 1958 ec. 14).

pfluegeri. This work, to be submitted to the Dana ports, is not yet in press. There is no published in-mation on the early development of *pfluegeri*. De lva (1963:125, 130) has compared the juvenile (the me individual shown in Fig. 1) with those of the trapturus albidus, *Istiophorus albicans* (=*I. atypterus*), and *Makaira nigricans*.

3.3 Adult Phase

3.31 Longevity

As noted in 3.1 and in Figure 3, few fish are more an 3 yr old (age class II). However, this is theoretical d no detailed age and growth study has been made.

3.4 Nutrition and Growth

Ovchinnikov (1970:Table 7) reported on stomach ntents of specimens from northeastern Brazil. ese data differ from those given for white marlin d Atlantic sailfish from the same region. Unfor-nately incidence of a food item is reported only in rcent of total numbers of stomachs examined and asmuch as no item appeared in fewer than 11% of e stomachs, we must conclude that only nine omachs were studied, a small sample for this type of udy. All but one of the food items are of the type at is found at or near the surface and under drift aterials. *Alepisaurus* sp. is reported from 22.5% of e stomachs (2 stomachs?). Presumably a gempylid is involved instead.

It is expected that *T. pfluegeri* feeds on small and edium-sized fishes and cephalopods in surface ters and that availability will largely determine the ture of the composition of the diet from place to ace or season to season.

3.5 Behavior

Nothing is known of migrations or local movements. southeastern Florida where there is good sport hing effort throughout the year, fish have been taken in every month. The species seems always to be present though in small numbers. There is no known area of postspawning aggregations as is the case with the white marlin. The scarcity of fish makes any tagging program unfeasible.

3.52 Schooling

Fishermen have experienced on several occasions a "double-header," that is, a situation where two fish cross the stern and take both trolled baits. Such pair formation is known for other billfishes and is common in the dolphin, *Coryphaena hippurus*. Sexes of the paired fish are unknown and frequently only one of the two involved is boated. In *C. hippurus*, the pair is usually but not always male and female. Such pair formation is more likely related to hunting procedure.

4 POPULATION

4.1 Structure

4.11 Sex ratio

Most fish examined at taxidermy shops have the internal organs soft, largely disintegrated in the case of the gonads. For this reason we have few specimens for which sex determination is positively given. Of 18 such identifications 9 each were male and female. Presumably the sex ratio is 1:1 as these few data suggest.

4.12 Age composition

For reasons explained above most of the population consists of year classes 0, I, and II, few fish entering year class III. They enter the sport fishery when they are 1 yr old and probably mature at about their second birthday. Maximum age is currently indicated at less than 4 yr, but there is no appreciable growth at this age and a few fish may live another year.

Beyond this nothing is known of population structure.

5 EXPLOITATION

5.1 Fishing Equipment

5.11 Gears

The longbill spearfish is fished by anglers in the same manner that they fish for other billfishes. No one specifically fishes for spearfish. Gear and tackle preferences vary but all are well documented in the sport fishing literature. Large baits such as those used by some anglers for blue marlin will not succeed with the much smaller spearfish. As many as 10 fish per year are sent to each of the two principal taxidermy

shops in southeastern Florida. Catches everywhere are low and probably fewer than 100 spearfish per year are caught by anglers in the western Atlantic.

Commercial longlining vessels take spearfish along with tunas, swordfish, other istiophorids, sharks, and miscellaneous other fishes. Again spearfish catches are incidental and the gear used is well described in various trade and government journals. Japanese data grouped spearfish and sailfish under "other marlin" until 1965, and since that time have listed them as "spearfish and sailfish." These data may include small marlin. It is assumed by Japanese biologists, as noted by Wise and Davis (1973:20), that nearshore records apply to sailfish, offshore records to spearfish. There is no valid reason to exclude spearfish from nearshore catches and no doubt some sailfish are included in the offshore data. Wise and Davis (1973:3) record 610,000 fish in this category caught between 1958 and 1969 with peak years of 118,000 in 1964 and 1965, the years of maximum Japanese effort, about 85 and 97.5 million hooks. Even assuming that only a small percentage are spearfish it is obvious that many more spearfish are caught commercially than by sport fishermen.

5.12 Boats

Fishing vessels, whether for sport fishing or longlining are those generally employed for the purpose and are well described in the voluminous literature on the subject.

5.2 Fishing Areas

5.21 General geographic distribution

See 5.22.

5.22 Geographic ranges

Inasmuch as all fish studied were obtained by the sport fishery or the longline fishery, it follows that the known distribution of the species, described in detail above, is that of the fishery.

The sport fishery is restricted by logistical reasons to coastal areas near ports where marina facilities exist for private yachts and where charter boats are available. Oceanic big game fishing is an expensive pastime and except for some coastal cities in the United States at which relatively inexpensive charters are available, most participants are from upper and middle income groups. Therefore, the main sport fishing centers for all istiophorid fishes are coastal tourist cities and islands with first-class tourist accommodations. Few boats fish more than 50 miles from port except along the southeastern coast of Brazil where operations may extend beyond 200 miles. Recently sport fishing boats, fishing out of St. Petersburg, Fla., and using a mother ship, have fished over the 100-fathom curve about 100 miles from shore (de Sylva, pers. comm.).

In contrast the longline fishery is largely ocea
The extent of this fishery is easily seen by the
amination of the numerous charts in Ueyanagi et
(1970). Specifically Figure 7 in that work shows
geographic distribution of catches of the long
spearfish, but the charts of the other more comm
istiophorids such as the blue and white marlins gi
better view of the geographic distribution of
fishing effort.

6 PROTECTION AND MANAGEMENT

No data.

7 POND FISH CULTURE

Not applicable.

ACKNOWLEDGMENTS

Albert Pflueger, Jr. has continued to m
specimens of spearfish and other species that enter
taxidermy plant available for study by the writer
his colleagues. I am indebted to Jon C. Staiger, Jar
P. Pardew, and Charles Getter for assitance in gath
ing and processing data and to William J. Richa
Catherine H. Robins, Martin A. Roessler, and Fra
Williams for advice and comments on the manuscr
William J. Richards generously provided data
specimen 47, collected by the RV *Oregon II*. Donal
de Sylva and I worked together on billfish projects
many years; I appreciate his advice and assistan
Support for this research was provided by
Maytag professorship in ichthyology. My partici
tion in the Kona Conference was through spe
funds provided by the Dean of the School of Ma
and Atmospheric Science, F. G. Walton Smith.

LITERATURE CITED

AGASSIZ, L.
1843. Recherches sur les poissons fossiles. Poissons
cloides. Neuchâtel 5:1-160.
ANONYMOUS.
1963. Spearfish: New species identified. Commer. F
Rev. 25(8):53.
1964a. Tuna distribution studies in North Atlantic
tinued. Commer. Fish. Rev. 26(2):37-40.
1964b. Tuna and swordfish distribution studies in wes
North Atlantic. Commer. Fish. Rev. 26(9):28-31.
1972. Some spear. Fla. Sportsman 3(4):40.

ILEY, R. M, J. E. FITCH, E. S. HERALD,
A. LACHNER, C. C. LINDSEY, C. R. ROBINS, and
B. SCOTT.
1970. A list of common and scientific names of fishes from the United States and Canada. 3rd ed. Am. Fish. Soc., Spec. Publ. 6, 149 p.

NAPARTE, C. J. L.
1841. Pesci. Iconografia della fauna Italica . . . 3(1). Rome, Salviucci, n. p.

IGGS, J. C.
1958. A list of Florida fishes and their distribution. Bull. Fla. State Mus., Biol. Sci. 2:223-318.

NESTRINI, G.
1872. Pesci. *In* E. Cornalia, Fauna d'Italia, Part 3. Vallardi, Milano, 208 p.

VALIERE, A.
1962. Notizie su biologia e pesca di *Tetrapturus belone* Raf. cenni sul'adulto e descrizione di un suo rarissimo stadio giovanile. Boll. Pesca Piscic. Idrobiol., New Ser. 15(2): 171-176.

BUEN, F.
1950. Contribucion a la ictiologia. 3. La familia Istiophoridae y descripción de una especie uruguaya (*Makaira perezi* de Buen). Publ. Cient. Serv. Oceanogr., Montevideo 5:163-178.

SYLVA, D. P.
1963. Postlarva of the white marlin, *Tetrapturus albidus*, from the Florida Current off the Carolinas. Bull. Mar. Sci. Gulf Caribb. 13:123-132.
1974. Synopsis of biological data on the Mediterranean spearfish, *Tetrapturus belone* Rafinesque. *In* R. S. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972, Part 3. Species synopsis, p. 122-132. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF 675.

DMAN, D. S., and E. ROMAN.
1959. Puerto Rico game fishing. Sea Front. 5:114-122.

EY, Z.
1928. Big game fishing in New Zealand seas. Nat. Hist. 28:46-52.

RASAKA, K., and H. NAKAMURA.
1947. On the Formosan spearfishes. Bull. Oceanogr. Inst. Taiwan 3:9-24.

ES, H. D.
1958. A partially annotated checklist of the marine fishes of Texas. Publ. Inst. Mar. Sci., Univ. Tex. 5:312-352.

WARD, J. K., and S. UEYANAGI.
1963. Japanese standard common names of Istiophoridae. Indo-Pac. Fish. Comm. FAO (Food Agric. Organ. U.N.), Occas. Pap. 63(12), 3 p.

MONTE, F. R.
1955. A review and revision of the marlins, genus *Makaira*. Bull. Am. Mus. Nat. Hist. 107:323-358.

RRETT, N. R.
1970. Gonad development in billfish (Istiophoridae) from the Indian Ocean. J. Zool. (Lond.) 160:355-370.

GDALSKI, E. C.
1958. Angler's guide to the salt water game fishes—Atlantic and Pacific. Ronald Press Co., N.Y., 506 p.

KAMURA, I., T. IWAI, and K. MATSUBARA.
1968. A review of the sailfish, spearfish, marlin and swordfish of the world. [In Jap.] Kyoto Univ., Misaki Mar. Biol. Inst., Spec. Rep. 4, 95 p.

NARDO, G. D.
1832. [*Skeponopodus*]. Ber. ü Verein. Dtsch. Naturf. u. Aerzte:99.
1833. De Skeponopodo novo piscium genere et de Guebucu marcgravii, specii illi cognata. Isis (Oken) 26:415-419.

OVCHINNIKOV, V. V.
1970. Mech-ryba i parusnikovye (Atlanticheskii okean. Ekologiya i funktsional'naya morfologiya) (Swordfishes and billfishes in the Atlantic Ocean. Ecology and functional morphology). Atl. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr., Kaliningrad, 106 p. (Translated by H. Mills, Israel Program Sci. Transl., 1971, 77 p.; available U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA, as TT 71-50011.)

PENRITH, M. J., and F. H. TALBOT.
1973. An additional billfish (Pisces: Istiophoridae) recorded from South Africa. S. Afr. J. Sci. 69:121-122.

PERONACI, A.
1966. La fauna delle acque interne e del mare. Editoriale Olimpia, Firenze, 537 p.

RADCLIFFE, L.
1926. "Opah" and "Skilligalee" landed at Boston Fish Pier. Copeia 151:112.

RAFINESQUE, C. S.
1810. Caratteri di alcuni nuovi generi e nuove specie di animali e piante della Sicilia con varie osservazioni sopri i medesimi. Palermo, 105 p.

RIVAS, L. R.
1956. Definitions and methods of measuring and counting in billfishes (Istiophoridae, Xiphiidae). Bull. Mar. Sci. Gulf Caribb. 6:18-27.

ROBINS, C. R.
1958. Check list of the Florida game and commercial marine fishes including those of the Gulf of Mexico and the West Indies, with approved common names. Fla. State Board Conserv., Educ. Ser. 12:1-44.
1974. The validity and status of the roundscale spearfish, *Tetrapturus georgei*. *In* R. S. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Lailua-Kona, Hawaii, 9-12 August 1972, Part 2. Review and Contributed Papers, p. 54-61. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF 675.

ROBINS, C. R., and D. P. DE SYLVA.
1960. Description and relationships of the longbill spearfish, *Tetrapturus belone*, based on Western North Atlantic specimens. Bull. Mar. Sci. Gulf Caribb. 10:384-413.
1963. A new western Atlantic spearfish, *Tetrapturus pfluegeri*, with a redescription of the Mediterranean spearfish *Tetrapturus belone*. Bull. Mar. Sci. Gulf Caribb. 13: 84-122.

SMITH, J. L. B.
1956. Swordfish, marlin and sailfish in South and East Africa. Rhodes Univ., Dep. Ichthyol., Ichthyol. Bull. 2:25-34.

SPRINGER, V. G., and H. D. HOESE.
1958. Notes and records of marine fishes from the Texas coast. Tex. J. Sci. 10:343-348.

SWANN, R.
1957. A stranger takes the bait. Tex. Game Fish 15(12): 15, 28.

TORTONESE, E.
1962. Osservazioni comparative intorno alla ittiofauna del Mediterraneo e dell'Atlantico occidentale (Florida e Isole Bahamas). Natura (Milano) 53:1-20.

UEYANAGI, S., S. KIKAWA, M. UTO, and Y. NISHIKAWA.
1970n Distribution, spawning, and relative abundance of billfishes in the Atlantic Ocean. [In Jap., Engl. synop.] Bull. Far Seas Fish. Res. Lab. (Shimizu) 3:15-55.

UEYANAGI, S., and H. WATANABE.
1965. On certain specific differences in the vertebrae of the istiophorids. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 22:1-8.

VERANY, J. B.
1847. Aggiunto al catalago de' pesci della Liguria. Ottava Riunione Sci. Ital. Genoa 1846:492-494.

WISE, J. P., and C. W. DAVIS.
1973. Seasonal distribution of tunas and billfishes in Atlantic. U.S. Dep. Commer., NOAA Tech. Rep. NM SSRF 2, 24 p.

Synopsis on the Biology of the Shortbill Spearfish, *Tetrapturus angustirostris* Tanaka, 1914 in the Indo-Pacific Areas

SHOJI KIKAWA[1]

IDENTITY

1.1 Nomenclature

1.11 Valid name

etrapturus angustirostris Tanaka, 1914 (Fig. 1).

ginal description: Tanaka, Shigeho. 1914-1915. 'igures and descriptions of the fishes of Japan. Dai-:hi Shoin, Tokyo 18:295-318 (1914); 19:319-342 1915).

1.12 Objective synonymy

ll synonyms that appear in the literature seem to subjective (See 1.21 below).

1.2 Taxonomy

1.21 Affinities

Suprageneric

Phylum Chordata
Subphylum Vertebrata
Superclass Gnathostomata
Class Osteichthyes
Subclass Actinopterygii
Order Perciformes
Suborder Xiphioidei
Family Istiophoridae

Generic

enus *Tetrapturus* Rafinesque, 1810.

ginal description: Rafinesque, C. S. 1810. arratteri di alcuni nuovi generi e nuove specie di nimali principalmente di pesci e piante della Sicil-a, con varie osservazioni sopra i medesimi. 105 p. alermo.

he generic concept of *Tetrapturus* has been a con-ed one. Hirasaka and Nakamura (1947) divide ophoridae into two subfamilies, Tetrapturinae and rlinae. They place *Tetrapturus, Istiophorus,* and *ikia* in the former, and *Marlina* and *Eumakaira* in the latter. In *Tetrapturus,* they include a single species, *T. angustirostris,* and state that *T. belone* in the Mediterranean Sea and *T. brevirostris* in the Indian Ocean may be the same species, or very closely related. Their genus *Kajikia* embraces two species, *K. mitsukurii* (Jordan and Evermann), the Indo-Pacific striped marlin, and *K. formosana,* new species, which was shown later by Ueyanagi (1957) to be the juvenile form of the striped marlin.

Matsubara (1955) disagrees with Hirasaka and Nakamura (1947) and states that the number of vertebrae and other diagnostic characters they used to establish the subfamilies do not deserve a generic rank. Matsubara also considers *Makaira* as valid and excludes *Kajikia, Marlina,* and *Eumakaira* as synonyms of *Makaira.* According to his classification of Japanese fishes, *Tetrapturus* includes *T. angustirostris,* and *Makaira* includes four species: *M. mitsukurii* (Jordan and Evermann), the Indo-Pacific striped marlin; *M. formosana,* now a synonym of *M. mitsukurii; M. mazara* (Jordan and Snyder), the Indo-Pacific blue marlin; and *M. marlina* Jordan and Hill, the black marlin.

LaMonte (1955) states that *Tetrapturus* can be distinguished from *Makaira* by its much more slender body, shorter spear, shorter pectorals, lower lobed and evenly high dorsal, and smaller caudal span. She recognizes *T. belone* Rafinesque, the Sicilian fish, and *T. angustirostris* Tanaka as valid species of the genus.

Royce (1957), who analyzed morphometric data from specimens taken in the central Pacific Ocean, states that the variation in the height of the dorsal lobe and the length of the pectoral fins is such that these characters are not useful in distinguishing one species from another.

With regard to Tetrapturinae and Marlinae as set forth by Hirasaka and Nakamura (1947), Robins and de Sylva (1960, 1963) support this basic dichotomy but state that these subdivisions need not be accorded subfamily rank. To such definitions, they add the character of negative allometry of the snout in the "Tetrapturinae" and positive allometry in the "Marlinae," but deny the aid of the pectoral fins for generic definition because this character, mainly on the basis of which LaMonte separates *Tetrapturus* from *Makaira,* shows marked allometry in the spearfish. Robins and de Sylva state that *T. angustirostris,*

[1] Far Seas Fisheries Research Laboratory, Shimizu, Japan.

Figure 1.—*Tetrapturus angustirostris* Tanaka, 1914 (from Hirasaka and Nakamura, 1947).

T. belone, T. pfluegeri, T. albidus, and *T. audax* form a series of equally distinct species. They also point out that the placement of *T. albidus* and *T. audax* in *Makaira* is not in accordance with their relationship because they must be more closely associated with *T. angustirostris* and *T. belone* and indeed with *Istiophorus* than with any of the Makairinae (= Marlinae). These authors feel that the characters separating the above five species do not deserve even subgeneric rank and conclude that all should be placed in the genus *Tetrapturus.*

Morrow (1964) agrees with Robins and de Sylva (1960, 1963) in placing *T. angustirostris* nearer to the Atlantic and Pacific billfishes, the striped marlin, and the white marlin than to the blue marlin and the black marlin in the phylogeny of the Istiophoridae.

The basic dichotomy in recognizing the Istiophoridae, viz. the *Tetrapturus* type and *Makaira* type (*Makaira* as constituted by Robins and de Sylva, 1960), is also supported by Ueyanagi (1963b). He confirms that, for the growth of the snout, *Tetrapturus* and *Istiophorus* have a positive allometry during the larval stage[2] and a negative allometry after the young stage,[2] while *Makaira* has a positive allometry after the young stage. The same subdivision, according to Ueyanagi, is also true of the morphological change in the dorsal fin.

Howard (in Howard and Ueyanagi, 1965) mentions that Nakamura's divisionary characters may have some phylogenetic value and feels that the hierarchy proposed by Robins and de Sylva, which takes into consideration the only presently known characters which may have phylogenetic significance, is the most usable and practical offered so far.

Nakamura, Iwai, and Matsubara (1968) include three genera, *Istiophorus, Tetrapturus,* and *Makaira,* in the Istiophoridae. They separate *Istiophorus* from the other two by its much higher, saillike first dorsal fin and considerably elongated ventral fins with well-developed fin membrane. The distinguishing characters between *Tetrapturus* and *Makaira* are t height of the first dorsal fin, the degree in lateral co pression of the body and the dorsal profile from t preocular region to the first dorsal origin. They pla in *Tetrapturus* not only the spearfish group of t world, but also the Indo-Pacific striped marlin a white marlin, and thus agree with Robins and Sylva (1960, 1963).

The generic concept of *Tetrapturus* has, as se above, changed from what originally referred only the spearfish group to where it now includes spec ranging from the shortbill spearfish to the Inc Pacific striped marlin. Most authors have used t form of the dorsal fin and the nature of the ventral fi in separating *Istiophorus* from *Tetrapturus.* T dichotomous system discussed above puts *Istiophor* nearer to *Tetrapturus* rather than to *Makaira,* but appears that there is current agreement that these characters deserve generic rank although the form the first dorsal fin changes considerably with growt

The diagnostic characters of *Tetrapturus,* especi ly for adults, given by Nakamura et al. (1968) are follows:

> The first dorsal fin is slightly higher than the body depth. T ventral fin rays are rather long, its fin membrane not well dev oped. The body is laterally compressed. The dorsal profile straight from the preocular region to the origin of the first dor fin, except for the striped marlin and the white marlin. The c nium is narrow in width and slender. The neural and haer spines on the central vertebrae nearly form a parallelogram. T number of vertebrae are 12 + 12 = 24. The lateral apophysi not well developed.

Specific

Species *Tetrapturus angustirostris* Tanaka, 1914. (Fig. 1)

(Original combinati

Identity of type specimen:

Length: 2,000 mm measured from tip of spear upper lobe of caudal.

[2]For these definitions, see Ueyanagi (1963a).

)eposited at: Science College Museum, Tokyo University.
'atalog No.: 4187.
teported by: Shigeho Tanaka.

pe locality: Sagami Bay, southern coast of central Japan.

ıgnosis:

1. The snout is very short and the body deeply compressed.
2. The first dorsal fin at the anterior lobe is nearly the same as the body depth, then being evenly high posteriorly.
3. The second anal fin is situated anterior to the second dorsal fin.
4. The pectoral fins are narrow and short.
5. The anus is situated far anterior to the origin of the first anal fin.
6. The number of first dorsal spines ranges from 46 to 50.
7. A paired gonad is asymmetrical and Y-shaped, the left lobe is much longer.

bjective synonymy:

Tetrapturus illingworthi Jordan and Evermann, 1926.
Tetrapturus kraussi Jordan and Evermann, 1926.
Tetrapturus brevirostris Fowler, 1928.
Psudohistiophorus angustirostris de Buen, 1950.
Psudohistiophorus illingworthi de Buen, 1950.
Tetrapturus indicus Deraniyagala, 1951.

tificial key to genus: The following key to the ecies of *Tetrapturus* is based on Nakamura et al. 968), who constructed a key to all the species of the phiidae and Istiophoridae.

First dorsal fin is rather high at the anterior part, being nearly at the same height to the posterior. Anus situated far anterior to the origin of first anal fin. Second anal fin situated anterior to the second dorsal fin.

b_1 Pectoral fins narrow and short . . . . . *T. angustirostris* Tanaka

c_1 Snout remarkably short . . . . . *T. belone* Rafinesque

b_2 Pectoral fins broad and long . . . . . *T. pfluegeri* Robins and de Sylva

First dorsal fin is slightly higher than, or nearly the same as, body depth at the anterior part, then gradually lowered. Anus situated just anterior of the origin of first anal fin. Second anal fin and second dorsal fin nearly opposite.

d_1 Pectoral fins broad with rounded tip. Tip of first dorsal fin and first anal fin roundish . . . . . *T. albidus* Poey

d_2 Pectoral fins narrow with pointed tip. Tip of first dorsal fin and first anal fin pointed . . . . . *T. audax* (Philippi)

1.22 Taxonomic status

Recent morphometric studies on external characters of *T. angustirostris* suggest that there may be clinal or population variations in some of these characters among specimens from different oceans. It seems natural, therefore, to think it polytypic (see 1.31).

1.23 Subspecies

None.

1.24 Standard common names and vernacular names

Country	*Standard common name*	*Vernacular name*
Japan	Furaikajiki	Sugiyama Sanmakajiki
United States	Shortbill spearfish	Shortnose spearfish Slender spearfish
Latin America	Pez aguja corta	

1.3 Morphology

1.31 External morphology

Generalized: The billfishes are generally large and elongate, and are usually characterized by a long projected spearlike snout (the snout is as long as 30% of the eye-fork length or more, except for the shortbill spearfish, which is the only species with an especially short snout). The ventral fins are more or less diminished. Other fins, especially the caudal fin, are well developed. The form of the first dorsal fin varies markedly according to the species. The vertebral bones are peculiar in structure with well-developed platelike neural and haemal spines and characteristically extended anterior neural and anterior haemal zygapophyses. The number of vertebrae are 24 (12 + 12 or 11 + 13) (Ueyanagi, 1963b).

The external morphology of *T. angustirostris* as described by Nakamura et al. (1968) is as follows:

ID, XLVII-L; IID, 6-7; IA, 12-15; IIA, 6-7; P, 18 - 19; V, I, 2. Body elongate and very laterally compressed (body length about 16.1-22.2 times body width). Body height remarkably low (body length about 8.3-10.4 times body height). Snout short (head length about 1.6 times upper jaw length), and round in cross section. Body densely covered with slender, bony scales whose tips are 3-5 cusped. Minute filelike teeth on both jaws and palatine. Lateral line curved above pectoral fin and straight to caudal portion. Head large (body length about 4.2-4.7 times head length). Eyes moderate in size, without ridge over the orbit. Dorsal profile straight from snout to insertion of first dorsal fin. Caudal fin deeply forked, both upper and lower lobes slightly narrow in width. Two lateral keels on either side of caudal peduncle. Pectoral fin somewhat low-situated (head length about 1.6-2.3 times pectoral length), with pointed tip. First dor-

sal fin starting above the posterior margin of preopercle, slightly higher than body height at its anterior end, down at the vicinity of 10th spine, again up to nearly the same height posteriorly and terminating just in front of insertion of second dorsal fin. Second dorsal fin and second anal fin small, nearly the same in size, and the latter situated anterior to the former. First anal fin somewhat small and crescent-shaped. Ventral fin longer than pectoral fin.

Fin membrane of first dorsal fin deep blue in color without spots. Other fins brown or dark brown. Basal part of first dorsal fin and second dorsal fin silvery white. Dorsal part of body deep blue, side somewhat brownish blue and belly silvery white.

Olfactory rosette radial in form, comprising 45-47 olfactory laminas. Gonad asymmetrical, Y-shaped; left lobe remarkably developed. Anus situated far anterior to insertion of first anal fin.

Variation in counts of spines and fin rays as provided in the literature are shown in Table 1.

Nakamura et al. (1968), who examined nine specimens from the northwestern Pacific, southeastern Pacific, and Indian Ocean, give the following ranges of fin ray counts: first dorsal spines, 47-50, second dorsal fin ray, 6-7; first anal spines, 12-15; second anal fin rays, 6-7; and pectoral fin rays, 18-19. These all fall within the ranges given in Table 1.

Merrett (1971) provides some morphometric relations for *T. angustirostris* (Table 2).

Geographic variation: Penrith (1964) suggests that there are geographic variations in pectoral fin length; the length of the pectoral fin forms a cline of increasing length eastward through the Indo-Pacific region. Penrith compared body proportions of a specimen taken in the Indian Ocean off South Africa with those of a Chilean specimen measured by Robins and de Sylva (1960) and noted that the values of these two specimens of *T. angustirostris,* taken at the extremes of the known range, agree very well except for the length of the pectoral fin. The ratio of the length of the pectoral fin to the body length is 9.8% for South African fish of 1,456 mm and 15.0% for Chilean fish of 1,822 mm, respectively. He adds these the value of about 11% for Japanese fish, jud from the plates in Nakamura (1951) and Tan (1914). Penrith is of the opinion that although ther marked allometric growth of the pectoral fins in *pfluegeri* (Robins and de Sylva, 1963), it seems dou ful that there is any comparable allometric growth *T. angustirostris*, because he sees no sign of allomet growth over a range of 1,470-1,857 mm (measu from tip of snout to fork) in the tables given by Roy (1957) for Hawaiian fish. The difference in s between the South African fish and Chilean fish nearly comparable to the above range. This Penrith to believe that the large difference in t ratios of the pectoral fin length to the body leng between these two specimens may be due geographic variation and not to the allometric grow of the pectoral fins. He also shows the following valu for the pectoral fin length as expressed as a percenta of the first dorsal height: West Indian Ocean 76.6 Hawaii (Royce, 1957) 87.4-100.4% with a mean of 9 and in the extreme east, off California 100.6% (Cra 1958), and Chile 100.8% (Robins and de Sylva, 196

Merrett (1971) analyzed the morphometric data *T. angustirostris* from the equatorial western Indi Ocean from the viewpoint of specific identity a compared the results with those from the Paci Ocean given by Royce (1957). His comparison of t mean values of four characters, i.e., the length of t pelvic fin, the anterior height of first dorsal fin to t fork length, the anterior height of first anal fin to t anterior height of the first dorsal fin and the length the 20th dorsal spine, shows that the values are in cases smaller for the Indian Ocean fish. Merrett (19 suggests that these probably reflect populati differences in the species from the Pacific and Indi Oceans, since the same trend is apparent throughou He gives the mean value of the pectoral fin length body length of 11.6% for the six specimens he exa ined. According to Merrett, this tends to confirm t

Table 1.—Individual variation in counts of spines and fin rays of *Tetrapturus angustirostris*.

Spines and fin rays	Counts (upper figure) / Frequency (lower figure)								
Dorsal spines	44	45	46	47	48	49	50	51	52
			3	1	6	5	3	2	
Second dorsal fin rays		5	6	7	8	9			
			7	2					
Anal spines		11	12	13	14	15	16	17	
			2	1	3	1	1		
Second dorsal fin rays		5	6	7	8	9			
			3	6					
Pectoral fin rays		16	17	18	19	20	21		
			2	5	1				

Source: Tanaka, 1914; Robins and de Sylva, 1960; Ueyanagi, 1962; Watanabe and Ueyanagi, 1963; Merrett, 1971.

Table 2.—Various morphometric relations of *Tetraptu angustirostris*.

Measurement or ratio of measurements	No.	Minimum	Maximum	Mean	Standa deviati
Pelvic fin length	5	28.2	34.7	30.0	2.5
Height DI / Fork L.	4	0.117	0.134	0.123	-
Height AI / Height DI	5	0.556	0.621	0.580	0.027
Height 20th ray of DI	6	14.1	16.9	15.0	0.9

pothesis in the geographic variation proposed by nrith (1964).

orphological definition of subpopulations: The owledge of subpopulations is insufficient to estab-h such a definition.

escription of morphological changes which occur ring growth: Ueyanagi (1963b) reports the change the length of the snout throughout growth as shown Figure 2. The relation between snout length and dy length shows a marked positive allometric owth until the larva attains a body length of 20-30 m (measured from posterior margin of eye to end of ntral ray of caudal), then suddenly changes to a gative allometric growth throughout the young and mature stages. The morphological change in the out during the early developmental stages is the st remarkable feature in this species. Generally in e billfishes, the dorsal fin is very large and high over base during the larval stage but, except for its terior lobe, decreases in height in the adult stage. In e shortbill spearfish, however, the dorsal fin is near-the same height posteriorly even in the adult fish. om this, Ueyanagi (1963b) assumes that the form of e first dorsal fin of *T. angustirostris,* although un-own in the immature stage, changes little, and that e change is linear throught its life-span.

1.32 Cytomorphology

No information is available.

1.33 Protein specificity

No information is available.



ure 2.—Relation of snout length to body (eye-fork) length of *trapturus angustirostris* during growth (Ueyanagi, 1963b).

2 DISTRIBUTION

2.1 Total Area

Geographical distribution: The occurrence of the shortbill spearfish is reported from Japan, Taiwan, California, Chile, South Africa, and very wide areas including the equatorial western, central, and eastern Pacific, the South Pacific, and the equatorial western Indian Ocean. This species is not reported to occur in the Atlantic Ocean and the Mediterranean Sea; howevern close relatives are present in these waters.

According to Nakamura (1937, 1949), this species is largely oceanic in distribution and does not enter coastal waters. It occurs in warm waters to the east of Taiwan (30-300 miles offshore). Near Japan, it is distributed in ocean waters south of lat. 35°N, but the density is not high.

Royce (1957) reports the sporadic occurrence of this species in the central Pacific Ocean between lat. 10°N and 10°S; information is based on longline fishing surveys conducted by POFI[3] vessels. In the Hawaiian fishery, according to Royce, the shortbill spearfish is one of the miscellaneous spearfishes that compose only a small fraction of the total billfish catch.

Koga (1959) states that in the South Pacific around the Fiji Islands there is a tendency of slightly increasing hooked rate toward the south between lat. 20° and 30°S.

Howard and Ueyanagi (1965) mention that this pelagic fish is dispersed throughout tropical and subtropical areas of the Pacific and density is always low except in the northwestern Pacific between lat. 15° and 30°N, where the density appears to increase from November through February.

Kume and Joseph (1969a, b) report the occurrence of the shortbill spearfish only in the high-seas area beyond about 600-700 miles from shore in the eastern tropical Pacific Ocean.

According to Merrett (1971), *T. angustirostris* is apparently more abundant during the southeast monsoon period, although this indication is due to high catch rates in only one monsoon period.

Within the area of the Japanese longline fishery which extends over the entire Pacific and Indian oceans between lat. 40°N and 50°S, this species is caught sporadically in areas roughly between lat. 30°N and 30°S. The catch report form used by the Japanese commercial longline fishery however, is such that catch data are not available for the shortbill spearfish and sailfish; these species are combined in a single column of the report. This pooling of the species in the catch record is related to the limitation in the number of columns on the punch card used for preparing the yearly statistics on the Japanese longline

[3]Pacific Oceanic Fishery Investigations; now the Southwest Fisheries Center Honolulu Laboratory, National Marine Fisheries Service, NOAA.



Figure 3.—Distribution of *Tetrapturus angustirostris*, occurrence by 1° square (Howard and Ueyanagi, 1965). Shaded lines fo delimitation from combined shortbill spearfish and sailfish data from Japanese longline fishery in 1970.

fishery. This makes it impossible to separate these species and to delineate areas of their distribution in detail. The approximate geographical range in distribution of *T. angustirostris* is shown in Figure 3.

2.2 Differential Distribution

2.21 Spawn, larvae, and juveniles

Areas of occurrence and seasonal variation: Nakamura (1937) reports that the fish taken in waters adjacent to Taiwan in November released ripe eggs and assumed that spawning of *T. angustirostris* takes place in winter in warm offshore currents with surface water temperature of about 26°C. Fish with ripe ovaries are also reported to occur in the central Pacific Ocean by Royce (1957) and in the western Indian Ocean by Merrett (1971).

Ueyanagi (1962, 1963a, 1964) assumes from the occurrence of larvae and mature fish that spawning of this species is more active in winter than in summer, although fish are reported to spawn in the tropical and subtropical waters between lat. 25°N and 25°S. The distribution of larval fish from the western to the central Pacific Ocean as determined by Ueyanagi is in Figure 4. The larvae were collected from May through February or almost throughout the entire year. The sites of capture of larval fish are concentrated in an area between lat. 15° and 25°N in the western Pacific Ocean, reflecting the increased effort (tows) in this region. From the Indian Ocean, two larval specimens presumably identified as *T. angustirostris* were reported. They were collected at the Dana station 3855 southwest off Sumatra (Jones and Kumaran, 1964). Ueyanagi (1962) provides a comparison of the seasonal occurrence of larvae of the shortbill spearfish with that of the blue marlin in the western subtrop Pacific Ocean (Table 3).

For the western Indian Ocean, Merrett (19 provides some data to indicate spawning of *angustirostris* and states that this species is m abundant during the southeast monsoon season, wh the maturity of females is more advanced and the s face temperatures are at the lowest (mean 25.5°C). mentions that this tends to confirm the suggestion Ueyanagi (1964) that spawning activity is greater d ing the winter months.

2.22 Adult

Areas of occurrence and seasonal and ann variation: No information on annual variation available. For seasonal variation, see 2.1.



Figure 4.—Distribution of larval shortbill spearfish. Bla dots from Ueyanagi, 1963a; open circle from Jones a Kumaran, 1964.

Table 3.—Larval occurrences of shortbill spearfish and blue marlin during winter and summer months in the western subtropical Pacific Ocean (lat. 15°-25°N) (Ueyanagi, 1962).

Season	No. of hauls	Species	Percent occurrence	Average no. per haul
Dec.-Feb.	69	Shortbill spearfish	14.5	0.25
		Blue marlin	4.4	0.16
May-Aug.	351	Shortbill spearfish	8	0.12
		Blue marlin	37.3	1.29

2.3 Determinants of Distribution Changes

To determine the vertical distribution of billfishes in the equatorial Indian Ocean, Merrett (1971) relates the percent catch rates of each species taken by the longline fishery to depth soundings and states that no *T. angustirostris* was caught in waters shallower than 500 fathoms. The highest catch rate for this species was obtained in the 501-1,000 fathom range and the catch rate decresed in deeper waters. Data are too sparse to be really significant, however.

2.4 Hybridization

2.41 Hybrids

No information is available.

2.42 Influence of natural hybridization in ecology and morphology

No information is available.

3 BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

Hermaphroditism, heterosexuality, intersexuality: Heterosexual.

Nature and extent of hermaphroditism: No example of hermaphroditism of *T. angustirostris* has been reported thus far.

Sexual dimorphism: Despite many morphological studies, there is no report dealing with sexual dimorphism. Concerning the difference in size by sexes, Koga (1959) reports that larger fish are more frequently female than male in the size composition of longline catches made in the southeastern Pacific Ocean from September through November. Merrett (1971) states that no limitation of size by sex is observed in the longline catch from the equatorial western Indian Ocean.

3.12 Maturity

Size and weight at sexual maturity: The size of fish in running ripe condition reported thus far are as follows: a 1,524-mm specimen (measured from bill tip to origin of lateral keels on caudal peduncle) from the waters adjacent to Taiwan (Nakamura, 1937), a 1,638-mm specimen (measured from bill tip to caudal fork) from the central Pacific (Royce, 1957), and a 1,390-mm specimen (measured from center of orbit to shortest caudal ray) from the equatorial western Indian Ocean (Merrett, 1971). The specimen reported by Merrett was in the ovulation stage, the maturity stage was VI, and the fish weighed 18.6 kg (41 lb).

3.13 Mating

Monogamous, polygamous, promiscuous: Although the evidence is not certain, the billfishes seem to be promiscuous.

3.14 Fertilization

External.

3.15 Gonads

Merrett (1970) describes the ovulation for *T. angustirostris,* the final phase in the course of oogenesis which follows a point at which the eggs attain 750 μm in size. He states that immediately prior to this, the ovary becomes more vascular and the follicles, both theca externa and granulosa layers, then burst. On bursting, the released eggs swell by the uptake of liquid. The increase in size is a mean diameter of from 845 to 1,442 μm. Clearly, ovulation is not an all or none process, since barely half of the eggs in the ovary are estimated to be shed. Ovulation is observed to take place first centrally, and then radially outwards. His observations further suggest a continuous availability of spermatozoa in mature males by the differential maturation of the testicular lobules in conjunction with the possession of a muscular seminal vesicle. The male cycle appears dissimilar to the cycle in other teleosts while the female cycle is similar to other teleosts, as maturation is synchronous throughout the ovary.

Merrett (1971) estimated the mean fecundity of the Indian Ocean istiophorids and found that considerable variation occurs in the fecundity of *Istiophorus platypterus* and that such variation is apparently related to fish size. Although the relation

between egg number and fish size is not given, he provides the estimates of 6,200,000 and 2,100,000 eggs per spawning for two *T. angustirostris,* both 1,390 mm long (measured from center of orbit to shortest caudal ray). The latter estimate is made on unovulated eggs only.

3.16 Spawning

Nakamura (1938) assumes that spawning of the billfishes probably does not, as in other fishes, take place in large groups over a very short period of time, but probably is continuous over a long period and over a broad area of the sea. The reason for this, he states, is that it is known that in a single area fish with ripe eggs and those with unripe eggs are found mingled together over a period as long as 2 mo.

Nakamura (1943) mentions that spawning of the sailfish in a group takes place as individuals become ripe and that the male fish follows the female at the time of breeding.

3.17 Spawn

Merrett (1970) shows the frequency distribution of the diameters of eggs shed by this species, which ranges roughly from 1.30 to 1.60 mm with a mean of 1.442 mm.

According to Nakamura (1937, 1938), the eggs released from the ovary and preserved in alcohol are spherical and about 1 mm in diameter. The ovarian eggs, which are nearly mature (not fully ripe), are almost colorless and translucent, with slightly yellowish brown oil globules which unite until they grow into two fairly large globules. At this stage, the egg is nearly 0.8 mm in diameter.

3.2 Preadult Phase

3.21 Embryonic phase

No information is available.

3.22 Larvae phase

Morphology of this fish at the postlarval stage (Fig. 5) was studied based on about 90 specimens ranging in total length from 2.5 to 83 mm collected from the tropical and subtropical areas of the Pacific Ocean (Ueyanagi, 1962).

The ratio of the snout length to body length increases until the larva grows to about 50 mm in length. Among billfishes, *T. angustirostris* ranging from 20 to 50 mm in length has the largest snout relative to the body length.

The ratio of the head length to body length increases rapidly until about 7 mm in length, then increases less rapidly up to about 50 mm in length. The ratio of the head length is about 50% at 7 mm and about 60% at 50 mm in length. The ratio seems to continue to decrease gradually afterward. A wide space between the anus and the origin of the first anal fi which is characteristic of this species, is already see in larval fish larger than 20 mm long.

The pterotic and preopercular spines are formed about 4 mm in length. As growth proceeds, this spin tion and the serration over the orbit and on the low edge of the lower jaw are rapidly developed until the eventually undergo degeneration at about 20 mm length. At this stage, the minute spiny scales begin appear on the opercle and the surface of the body. larvae greater than 70 mm, the spiny scales cover th whole surface of the body and the lateral line becom evident. The palatine teeth appear at about 4 mm length. There are 30 or more teeth on the upper jaw fish about 20 mm in length. In larvae larger than 3 mm in length the teeth are beginning to degenerat

The vertical fins are membraneous and not y differentiated in larvae smaller than 4 mm. The ver tral fins are formed in larvae over 4 mm long and a about 7 mm, the dorsal fin and the caudal fin ar clearly differentiated. The fin rays and the hypura plate are also formed. At about 20 mm in length, th full complement of fin rays is reached. As develop ment further advances, the dorsal fin becomes highe and saillike. The pectoral fins and the ventral fin become more elongate and the caudal fin is forked

Melanophores are found on the body from the ver early stage of development. At 2.4 mm in length, th melanophores appear on the head, the dorsal and th lateral sides of the trunk, and the dorsal wall of th body cavity. They are also visible along the urostyle inside the isthmus, on the tip and the mid-part of th ventral side of the lower jaw, and on th branchiostegal membrane. The melanophores on th branchiostegal membrane is characteristic of *T angustirostris* at the larval stage. As developmen proceeds, the pigmentation on the side of the body ex tends posteriorly and ventrally and covers the greate part of the caudal region at about 7 mm in length. Th chromatophores appear on the dorsal fin in larva over 15 mm, and at about 20 mm the whole side of th body is pigmented except for the tip of the hypura plate. A trace of the melanophores on th branchiostegal membrane is visible even in specimen as large as 76 mm in length.

3.23 Adolescent phase

The following description is for a young specime 514 mm in standard length (measured from bill tip t end of hypural plate) reported by Watanabe an Ueyanagi (1963).

The snout is elongate, being 26% as long as the bod length (posterior margin of eye to end of central ray caudal fin). The ratio of the upper jaw length to th lower jaw length is approximately 2:1. There is dense growth of minute conical teeth on both upp and lower jaws, giving a filelike touch. The outer teet are larger and found in rows and the inner teeth ar coarsely implanted. Palatine teeth present. The gi



Figure 5.—Postlarval to young specimens of *Tetrapturus angustirostris*. a. 4.5-mm SL (Ueyanagi, 1963a), b. 13.0-mm SL (Ueyanagi, 1963a), c. 27.2-mm SL (Ueyanagi, 1963a), d. 514.0 SL (Watanabe and Ueyanagi, 1963). SL = standard length: from tip of snout to end of hypural bone.

rs are not formed and, instead, there is a thick th of villiform minute conical teeth on that part ıe gill, giving a rough touch.

ıe height of the first dorsal fin is nearly the same throughout its entire base, but the anterior lobe is not yet formed and the fin rays in the central part are slightly higher compared with those of the adult. First and second dorsal fins are still continuous. Number of

dorsal spines and fin rays are 51 + 6. The grooves in which the first dorsal fin and the first anal fin are folded are formed throughout the entire base of each fin. The length of the pectoral fin nearly equals the height of the body. The ventral fins are considerably elongated, reaching the vicinity of the anus. The ventral groove in which the ventral fins are folded is developed. The spread of the caudal fin is not large compared with that of the adult. The space between the anus and the origin of the first anal fin is large, reaching about 26% of the length of the body cavity.

The body cavity is extremely long, extending to the first anal fin beyond the anus. The arrangement of the viscera is almost identical with that in the adult. Although completely immature, the left lobe of the gonad is longer than the right lobe, and elongated a little beyond a point at which the right lobe joins.

Scales cover the opercle and the whole surface of the body. They are cycloid with less exposed portion and appear to be at the early stage of development. The shape of the scales depends on the location of the scales on the body; those on the dorsal and the ventral parts are slender compared with those on the lateral part of the body. The lateral line is simple: it starts above the pectoral fin, being bent downward and running straight along the lateral median line from a point posterior to the pectoral fin to the caudal peduncle.

Sun (1960) reported on a juvenile specimen he identified as *T. angustirostris* from the Pacific Ocean. Ueyanagi (1962), however, believes this specimen to be the young form of *Eumakaira nigra* (= *Makaira mazara*), the Pacific blue marlin, in the light of features such as the remarkable shortness of the snout, the evenly high dorsal fin throughout its entire base and the form of the lateral line.

3.3 Adult Phase

3.31 Longevity

Nakamura (1938) states that fish over 20 kg are rare. Royce's (1957) largest specimen weighs 51 pounds (about 23 kg) and measures 1,859 mm from the bill tip to the caudal fork. According to the length composition of the South Pacific shortbill spearfish reported by Koga (1959), the largest fish is in the 151- to 155-cm (eye-fork length) class for males and in the 146- to 150-cm class for females. The largest specimen ever reported is a fish of 114 pounds (about 51.8 kg) given in Table 3 of Royce's paper (1957). This fish was landed at the Honolulu market in 1951.

3.32 Hardiness

Nothing has been reported on hardiness. Since billfishes are large and active dwellers of the ocean, their life will no doubt be very short in artificially confined environments. When hooked on longline gear, they often jump up above the sea surface, thrash violently and eventually entangle themselves w branch lines into almost exhaustion before they a landed.

3.33 Competitors

Billfishes, together with tunas, constitute a group apex predators in tropical and subtropical wate They appear to compete with one another in feedin since previous reports (Suda, 1953; Yabuta, 195 Watanabe, 1960; Koga, 1967; Mori, 1967[4]) indica that they have no marked food preference and there no significant differences in their stomach conten Sharks or other kinds of fishes such as dolphin barracudas, Spanish mackerels also probably co pete in feeding with tunas and billfishes.

3.34 Predators

When hooked on longline gear, it is well known th tunas and billfishes are frequently attacked by shar or perhaps killer whales. Although information lacking, sharks and killer whales probably prey tunas and billfishes.

3.35 Parasites, diseases, injuries, and abnormalities

No information is available.

3.4 Nutrition and Growth

3.41 Feeding

Information is scarce on diel feeding habit manner, frequency, and variation of feeding habits relation to environmental and physiological co ditions. Yabuta (1953) notes that different kinds food organisms tend to appear in stomachs of tun and billfishes including *T. angustirostris* as the seaso changes. It has not been determined, howeve whether this difference is due to the difference in fi species from which stomachs were collected or other reasons.

3.42 Food

Types eaten and their relative importance in th diet: The stomach contents of billfishes comprise very large variety of crustacea, mollusca, tunicat pisces, and coelenterata. Of the food items, squid an fishes are the most important in frequency an number (Suda, 1953; Yabuta, 1953; Watanabe, 196 Koga, 1967; Mori, 1967 (see footnote 4); Kog Imanishi, and Tawara, 1972). These reports agree that the billfishes, as do tunas, feed on commonly pre

[4]Mori, K. 1967. Interim report on the investigation of stomac contents of billfishes. Report presented at Tuna Fisheries Researc Meeting, 1966, 5 p.

and easily available organisms in their habitats. itanabe (1960) stated that there is no significant ference between the stomach contents of tunas and lfishes except that billfishes feed more frequently juvenile tunas and billfishes than tunas do. The age organisms which are the most common in the machs of billfishes in the North Pacific Ocean are uid and fishes of the Lepidotidae, Alepisauridae, inaceidae, and Katsuwonidae.
Koga et al. (1972) examined the stomach contents the shortbill spearfish in the central South Pacific ean and pointed out that the number of species of n eaten by this species is limited compared with the iped and blue marlins, although the food of this ecies is almost identical with the other marlins with pect to the cephalopods and crustaceans. They ted that the members of the Myctophidae, iacanthidae, and *Polyipnus* which are considered to deepwater dwellers are lacking in the stomachs of e shortbill spearfish. These authors assume from s that the shortbill spearfish swim in a shallower er of water than do the striped and blue marlins.

lume of food eaten during a given feeding iod: Information is lacking on the absolute ume of food eaten by this species. Koga et al. 72) examined the relative volume of stomach con- ts of tunas and billfishes by classifying it into five egories and mentioned that the shortbill spearfish l yellowfin tuna more often have stomachs filled h food than the albacore, striped marlin, and blue rlin do. They also stated that the shortbill spear- n and albacore tend to eat smaller food items than the other tunas and billfishes.

3.43 Growth rate

lative and absolute growth pattern and rates: The nual growth rate has been estimated for the Atlan- sailfish (de Sylva, 1957), the western Pacific lfish and striped marlin (Koto and Kodama, 1962a; to, 1963) and the black marlin in the South China a (Koto and Kodama, 1962b). Age determination ng hard tissues such as scales, otoliths, and contra difficult to apply to the billfishes (Koto and dama, 1962a). Information on the absolute growth lacking for the shortbill spearfish. (For relative wth, see 1.31.)

3.44 Metabolism

No information is available.

3.5 Behavior

3.51 Migration and local movement

Data available thus far are too limited to consider movement of this species.

3.52 Schooling

Composition of stocks by size, age, and sex: See 4.13.

Mixing between schools: Howard and Ueyanagi (1965) mention that the two areas of greatest density of the Pacific shortbill spearfish appear to correspond with the overlapping borders of distribution of striped and blue marlins, and speculate whether, if more data were available, the same might be found for all such areas of overlap between the distribution of striped and blue marlins. For schooling habit, see 3.16

3.53 Response to stimuli

No information is available. For response when hooked in the longline gear, see 3.32.

4 POPULATION

4.1 Structure

4.11 Sex ratio

According to Nakamura (1944), the sex ratio of the shortbill spearfish and sailfish is nearly 1:1 during the spawning season in waters adjacent to Taiwan. For the swordfish and the striped, blue, and black marlins, the size difference with sex and the seasonal change in sex ratio are noted (Nakamura, 1944; Nakamura, Yabuta, and Ueyanagi, 1953; Ueyanagi, 1953). Koga (1959) states that size difference with sex is also evident for the shortbill spearfish in the South Pacific; the female is larger in average length than the male. The catch data collected during 1966-70, however, suggest that the difference in size between the sexes for this species is not so definite as first expected (Koga et al., 1972). The sex ratio from September through November in the South Pacific, according to Koga's data, is 1:1.54 (M:F). The preponderance of females in the catches of this species is also noted in the equatorial western Indian Ocean, although the longline catches are quite few (Merrett, 1971).

4.12 Age composition

No information is available on age.

4.13 Size composition

Size composition of the shortbill spearfish from the Pacific Ocean is reported by various authors (Royce, 1957; Koga, 1959, 1967; Kume and Joseph, 1969b; Strasburg, 1970; Koga et al., 1972). The size composition of fish from the Indian Ocean is given by Merrett (1971). Although data on size compositions reported thus far are limited in number and fragmental seasonally, there appears to be a difference in size according to localities in the Pacific Ocean. As shown in



Figure 6.—Body (eye-fork) length and weight compositions of shortbill spearfish caught by longline fishery in the central South Pacific Ocean (lat. 16°-28°S, long. 166°E-174°W), November-December 1969. M = male; F = female; white triangle = average (Koga, Imanishi, and Tawara, 1972).



Figure 7.—Size composition of 63 shortbill spearfish captur in the eastern Pacific Ocean during 1963 and 1964 by the F *Shoyo Maru*. Number and location of fish measured are shov by 1° areas. (From Kume and Joseph, 1969b).

Figures 6 and 7, the average length of fish caught by the longline fishery is about 135 cm in the central South Pacific and about 150 cm in the eastern Pacific, suggesting an increasing tendency in size toward the east.

Kume and Joseph (1969b) report the weight-length relation for the shortbill spearfish taken in the Pacific Ocean east of long. 130°W. To describe the relationship, they use the equation $\log_{10} Y = \log_{10} a + \log_{10} X$ and fitted the data by the least squares method, where Y is the weight in kilograms and X the eye-fork length in centimeters. The data employed are of two types, 1) gilled and gutted and 2) whole, both from the Japanese longline fishery. The regression coefficient (b) and the Y intercept ($\log_{10} a$) they obtained are 3.9195 and -7.2239 for the gilled and gutted, and 3.7242 and -6.8146 for the whole, respectively. The number of fish and the size range of specimens examined are 89 fish and 102.4-167.0 cm long for the gilled and gutted, and 19 fish and 128.0-156.0 cm for the whole, respectively. Koga et al. (1972) use the same weight-length relation and obtain the value of 4.5926 as the regression coefficient for the South Pacific shortbill spearfish.

4.2 Abundance and Density (of Population)

4.21 Average abundance

The population size is not yet estimated.

4.22 Changes in abundance

Because of the inadequacy of the catch statistics this species (see 2.1), the changes in abundance ca not be considered at present.

4.23 Average density

Annual mean density for any given area or a given year has not been calculated.

4.24 Changes in density

The apparent relative abundance is low throughou areas of the longline fishery compared with the stripe and blue marlin. In 1958, it ranged from 0.16 fish 0.3 fish per 100 hooks depending on areas of t fishery in the South Pacific between lat. 20° and 30° and between long. 170°E and 170°W (Koga, 1959 Concerning variation with depth, there is an indic tion in the equatorial western Indian Ocean that t highest catch rate for this species was in the 501-1,0 fathoms range (Merrett, 1971; see 2.3). Merrett i dicates the value of approximately 0.03 fish per 1 hooks in this depth range.

4.3 Natality and Recruitment

4.31 Reproduction rates

No information is available.

4.32 Factors affecting reproduction

No information is available, except for the gene recognition that the transportation, dispersion, a mortality of the larval fish population may be grea

cted by physical and biological factors such as, for mple, the ocean currents, water temperature, food ply, and predation.

4.33 Recruitment

o information is available.

4.4 Mortality and Morbidity

o information is available.

4.5 Dynamics of Population (as a Whole)

nowledge is limited. See 4.24 for information on arent abundance.

4.6 The Population in the Community and the Ecosystem

nowledge is very limited and no particular infor- ion is available.

EXPLOITATION

5.1 Fishing Equipment

5.11 Gear

he tuna longline fishery is essentially responsible all of the commercial shortbill spearfish catch from Indo-Pacific areas. The longline gear has been eloped to catch tunas, billfishes, and sharks that more or less sparsely distributed in a depth range roughly from 50 to 150 m. The form of the gear set in the water is as shown in Figure 8.

A single unit of gear consists of the main line, a number of branch lines, each with a hook, the float line and the buoy. The depth at which the hook hangs from the main line can be adjusted by the length of the float line. The construction of the gear differs depending on the species of fish sought, the conditions of fishing grounds and the efficiency of fishing vessels. When aiming mostly at the albacore or a species of salmon shark, as seen in Table 4, the gear ordinarily used by the fishermen has relatively short branch lines with short spacing between them. In such a construction the branch lines in a single unit of gear tend to increase in number. With increasing size of the species of fish sought such as yellowfin and bigeye tunas, the length of the branch lines and the spacing between them tend to increase and the number of the branch lines in a single unit tends to decrease. This appears to come from the fishermen's experience in improving fishing efficiency by considering the density of schools of the fish sought, and in preventing the gear from tangling while it is in the water or is being retrieved. The relation between the length of the branch lines and the spacing between them is given in Figure 9.

The length of the main line tends to increase according to the size of the vessels and the power of a longline hauler rather than to the size of fish sought.



ıre 8.—An example of the longline gear: A, buoy; B, main line; C, branch line. (Katsuo-Maguro Nenkan, 1969).



Figure 9.—Relation between length of branch line and distance between branches (Katsuo-Maguro Nenkan, 1969).

Table 4.—Longline gear construction by the species of fish sought (Katsuo-Maguro Nenkan, 1969).

Fish sought	No. of branch lines	Distance between branches	Length of branch lines	Hook size	Length of main line (single unit)
		m	*m*		*m*
Albacore, salmon shark	6-9	20-40	12-18	#34-36	140-360
Yellowfin tuna	5-6	40-50	20-22	#36-38	240-360
Bigeye tuna	4-5	50-60	22-25	#37-38	250-360

The adjustment of fishing depth is achieved by slackening the main line while setting the gear as well as by adjusting the length of the float line. With the float line of 20-40 m and the slackening rate of the main line of 20-50%, the adjusted depth of hooks appears to range from 60 to 150 m (Katsuo-Maguro Nenkan, 1969).

Recently, much effort has been directed at developing labor-saving devices and devices for improving fishing efficiency in the Japanese longline fishery. This effort has led to the gradual improvement of devices within the actual fishery and the development of essentially different methods which may represent breakthroughs in fishing technology. The former is represented by the introduction of two similar devices of longlining, the reel-type and the tub-type gear with improvement in related equipment such as the gear-transporting conveyor and guide stand system, and the latter, for example, by the development of the so-called wire fishing or automatic snapping (Katsuo-Maguro Nenkan, 1969).

5.12 Boats

Most Japanese tuna longline vessels which have recently been built are in the 240-ton class. Table 5 gives the specification of larger vessels in the 340-ton class. Recently, many of these vessels have be equipped with refrigerators capable of preserving t catch at very low temperatures of 40° or -45°C.

5.2 Fishing Area

See 2.1, 2.21, 2.22, 2.3.

5.3 Fishing Season

See 2.1, 2.21, 2.22, 2.3.

5.4 Fishing Operations and Results

5.41 Effort and intensity

Fishing effort for billfishes can practically be e pressed by the fleet size of the tuna longline fishery c in more detail, by the number of hooks used. For t shortbill spearfish, however, this has never been e timated. To assess the fishing effort exerted on a give species of fish, information is first required on t catches of that species and its inherent geographic pattern of density in addition to the crude total effo In this sense, the improvement of data format fro commercial vessels for preparing the catch statisti

Table 5.—Specification of 340-ton class tuna longline vessels.

Items	Vessel 1	Vessel 2	Vessel 3	Vessel 4
Gross ton T	339.79	338.87	338.77	238.68
L (m)	43.74	43.00	43.00	42.80
B (m)	7.80	7.80	7.70	8.00
D (m)	3.80	3.80	3.75	3.80
L/B, L/D, B/D	5.61, 11.51, 2.05	5.51, 11.31, 2.05	5.58, 11.46, 2.05	5.35, 11.26, 2.11
LBD, T/LBD	1,297, 0.26	1,274, 0.27	1,241, 0.27	1,301, 0.26
Fish hold (m^3)	440.4	400.2	345.6	402.9
Freezing room (m^3)	72.2	72.0	87.5	73.5
Freezing capacity/day (t)	13.0	10.0	11.7	7.0
Fuel oil tank (m^3)	230.0	184.9	203.5	200.8
Fresh water tank (m^3)	28.0	21.5	22.3	24.4
Main engine:				
Model	4.D.S.	4.D.S.	4.D.S.	4.D.S.
Rating horse power (PS)	750	750	800	800
Rating revolutions	320	320	330	330
Cylinder (mm)	6 × 370 × 520	6 × 370 × 520	6 × 350 × 500	6 × 350 × 500
Propeller	4 × 1,980 × 1,250	4 × 2,060 × 1,195	4 × 2,000 × 1,240	4 × 2,050 × 1,175
Auxiliary engine:				
Model number	4.D×2	4.D×2	4.D×2	4.D×2
Rating horse power	130	130	130	125
Rating revolutions	720	720	720	750
Generator	100KVA × 230KVA × 1	100KVA × 2	100KVA × 2 30KVA × 1	100KVA × 2 25KVA × 1
Refrigerator	NH_3 26.7R.T × 3	NH_3 27.5R.T × 3	NH_3 44RT × 1 40R.T × 1	NH_3 26.7R.T × 3
Wirless telegram	200W	250W	250W	250W
Direction finder	1	1	1	1
Fish finder	1	1	1	1
Radar	1	1	1	1
Loran	1	1	1	1
Gyro	1	1	1	1

Japan must be made for the shortbill spearfish (see).

5.42 Selectivity

he longline gear takes fish distributed at a depth ge roughly from 50 to 150 m, while the pole-and- and the purse seine gears catch those living near surface. For tunas, this means that the longline- ery exploits larger fish than does the surface ery.

oncerning the effect of selective properties of the gline gear on growth studies, Suzuki (1971) set h the following hypothesis and examined the sibility of gear selectivity in connection with Lee's nomenon: If a fishery exploits fish within a certain range, the fishery tends to catch larger in- iduals out of the young fish group whose size is r the lower limit of the selective range of the gear smaller ones out of the old fish group whose size is r the upper limit. The catches, therefore, tend to resent larger members for the young group and aller ones for the old group compared with fish ac- lly distributed in the sea. Applying this to the gline fishery, samples from the catch would result erroneously larger estimates of growth coefficient smaller estimates of asymptotic length.

PROTECTION AND MANAGEMENT

rotection and management are the most impor- t and urgent issues we now face in the world tuna ources. Presently, little attention is paid to the rtbill spearfish stock(s) because it is less important he commercial and sport fisheries. This, however, s not appear to fix the relative value of the short- spearfish in the future management of the fishery.

POND FISH CULTURE

Not applicable.

LITERATURE CITED

AIG, W. L.
1958. The shortnose spearfish *Tetrapturus angustirostris* Tanaka. A species new to the near shore fauna of the eastern Pacific Ocean. Calif. Fish Game 44:275-280.

BUEN, F.
1950. Contribución a la ictiología. 3. La familia Istiophoridae y descripción de una especie uruguaya (*Makaira perezi* de Buen). Publ. Cient. Serv. Oceanogr., Montevideo 5:163-178.

RANIYAGALA, P. E. P.
1951. The Istiophoridae and Xiphiidae of Ceylon. Spolia Zeylan. Bull. Natl. Mus. Ceylon 26:137-142.

SYLVA, D. P.
1957. Studies on the age and growth of the Atlantic sailfish, *Istiophorus americanus* (Cuvier), using length-frequency curves. Bull. Mar. Sci. Gulf Caribb. 7:1-20.

FOWLER, H. W.
1928. The fishes of Oceania. Mem. Bernice P. Bishop Mus. 10, 540 p.

HIRASAKA, K., and H. NAKAMURA.
1947. On the Formosan spearfishes. Bull. Oceanogr. Inst. Taiwan 3:9-24.

HOWARD, J. K., and S. UEYANAGI.
1965. Distribution and relative abundance of billfishes (Istiophoridae) of the Pacific Ocean. Univ. Miami Inst. Mar. Sci., Stud. Trop. Oceanogr. 2, 134 p.

JONES, S., and M. KUMARAN.
1964. Distribution of larval billfishes (Xiphiidae and Istiophoridae) in the Indo-Pacific with special reference to the collections made by the Danish Dana Expedition. Proc. Symp. Scombroid Fishes, Part 1, p. 483-498. Mar. Biol. Assoc. India, Mandapam Camp.

JORDAN, D. S., and B. W. EVERMANN.
1926. A review of the giant mackerel-like fishes, tunnies, spearfishes and swordfishes. Occas. Pap. Calif. Acad. Sci. 12:1-113.

KATSUO-MAGURO NENKAN.
1969. Katsuo-Maguro Nenkan (Skipjack-Tuna Yearbook). [In Jap.] Suisan Shinchosha, Tokyo, 950 p.

KOGA, S.
1959. Fishing condition of tunas and marlins in the western South Pacific Ocean. [In Jap., Engl. abstr.] Bull. Fac. Fish. Nagasaki 8:57-79.
1967. Studies on the fishery biology of the tuna and marlin in the Indian and the South Pacific Ocean. [In Jap., Engl. summ.] J. Shimonoseki Coll. Fish. 15:51-256.

KOGA, S., H. IMANISHI, and S. TAWARA.
1972. A fishing condition of the tuna and marlin in the central South Pacific. [In Jap., Engl. abstr.] J. Shimonoseki Coll. Fish. 20:213-233.

KOTO, T.
1963. Some considerations on the growth of marlins, using size frequencies in commercial catches. III. Attempts to estimate the growth of striped marlin, *Tetrapturus audax* (Philippi) in the Western North Pacific Ocean. [In Jap.] Rep. Nankai Reg. Fish. Res. Lab. 17:63-85.

KOTO, T., and K. KODAMA
1962a. Some considerations on the growth of marlins, using size-frequencies in commercial catches. I. Attempts to estimate the growth of sailfish. [In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. Lab. 15:97-107.
1962b. Some considerations on the growth of marlins, using size-frequencies in commercial catches. II. Attempts to estimate the growth of so called white marlin, *Marlina marlina* (J. and H.) [In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. Lab. 15:109-126.

KUME, S., and J. JOSEPH.
1969a. The Japanese longline fishery for tunas and billfishes in the eastern Pacific Ocean east of 130°W, 1964-1966. [In Engl. and Span.] Bull. Inter-Am. Trop. Tuna Comm. 13:277-418.
1969b. Size composition and sexual maturity of billfish caught by the Japanese longline fishery in the Pacific Ocean east of 130°W. [In Engl., Jap. summ.] Bull. Far Seas Fish. Res. Lab. (Shimizu) 2:115-162.

LAMONTE, F. R.
1955. A review and revision of the marlins, genus *Makaira*. Bull. Am. Mus. Nat. Hist. 107:323-358.

MATSUBARA, K.
1955. Fish morphology and hierarchy. Part I. Ishizaki-Shoten, Tokyo, 789 p.

MERRETT, N. R.
1970. Gonad development in billfish (Istiophoridae) from the Indian Ocean. J. Zool. (Lond.) 160:355-370.
1971. Aspects of the biology of billfish (Istiophoridae) from the equatorial western Indian Ocean. J. Zool. (Lond.) 163:351-395.

MORROW, J. E.
1964. Marlins, sailfish and spearfish of the Indian Ocean. Proc. Symp. Scombroid Fishes, Part 1, p. 429-440. Mar. Biol. Assoc. India, Mandapam Camp.

NAKAMURA, H.
1937. Habit of the spearfishes of Formosan waters with special reference to the shortbill spearfish. [In Jap.] Zool. Mag. 49:233-238.
1938. Report of an investigation of the spearfishes of Formosan waters. [In Jap.] Rep. Taiwan Government General Fishery Experiment Station, No. 10. (Translated by W. G. Van Campen, 1955; U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 153, 46 p.)
1943. Tunas and billfishes. Kaiyō no kagaku (Science of the sea). J. Oceanogr. Soc. Jap. 3:445-459.
1944. Studies of the spearfishes of Formosan waters. 7. Sexual difference in size. Bull. Soc. Nat. Hist. Taiwan 34 (250):242-246.
1949. The tunas and their fisheries. [In Jap.] Takeuchi Shobōn, 118 p. (Translated by W. G. Van Campen, 1952, U.S. Dep. Int., Fish. Wildl. Serv., Spec. Sci. Rep. Fish. 82, 115 p.)
1951. Tuna longline fishery and fishing grounds. Association of Japanese Tuna Fishing Cooperatives, Tokyo. (Translated by W. G. Van Campen, 1954, U.S. Dep. Int., Fish. Wildl. Serv., Spec. Sci. Rep. Fish. 112, 168 p.)

NAKAMURA, H., Y. YABUTA, and S. UEYANAGI.
1953. Relation between the spawning season and the sex-ratio of some fishes of the family Istiophoridae. [In Jap., Engl. abstr.] Contrib. Nankai Reg. Fish. Res. Lab. 1(13), 8 p.

NAKAMURA, I., T. IWAI, and K. MATSUBARA.
1968. A review of the sailfish, spearfish, marlin and swordfish of the world. [In Jap.] Kyoto Univ., Misaki Mar. Biol. Inst., Spec. Rep. 4, 95 p.

PENRITH, M. .
1964. A marked extension of the known range of *Tetrapturus angustirostris* in the Indian Ocean. Copeia 1964: 231-232.

RAFINESQUE, C. B.
1810. Caratteri di alcuni nuovi generi e nuove specie di animali e piante della Sicilia, con varie osservazioni sopri i medesimi. Palermo, 105 p.

ROBINS, C. R., and D. P. DE SYLVA.
1960. Description and relationships of the longbill spearfish, *Tetrapturus belone,* based on Western North Atlantic specimens. Bull. Mar. Sci. Gulf Caribb. 10:383-413.
1963. A new western Atlantic spearfish, *Tetrapturus pfluegeri,* with a redescription of the Mediterranean spearfish *Tetrapturus belone.* Bull. Mar. Sci. Gulf Caribb. 13:84-122.

ROYCE, W. F.
1957. Observations on the spearfishes of the central Pacific. U.S. Fish Wildl. Serv., Fish. Bull. 57:497-554.

STRASBURG, D. W.
1970. A report on the billfishes of the central Pacific Oce
Bull. Mar. Sci. 20:575-604.

SUDA, A.
1953. Juvenile skipjack from the stomach contents of tu
and marlins. [In Jap., Engl. summ.] Bull. Jap. S
Sci. Fish. 19:319-340.

SUN, G.
1960. Lichinki i mal'ki tuntsov, parusnikov i mech-r
(Thunnidae, Istiophoridae, Xiphiidae) tsentral 'noi i
padnoi chasti Tikhogo Okeana [Larvae and fry of tu
sail-fish and sword-fish (Thunnidae, Istiophoridae,
phiidae) collected in western and central Pacific].
Russ.] Tr. Inst. Okeanol. Akad. Nauk SSSR 41:175-1
(Engl. transl. by W. L. Klawe, Inter-Am. Trop. Tu
Comm., 1960, 18 p.)

SUZUKI, Z.
1971. Comparison of growth parameters estimated for
yellowfin tuna in the Pacific Ocean. [In Jap., En
synop.] Bull. Far Seas Fish. Res. Lab. (Shimizu)
89-105.

TANAKA, S.
1914-1915. Figures and descriptions of the fishes of Jap
Daiichi Shoin, Tokyo 18:295-318 (1914); 19:319-3
(1915).

UEYANAGI, S.
1953. The sexual difference of the sizes of marlins.
Jap., Engl. abstr.] Contrib. Nankai Reg. Fish. Res. La
1(14), 5 p.
1957. On *Kajikia formosana* (Hirasaka et Nakamur
[In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. La
6:107-112.
1962. On the larvae of the shortnosed spearfish, *Tetrapt*
us angustirostris Tanaka. [In Jap., Engl. synop.] Re
Nankai Reg. Fish. Res. Lab. 16:173-189.
1963a. Methods for identification and discrimination of t
larvae of five istiophorid species distributing in the Ind
Pacific. [In Jap., Engl. synop.] Rep. Nankai Reg. Fis
Res. Lab. 17:137-150.
1963b. A study of the relationships of the Indo-Pacific isti
phorids. [In Jap., Engl. synop.] Rep. Nankai Reg. Fis
Res. Lab. 17:151-165.
1964. Description and distribution of larvae of five isti
phorid species in the Indo-Pacific. Proc. Symp. Scor
broid Fishes, Part 1, p. 499-528. Mar. Biol. Assoc. Indi
Mandapam Camp.

WATANABE, H.
1960. Regional differences in food composition of the tun
and marlins from several oceanic areas. [In Jap., En
abstr.] Rep. Nankai Reg. Fish. Res. Lab. 12:75-84.

WATANABE, H., and S. UEYANAGI.
1963. Young of the shortbill spearfish, *Tetrapturus angu*
tirostris Tanaka. [In Jap., Engl. synop.] Rep. Nank
Reg. Fish. Res. Lab. 17:133-136.

YABUTA, Y.
1953. On the stomach contents of tuna and marlin from t
adjacent seas of Bonin Islands. Contrib. Nankai R
Fish. Res. Lab. 1(15), 6 p.

Synopsis of the Biology of the White Marlin, *Tetrapturus albidus* Poey (1861)[1]

F. J. MATHER III, H. L. CLARK, and J. M. MASON, JR.[2]

IDENTITY

1.1 Nomenclature

1.11 Valid name

Tetrapturus albidus Poey, 1861 (Fig. 1).

y, Felipe. July 1861 Memorias sobre la historia atural de la Isla de Cuba. Habana, vol. 2, 442 p. ef. p. 237-243, 258-260; pl. 15, fig. 1, 3; pl. 16, figs. -13; pl. 17, figs. 1-11, 21, 26).



Figure 1.—White marlin, *Tetrapturus albidus* Poey, 1860. (From Nakamura et al., 1968, Fig. 19.)

1.12 Objective synonymy

No type specimen exists, therefore there are no ective synonyms.

1.2 Taxonomy

1.21 Affinities

Suprageneric

Phylum Chordata
Subphylum Vertebrata
Superclass Gnathostomata
Class Osteichthyes
Subclass Actinopterygii
Order Perciformes
Suborder Xiphioidei
Family Istiophoridae

Generic

Genus *Tetrapturus* Rafinesque 1810.

Rafinesque, C. S. 1810, "Caratteri di alcuni nuovi generi e nuove specie de animale e piante della Sicilia." Palermo, 105 p., 20 pls. (ref. p. 54-55, pl. 1, Fig. 1). Type species; *Tetrapurus belone* Rafinesque, 1810.

We follow the generic concept of C. R. Robins and D. P. de Sylva, 1960, "Description and relationships of the longbill spearfish, *Tetrapturus belone*, based on western North Atlantic specimens," Bulletin of Marine Science of the Gulf and Caribbean, 10:383-413, 5 figs. (ref. mat. p. 383, 404-406, Fig. 5), and 1963, "A new western Atlantic spearfish, *Tetrapturus pfluegeri*, with a redescription of the Mediterranean spearfish *Tetrapturus belone*," Bulletin of Marine Science of the Gulf and Caribbean, 13:84-122, 5 figs. (ref. mat. p. 101-102).

The following are transcribed from Robins and de Sylva (1960), page 385:

Contribution No. 3110 from the Woods Hole Oceanographic itution, Woods Hole, MA 02543.
Woods Hole Oceanographic Institution, Woods Hole, MA 02543.

Tetrapturus Rafinesque

Tetrapturus Rafinesque, 1810: 54-55 (*T. belone,* type species by monotypy).

Skeponopodus Nardo, 1832: 99 (*nomen nudum*); 1833: 415-419 (*S. typus*, [= *T. belone* Rafinesque], type species by virtue of the name *typus*).

Tetrapterurus Bonaparte, 1841: 19 (emended spelling).

Tetrapterus Agassiz, 1843: 7, 89-92, table E (emended spelling).

Tetraplurus Vérany, 1847: 492-494 (misprint for *Tetrapturus?*).

Scheponopodus Canestrini, 1872: 112 (emended spelling).

Tetraperus Radcliffe, 1926: 112 (misprint for *Tetrapturus*).

Marlina Grey, 1928: 47 (*Tetrapturus mitsukurii* Jordan and Snyder,] = *T. Audax* Philippi], type species by monotypy; the use of *Marlina* at the generic level is probably a slip).

Kajikia Hirasaka and Nakamura, 1947: 13-14 (*Kajikia formosana*,] = *Tetrapturus audax* Philippi], type species by monotypy).

Pseudohistiophorus de Buen, 1950: 171 (*Tetrapturus illingworthi* Jordan and Everymann] = *T. angustirostris* Tanaka], type species by original designation).

Lamontella Smith, 1956: 32 (*Tetrapturus albida* [sic] Poey, type species by original designation and monotypy).

and p. 402-404:

Relationships.—The limits of the genus *Tetrapturus* are ill-defined in current literature, perhaps due in part to the prior lack of information about the type species *belone*. Nakamura (1949 and English translation, 1952) divided the Istiophoridae into two subfamilies Tetrapturinae and "Marlinae" (=Makairinae). Other workers either overlooked or failed to follow this lead. *Makaira*, as currently constituted, bridges this break and is unnatural; its species are in part assigned to *Tetrapturus* below. We agree with the basic dichotomy suggested by Nakamura although the divisions need not be designated subfamilies. To the definitions of Nakamura (1938, 1949) we add the character of negative allometry of the bill in the "Tetrapturinae" versus positive allometry in the "Makairinae." Also the "Tetrapturinae" are small species, generally much less than 300 lbs. while the two members of the "Makairinae" reach about 2000 lbs. The world-record "striped marlin," a 692 lb. individual taken off Balboa, California, August 18, 1931, is a blue marlin, *M. nigricans*.

LaMonte (1955:325) separated *Tetrapturus* from *Makaira* largely on the basis of the length of the pectoral fin. As already noted, the pectoral fin shows marked allometry in the spearfish. Also, *Tetrapturus*, as here constituted, includes one species (*angustirostris*) that apparently has a short pectoral and several species with very long pectorals; therefore this feature will not aid generic definition. . . .

The relationships of *Tetrapturus* are with *Istiophorus* rather than *Makaira* as here restricted. Our concept of the phylogeny of the Istiophoridae is diagrammed in Figure 5. The species of *Istiophorus* are poorly known but it already appears that few species are represented. The only characters separating *Istiophorus* from *Tetrapturus* are the form of the dorsal fin and the nature of the pelvic fins. Even the character of the dorsal fin breaks down for when all species are considered *T. belone* and *T. angustirostris* are somewhat intermediate. Moreover the young of all included species have high dorsal fins, but this feature is extreme in *Istiophorus* . The structure of the pelvic fin has been insufficently investigated. We choose to retain *Istiophorus* and *Tetrapturus* as genera but a case could be made for ranking them as subgenera.

Figure 5 of Robins and de Sylva (1960) is reproduced here as Figure 2.



Figure 2.—Phylogenetic relationships of fishes of the fam Istiophoridae. Names in quotations are not employed by present writers. (From Robins and de Sylva, 1960, Fig. 5 caption.)

After examining Mediterranean material, Rob and de Sylva (1963) assigned the western Atlan specimens, which they had described as *T. bel* Rafinesque in their 1961 paper to a new species, *pfluegeri* Robins and de Sylva, and redescribed *belone* Rafinesque on the basis of the Mediterrane material which they had examined. Furth statements on the species of *Tetrapturus* and relationship of this genus to the genera *Istiopho* and *Makaira* transcribed from Robins and de Syl 1963, p. 100-102, follow:

. . . Aside from the 12 caudal and 12 precaudal vertebrae t all share, the important characters of the species of *Tetraptu* involve the number of dorsal spines, shape and pigmentatio the spinous dorsal fin, height and shape of the anal fin, lengt the pectoral fin in the adult, position of anus, shape of do profile from the base of the bill to the spinous dorsal, and length . . . .

In their delineation of *Tetrapturus* and placement of it w *Istiophorus* on one branch of the istiophorid dendrogram, Rob and de Sylva (1961: 402) referred to Nakamura (194 Nakamura (1938), though recording the vertebral differen neither defined nor suggested such a subdivision of Istiophoridae. This dichotomy actually dates from the work Hirasaka and Nakamura (1947). These subdivisions are natu but need not be accorded subfamily rank, although so ichthyologists will choose to do so. It was not our intent in earlier paper to suggest such ranking for in small families like Istiophoridae subfamilies are unnecessary and serve no imp tant purpose. These subdivisions are defined as follows (in e instance the first condition applies to that containing *Tetr turus* and *Istiophorus*, the second to that containing *Makaira*. Precaudal vertebrae 12, caudal vertebrae 12 *vs.* precau

ebrae 11, caudal vertebrae 13 (these vertebral differentiations precaudal and caudal elements are clear [see Nakamura, 8 and Pls. 4-5 of 1955 translation] and no variant is known for species). 2. The bill is proportionately longer in juveniles n in adults *vs.* the opposite condition (this is independent of ultimate length of the bill of any species). 3. Size relatively ll, less than 200 lbs., except for *T. audax* which sometimes ex- ds 400 lbs. (as previously noted [Robins and de Sylva, 1961:] the world-record 692-lb. striped marlin is misidentified and eally *Makaira nigricans*, the blue marlin) *vs.* size very large, h species approaching 2000 lbs. 4. Anterior lobe of dorsal fin h, equal to or exceeding the body depth at this point *vs.* erior lobe of dorsal low, less than the body depth at this point.

obins and de Sylva (1960, 1963) assign, as a result eir work, five distinct species to the genus *Tetrap-* s—*T. angustirostris, T. belone, T. pfluegeri, T. lus,* and *T. audax*—with a transition running *Tetrapturus* of the *angustirostris* type to that of *udax* type consecutively. Specific differences are ssed in Robins and de Sylva, 1963, p. 100-102, les 6 and 7.

Specific

e specimen: No type specimen exists.

e locality: North coast of Cuba (Poey, 1861).

gnosis: Fishes of the family Istiophoridae are dis- uished from all others by the extension of the up- jaw into a short pointed spear of nearly round s section, and a pair of small cartilaginous keels ither side of the base of the caudal fin. In the most ely related family, Xiphiidae, the upper jaw is ex- ed into a long sword, much flattened vertically sharp at the horizontal edges; there is a single e fleshy keel on either side of the base of the dal fin, and the ventral fins, present in the phoridae, are lacking.

its adult form, *T. albidus* is easily distinguished the other Istiophoridae. It differs from the *aira* species in: vertebral count—12 + 12 vs. 11 + maximum height of first dorsal fin—more than h of body vs. less than depth of body; body pe—slender (depth less than length of pectoral vs. stout (depth usually greater than length of oral fin); lateral line—distinct, single vs. obscure, le or complex; lobes of first dorsal and first anal —strongly rounded at extremities vs. pointed at emities; spear—long and slender vs. short and t; size—small (less than 100 kg) vs. large (ap- ching 1,000 kg). The above characters are from amura (1949), LaMonte (1955), and Robins and Sylva (1960, 1963). *Tetrapturus albidus* is easily nguished from *Istiophorus* by the size and shape e first dorsal fins. The first dorsal of *T. albidus* is nctly peaked anteriorly, and its height tapers eriorly. Its maximum height is less than twice the depth. The dorsal fin of *Istiophorus* is highest in middle portion of its length, and its maximum ht is much more than twice the body depth.

Tetrapturus albidus may be distinguished from *T. belone, T. pfluegeri,* and *T. angustirostris* by the strongly rounded lobes of its first dorsal and first anal fins, the shape of its dorsal fin, and the location of its anus. The anterior distal extremities of the first dorsal and first anal fins of the latter three species are pointed rather than rounded; their first dorsal fins are not as distinctly peaked anteriorly, and are higher in their middle portion than that of *T. albidus,* making the height of the fin more nearly uniform. The anus of *T. albidus* is near (4.8-5.8% of body length) the anal fin origin, but in the other three species, it is from 6.3 to 11% of the body length from the anal fin origin. The spear of *T. albidus* (26-35% of body length) is longer than that of *T. pfluegeri* (19-25% of body length) and much longer than those of *T. belone* and *T. angustirostris.* These data are from Robins and de Sylva (1963).

Tetrapturus audax is the species most similar in external appearance to *T. albidus.* Although the former is an Indo-Pacific species, the ranges of the two overlap at least off South Africa (Talbot and Penrith, 1962). The main difference noted by LaMonte (1955) was the lateral line—conspicuous in *T. albidus,* and invisible or very inconspicuous in *T. audax.* Robins and de Sylva (1960) separate these species on the basis of the strongly rounded tips of the spinous dorsal, anal, and pectoral fins, and the red flesh of *T. albidus,* and the pointed tips of these fins, and pale flesh, of *T. audax.* The experience of the senior author has been that roundedness vs. pointedness of pectoral fins is a very unreliable character in separating scombroid fishes and that the tip of the spinous dorsal of *T. audax* is not always sharply pointed. The best character in separating these species probably is the shape of the anal fin. In fact, this character easily separates *T. albidus* from all the other Istiophoridae.

The diagnosis of larval and juvenile *T. albidus* is little known. We can only reproduce the key of de Sylva (1963) for young specimens (50-500 mm in length) of the istiophorid species known in the western Atlantic.

A provisional key is presented to assist identification of young specimens of known istiophorid species in the western Atlantic, based upon published data and the writer's material . . .; this key should suffice for specimens from about 50 to 500 mm in length:

1a. Anus placed far forward, approximately midway between bases of anal and pelvic fins; pelvic fins short, not reaching nearly to base of anal fin; dorsal fin unspotted or mottled; dorsal spine count high, 45-53, modally 49 (see Robins and de Sylva, 1963, Table 5) . . . . . Longbill spearfish, *Tetrapturus pfluegeri*

1b. Anus placed close to anal fin base; pelvic fins long, reaching nearly to base of anal fin; dorsal fin variously spotted or mottled; dorsal spine count relatively low, 38-49 2

2a. Lateral line complex, partly forming a reticulated pattern (see Gehringer, 1957, Fig. 25; LaMonte, 1958, Pl. 75, Fig. 2; de Sylva, 1958, Pl. 81, Figs. 1-3; Caldwell, 1962, Fig. 1); bill short, only slightly longer than the mandible . . . . . Blue marlin, *Makaira nigricans*

2b. Lateral line simple, forming a single unit (LaMonte, 1958, Pl. 75, Fig. 3); bill long, greatly exceeding the length of the lower jaw 3

3a. Dorsal fin variously mottled, but without distinct ocelli at its base; anal fin pointed, pale, unmarked; dorsal spine count 40-49, modally 43 or 45; anal spine count 12-17, modally 13; dorsal soft-ray count 6-8, modally 7; anal soft-ray count 6-8, modally 7 (Voss, 1953, Fig. 4 and Gehringer, 1957, Fig. 15; see also Robins and de Sylva, 1963, Table 5, for frequency distribution) . . . . . Atlantic sailfish, *Istiophorus albicans*

3b. Dorsal fin variously mottled, but with four distinct ocelli near the base of dorsal fin (Fig. 1); anal fin broadly rounded, its middle rays dark; dorsal spine count 38-43, modally 42; anal spine count 13-19, modally 14 or 15; dorsal soft-ray count 5-7, modally 6; anal soft-ray count 5 or 6, modally 6 . . . . . .White marlin, *Tetrapturus albidus*

The present authors have identified the specimens of *T. albidus* they have examined by the combination of the above characters, but principally by the strongly rounded lobes of the first dorsal and first anal fins.

Subjective synonymy (from de Sylva, in press):

Tetrapturus belone: Sassi, 1846, Nuovi Ann. Sci. Rend. Sess. Soc. Accad. Sci. Hist. Bologna, ser. 2, VI: 391 (from Genoa; non *T. belone* Rafinesque, 1810). Placed in synonymy of *T. lessonae* Canestrini in Canestrini, 1861.

Tetrapturus lessonae Canestrini, 1861, Arch. Zool. Anat. Fisiol., Genoa, I (k): 259-261, pl. XVII (type locality: Ligurian Sea. Type specimen: a mount in the Museo di Storia Naturale di Genova). Placed in synonymy of *Makaira albida* (Poey) (= *T. albidus* Poey) in LaMonte (1955); reasons discussed.

Tetrapturus imperator: Jordan and Evermann, 1896, Bull. U.S. Natl. Mus., (47), I: 892 (*Tetrapturus albidus* Poey, 1860, considered a junior synonym of *T. imperator* Bloch and Schneider, 1801, which, however, is based upon *Xiphias gladius* Linnaeus, 1758). Placed in synonymy of *T. albidus* Poey in de Sylva (in press); reasons noted above.

Makaira lessonae Jordan and Evermann, 1926, Occas. Pap. Calif. Acad. Sci., XII: 56-57 (new combination based on *Tetrapturus lessonae* Canestrini, 1861). Placed in synonymy of *T. albidus* Poey in de Sylva (in press); reasons noted above.

Makaira albida Jordan and Evermann, 1926, Occas. Pap. Calif. Acad. Sci., XII: 66-67 (new combination based on *Tetrapturus albidus* (Poey, 1860)). Placed in synonymy of *T. albidus* Poey in de Sylva (in press); reasons noted above.

Tetrapturus belone: Legendre, 1928, Bull. Soc. zool. France, LIII: 391-392, fig. 1 (west of English Channel; *non T. belone* Rafinesque, 1810). Placed in synonymy of *Makaira albida* (Poey) (= *T. albidus* Poey) in LaMonte (1955); reasons discussed.

Tetrapturus belone: Desbrosses, 1938, Bull. Soc. zool. France, LXIII: 48-58 (200 miles west-northwest of Groix, Brittany; *non T. belone* Rafinesque, 1810). Placed in synonymy of *Makaira albida* (Poey) (= *T. albidus* Poey) in LaMonte (1955); reasons discussed.

Tetrapturus georgii: Tortonese, 1940, Boll. Mus. Zool. Anat. comp. Torino, XLVIII, ser. 3, no. 115, 3, 5, 6 (based on a mounted specimen in the Turin Museum, cat. no. 784; *non T. georgii* Lowe, 1840). Placed in synonymy of *Makaira albida* (Poey) (= *T. albidus* Poey) in LaMonte (1955); reasons discussed.

Lamontella albida Smith, 1956, Ichthyol. Bull. Rhodes, Univ. 32 (new generic name for *Tetrapturus albidus* Poey, 1 *Lamontella* placed in synonymy of *Tetrapturus* in Robins a Sylva (1960); reasons discussed.

Tetrapturus belone: Lozano Cabo, 1958, Trab. Inst. esp. Ocean XXV: 57, fig. 93 (Ceuta, Spain; *non T. belone* Rafinesque, 1 Placed in synonymy of *T. albidus* Poey, 1860, in Robins a Sylva (1960). Reasons not discussed.

Many additional name changes are listed Nakamura et al. (1968).

Artificial key: Adult fishes of the genus *Tetraptu* modified from Nakamura et al. (1968):

a[1] Anterior fin rays of first dorsal fin fairly l posterior rays about same height; vent situated cidedly anterior to origin of the first anal fin; ond anal fin anterior to second dorsal fin.

b[1] Pectoral fin narrow and short.

c[1] Snout very short ..... Shortbill spearfish *T. angustirostris* Tana

c[2] Snout fairly long ..... M iterranean spearfish *T. Belone* Rafinesq

b[2] Pectoral fin wide and long ..... Long spearfish *T. pfluegeri* Robins and de Sy

a[2] Height of anterior portion of first dorsal fin ab same as the body but as it goes gradual decreas in height towwards posterior end; vent direc anterior to the origin of the first anal fin; secc dorsal fin and second anal fin in parallel positio

d[1] Pectoral fin wide and rounded. The tip the first dorsal fin and first anal rounded..White marlin *T. albidus* Po

d[2] Pectoral fin narrow, and its tip point the tips of the first dorsal fin and first a fin pointed..... ..... Striped marlin *T. audax* (Philipp

1.22 Taxonomic status

The white marlin is a morphological species described and taxonomically understood. Nakam et al. (1968) in their review recognize only one spe of white marlin which is universally accepted in genus *Tetrapturus*.

1.23 Subspecies

We have found no descriptions of subspecies o *albidus*, or any evidence on which subspecies coul based.

1.24 Standard common names and vernac names (Rosa, 1950; Miyake and Hay 1972; de Sylva, in press)

Brazil - Bicuda, Agulhão branca, Espadarte mec british West Indies - White marlin

ıada and France - Makaire blanc
ɔa - Aguja blanca, Cabezona, Blanca, Aguja de
ɔaladar
ɔan - Nishimakajiki, Makajiki
rea - Baeg-sae-chi
rocco - Espadon
tugal - Espadim branco do Atlantico, Espadim
ranco, Agulha, Espadim pequeño
ıth Africa - Wit marlyn (Afrikaans)
ain - Pez aguja, Alfiler, Cometa, Altón
ited States - White marlin, Skilligalee (New
Ingland)
SR - Belyi marlin
ıezuela - Aguja blanca

1.3 Morphology

)ur description of the morphology of *Tetrapturus idus* is based on Poey (1861), LaMonte (1955, 8a, 1958b), Robins and de Sylva (1960, 1963), kamura et al. (1968), Robins (1974), unpublished a collected by Mather and his colleagues, and er sources as cited. Proportions are expressed in cent of body length, with most measurements en as defined by Rivas (1956). Additional asurements were taken as specified. The following ɔreviations from Robins and de Sylva (1960) are d:

D^1 = first dorsal fin
D^2 = second dorsal fin
C = caudal fin
A^1 = first anal fin
A^2 = second anal fin
P^1 = pectoral fin
P^2 = pelvic fin
orig. = origin (in reference to fins)
c.p. = caudal pedencle

e word "ray," if not qualified, is used to designate elements without any implication as to whether y are spines or segmented rays.

1.31 External morphology

he body of the white marlin is of a modified form shape, slender and notably compressed rally (depth: greatest 14-19, near posterior margin of opercle; at orig. A, 12-16; c.p. 3.5-4.2; width: greatest 5.2-9.4, near orig. A; at c.p.[in front of keels] 2.7-3.8). The bill is like a spear, or, as the English name of the fish implies, a marlin spike. Its length is 25-35 from the edge of the eye, or 14-18 from the tip of the mandible. At the latter point, it is 0.81-1.3 deep and 1.2-2.0 wide, and at one third of the distance from the tip of the bill to the tip of the mandible Poey found the depth to be 0.5-0.6 of the width.

The head (24-27) tapers very rapidly from a pronounced dorsal hump over the opercles to a beak-like mouth formed by the bill and the sharply pointed lower jaw. Its dorsal and ventral profiles are concave anteriorly and convex posteriorly. The mouth opening is large (maxillary 15-17), extending beyond the posterior margin of the fleshy orbit by about one-third of its diameter, which is 2.8-3.2. The eye is about midway between the tip of the mandible and the operculum and the preoperculum about midway from the eye to the operculum. The nostrils, each consisting of two openings, are about two thirds of the diameter of the eye ahead of its anterior margin.

The body depth tapers gradually from the opercular area to the first anal fin, then more rapidly to the origin of the caudal.

A pair of horizontal cartilaginous keels (2.7-4.7), one above the other, is located on each side of the caudal peduncle, orginating in nearly the same longitudinal position as the caudal fin. The caudal fin is stiff, powerful, and deeply forked. Its spread is 33-43, and an angle of 80° between the posterior margin (disregarding the tips) and the centerline was reported for the type specimen. Poey lists 12 caudal rays for the type specimen, and Goode (1881) gives a count of 6 + 4 + 4 + 5 for a specimen collected off New Bedford, Mass.

The first dorsal, first anal, and pelvic fins disappear completely into dorsal and ventral groves when not in use. The pectorals fit flush with the sides of the fish in slight depressions in the body, but the second dorsal and second anal fins are not retractable. Poey (1861) described the characteristics and arrangement of the rays of these fins in detail. Their counts are: D_1 - 38-46 (usually 40-43); D_2 - 5-7 (usually 5-6); A_1 - 12-18 (usually 14-16); A_2 - 5-7 (usually 6); P_1 - 17-22 (usually 19-21); and P_2 - 5. Frequency distributions of D_1, D_2, A_1, A_2, and P_1 counts for white marlin from the eastern

Table 1.—Fin-ray counts of western[1] and eastern Atlantic white marlin, *Tetrapturus albidus*.

	Dorsal spines								D_2 rays			Anal spines						A_2 rays			P_1 rays[2]					
	38	39	40	41	42	43	44	45	5	6	7	13	14	15	16	17	18	5	6	7	17	18	19	20	21	22
tern Atlantic	1	3	8	10	11	9	-	-	20	21	1	4	18	18	5	-	-	2	41	-	1	2	6	23	9	-
tern Atlantic	-	1	7	16	19	8	5	1	26	30	-	2	9	28	12	3	2	5	50	1	-	2	10	30	13	1

Data from Robins and de Sylva (1960: Table 1).
Only the left pectoral fin was counted.
ɔurce: Robins, 1974, Table 1.

and western North Atlantic are presented by Robins (1974) (Table 1). The first dorsal fin originates over the posterior margin of the preopercle (first predorsal length 22-25) and extends nearly to the origin of the second dorsal (base of first dorsal 55-58). From a rounded anterior lobe (anterior height 14-23), somewhat higher than the depth of the body below it, the height of this fin diminishes, first rapidly and then more gradually (25th spine 4.0-6.2), until the last few rays are hidden in the dorsal groove. The first anal fin is somewhat similar, but lower and much shorter (anterior height 12-16, base 14-17). It originates 57-62 (first preanal length) behind the tip of the lower jaw, and 3.7-5.2 behind the vent. After the anterior lobe, which is usually broadly rounded, the height decreases rapidly, then more gradually, until the last rays are hidden in the ventral groove. The second dorsal and second anal fins are very similar, and are located nearly opposite each other (second predorsal length 77-83; second preanal length 75-81) close behind the first dorsal and first anal fins. They are small (anterior height D_2 - 3.1-4.7; A_2 - 2.8-3.9; length of base: D_2 - 3.6-4.6; A_2 - 4.0-4.9; length of last ray: D_2 - 4.2-7.2; A_2 - 4.8-6.7), with short slightly convex leading edges and long deeply concave trailing edges, terminating in long points. The ultimate rays are broad and flattened proximally and adhere closely to the body of the fish.

The pectoral fin is moderately long (19-27), and its tip is usually rounded. Its origin is just behind the opercle (prepectoral length 25-27) and low on the side of the fish (distance from orig. D_1 to orig. P_1 12-14). About eight of the lowest rays are very short, forming a subbrachial dilatation.

The pelvic fins, which also originate near the opercle (prepelvic length 26-29), are somewhat shorter (length 14-22) and are so narrow that their area is negligible, and they appear to consist of a single spine. Poey (1861, pl. 16, figs. 4 and 5; the explanation of the plate [p. 241] mistakenly refers to these figures as of the pectoral) describes the structure of this fin in detail.

The color of the white marlin has been described by Poey (1861), Goode (1881), LaMonte (1955), and Nakamura et al. (1968). The back and the top of the head are described as dusky blue, and the back also as brilliant greenish blue, darker above (LaMonte, 1955). The underparts are silvery white. The transition from the dark dorsal to the white ventral areas is gradual, passing through lines of rich purplish brown and smokey grey (Goode, 1881). The sides are crossed by vertical light blue or light lavender bars (LaMonte, 1955). The cheeks and opercles are dusky, with a pearly sheen. The membrane of the first dorsal fin is deep blue or bluish purple, with circular dusky spots near its base. The rays are dark blue or black. The first anal fin is dark blue or bluish purple distally, and pearly white proximally. The second dorsal is a deep blue, and the second anal is somewhat lighter b Outer surfaces of the pectorals are dusky wit pearly sheen, and the inner surfaces bluish pur The pelvics are blue black. The caudal is dusky b ish or brownish, sometimes with pearly reflecti in the lower lobe, and the caudal keels are blu black.

The colors change in response to the state of exc ment of the fish. Jaen (1964) states that whe billfish following a bait is pale green, it probably not strike, but if it is dark-colored and its fins shiny bright blue, it will strike immediately. The p toral fins of the white marlin "light up" especia brilliantly in these circumstances.

The lateral line of the white marlin is simple a more conspicuous than those of other Atlan istiophorids. The lateral line originates at the top the opercular aperture, whence it runs longitudina for a short distance, then curves away from the ba and reaches the median line of the fish near the tip the pectoral. The lateral line was described in so detail by Poey (1861) and Talbot and Penrith (196 but LaMonte (1958a) has made the most detail study of the lateral line, scales, and skin of the wh marlin now available. The lateral line is 4 mm wi (6-8 mm, according to Talbot and Penrith, 196 Specialized subcutaneous overlapping lateral li scales have a complicated system of holes and du leading through holes in the surface of the skin to t exterior. These scales roof a relatively wide co tinuous tube (Talbot and Penrith, 1963). Nonspeci ized subcutaneous scales are thorn shaped, regul and only occasionally and slightly overlapping. Th show very clearly externally, but are completely co tained in fibrous integument, and also complete covered by the outer skin of the fish. The surfa scales are very small and delicate, with round, radi ly marked bases and glassy perpendicular spin There are small round openings through the skin u der many of these scales. A histological description marlin skin by Rasquin is included in LaMor (1958a).

The gills include four double arches and an acc sory, reticulated as in swordfish (Poey, 1861). LaMo (1958b) states, for billfish in general "The wi laminated gills are paired; each pair is connected fr the base to within a short distance of the free margi as if woven together. Gill-rakers are absent vestigial. This gill structure is typical of the grou We have not found gill rakers on any of the wh marlin which we have examined.

Poey (1861), LaMonte (1958b), and Nakamura al. (1968) discuss the dentition of white marlin brief The dentition is weak, consisting of small patches denticles or villiform teeth on the jaws, palatin palatine membrane, and pharyngeals. Poey also me tions anteriorly directed denticles on the sides of t bill.

1.31a Internal morphology[3]

ie nostrils are separated by a cup-shaped space are enclosed in a common cavity (Poey, 1861). olfactory rosette is radially shaped and has 44-50 l lamina, the distribution of blood vessels on h cannot be seen with the naked eye (Nakamura ., 1968). The structure of the skull and bill are described by Poey (1861) and Nakamura et al. 3). The following description of the skull and of ebral characteristics are from the latter authors.

he cranium is hard, elongated, and on the whole, long and i. The post-occular portion of the head is short. The anterior, tral side of the vomer and the parasphenoid is rather thin. temporal crest and the pterotic crest run almost parallel to h other. There is a small and fairly well developed projection the upper back side of the frontal bone.

he haemal spines and the neural spines of the vertebrae form elongated parallelogram. The lateral apophysis is poorly eloped. There are 24 vertebrae (12 + 12 = 24).

vertebral characteristics of white marlin and r istiophorids were discussed and compared in il by Ueyanagi and Watanabe (1965), and Monod 8, Fig. 784) described the osteological structure of caudal region. Morrow (1957) studied and com- d the morphology of the pectoral girdle in the is *Makaira*, including *T. albidus*.

he internal organs of white marlin were described fly by Poey (1860) and Nakamura et al. (1968), Krumholz (1958) provided information on tive weights of some of the viscera. The following cription is abstracted from the more comprehen- work of LaMonte (1958b), from which we show ventro-lateral aspect of the viscera (Fig. 3). The neys lie at the top of the body cavity against the ebrae. The peritoneum separates them from the , physoclystic air bladder, which consists of a dou- row of bubblelike chambers closed in a common outer membrane. The stomach is at the anterior end of the body cavity, and below it, anteriorly, are the caecal mass, capped by the liver, and the heart. The spleen is below the stomach on the left side, behind the caecal mass. Adjacent to it is the intestine, which makes one coil upon itself and runs along the body wall to the vent which lies in a ventral groove with the urinogenital opening. The gonads lie on either side of the stomach. Krumholz (1958) presented the relative weights of different organs of 42 white marlin caught off the Bahamas in percentage of the weight of the fish, and discussed their variations with the size and sex of the fish.

No analyses of geographic variations in morphological characters were found in the literature, and consequently no subpopulations have been defined on this basis.

Morphological changes with growth have not been well defined, since nearly all of the available morphological data is for adults, and no thorough compilation and analysis of this have been published. Robins and de Sylva (1960), pointed out that the development of the bill in *Tetrapturus albidus* and other "Tetrapturinae" was negatively allometric. The bill length (30) of the 125 mm (body length) postlarva described by de Sylva (1963), however, falls in the range for adult material (25-35). The first dorsal fin of this specimen is high medially (25th ray 30) and saillike. The pectoral fin is much shorter than in adults (10.7 vs. 19-27) and the pelvic much longer (29.6 vs. 14-22). The body is more slender and compressed (greatest depth 11 vs. 12-18, greatest width 5.8 vs. 5.2-9.4, usually 7.5-8.5), with the greatest width near the orgin of the pectorals rather than near the origin of the first anal fin as in adults.

Data (unpublished) for two specimens of intermediate size (904- and 964-mm body length) examined by C. R. Robins and Mather are available. In general, these individuals had attained adult proportions except that the central part of the first dorsal fin remained higher (14.7 vs. 3.1-4.7), the pectorals

[3]his section heading has been introduced by the authors since utline provides no place for this material.



Figure 3.—Diagram of *Makaira albida*, showing ventral-lateral aspect, with caecal mass reflected to left. Approximately natural size. Dissection by Francesca R. LaMonte. Drawing by Janet Roemhild. (From LaMonte, 1958b, Fig. 13 and caption.)

shorter (17.5-19.2 vs. 19-27), and the pelvics longer (26.0-26.7 vs. 14-22). The larger of these was described by de Sylva (1963) as having the adult characteristics nearly fully developed.

1.32 Cytomorphology

The genetic make up of the white marlin has not been dealt with in any detail.

1.33 Protein specificity

Edmunds (1972) analyzed samples from over 100 white marlin captured in the middle Atlantic Bight, off the Bahamas, in the northern Gulf of Mexico, and off Venezuela electrophoretically for the forms of the proteins transferrin, esterase, 6-phosphogluconate dehydrogenase (PGD), malate dehydrogenase (MDH), hemoglobin, lactate dehydrogenase (LDH), and glucose 6-phosphate dehydrogenase (G6PD) present. He concluded that these analyses provided no basis for separating the white marlin from these areas into different subpopulations.

2 DISTRIBUTION

2.1 Total Area

The white marlin is distributed over nearly all of the Atlantic Ocean from lat. 35°S to 45°N, including the Gulf of Mexico and the Caribbean Sea (Figs. 4, 5). Most of this range is documented by catches of the Japanese longline fishery (Ueyanagi et al., 1970; Mather, Jones, and Beardsley, 1972). Data from other fisheries (Furnestin et al., 1958; Postel, 1959; Rodríguez-Roda and Howard, 1962; Robins, 1974), however, establish the presence of the species in the eastern North Atlantic near the Strait of Gibraltar. The white marlin has occasionally been recorded outside this area, perhaps on the basis of strays. Legendre (1928) and Desbrosses (1938) reported individual captures west of Brittany, France, Tortonese (1940, 1961, 1962, 1970), Lozano Cabo (1958), and Rodríguez-Roda and Howard (1962) recorded a few catches in the western Mediterranean. Also, Japanese longliners have taken a few white marlin near lat. 45°S and long. 50°W, and near lat. 40°S and long. 15°E (Ueyanagi et al., 1970). The distribution of the Japanese longline effort does not provide complete coverage of the Atlantic Ocean in all seasons. Therefore the seasonal distribution of the species and the limits of its normal range, particularly in the South Atlantic, may not be completely defined.

Pollutants such as chlorinated hydrocarbons (Harvey et al., 1972), petroleum (Horn, Teal, and Backus, 1970; Morris, 1971), and plastics (Carpenter et al., 1972) have found in the surface waters of areas inhabited by white marlin, but what effect, if any, these may have on this species is unknown.

The known distribution encompasses most of a (Rosa, 1965) ASW (Atlantic, SW) and ASE (Atlan SE), as well as the part of ANW (Atlantic, NW) sc ot lat. 45°N, and the Atlantic portions of P (Southern Ocean, W) near South America and Afr Natural regions (Rose, 1965) included are the N foundland waters (5.2.1); the Florida, Gulf Stre and Atlantic Drift Current waters (5.3.1-.2 [rarely]); the Sargasso Sea and Azores waters (5. .2); the western part of the Mediterranean Sea (5 [rarely]); the Gulf of Mexico, Carribean S Bahama, Atlantic North Equatorial Current, Cape Verde waters (5.6.1,-.2,-.3,-.4,-.5); the Gui region (5.7); the Atlantic South Equatorial Curr region, E. Brazilian, S.E. Brazilian, Bengu Current, and S.W. African waters (5.8.1-.2,-.3,-.4,- the Atlantic southern gyrals (5.9.1); and the Agul waters (1.5.4 [rarely]).

This area is so large that it is difficult to class The distribution varies seasonally, however, reach the higher latitudes in the respective hemisphe during the local warm seasons only. For the most pa white marlin are found in deep (over 100-m) b water with surface temperatures over 22°C a salinities of 35-37‰. Some seasonal feeding conc trations and migrations, however, occur in wat with characteristics differing from one or more these. Average air temperatures of regions wh white marlin occur are usually moderate to war 15°-28°C. Currents of from 0.5 to 2 knots occur o much of their habitat. The productivity of most of t distributional area is considered to be low.

2.2 Differential Distribution

2.21 Spawn, larvae and juveniles

The early stages of the life history of white mar are little known. As a result of collections of lar 3.0-3.5 mm long made from RV *John Elliot Pillsbu* 24 July-13 August 1964, de Sylva postulated th spawning grounds for white marlin in the west North Atlantic (Stephens, 1965). These were nor east of the Little Bahama Bank, off Abaco Islan northwest of Grand Bahama Island (both in reg 5.6.2), and southwest of Bermuda (region 5.4 Ueyanagi et al. (1970) state that white marlin migr to subtropical waters to spawn, with peak spawnin early summer. They record collections of larvae only a few of a very large number of collecting s tions (Fig. 6) in November-April. Four collecti were made in the central South Atlantic (near 20°N, long. 20°W). Two were in the western So Atlantic; one near lat. 22°S and long. 32°W, and near lat. 8°S and long. 35°W, just off the easte most part of Brazil. Two were in the central No Atlantic, near lat. 20°N and long. 35°W. Another in the western North Atlantic, near lat. 8°N, lo 50°W.



Figure 4.—Distribution of white marlin by season: a) May-October; b) November-April. (From Ueyanagi et al., 1970, Fig. 18 and caption.)

Ueyanagi (1959) suggested that some of the larvae ntified by Gehringer (1957, Figs. 6, 10, 14) as fish might have been white marlin. These cimens were collected in the western North Atlan- from Cuba to the Carolinas, in May-September. Hayasi et al. (1970)[4] state that the high hook rate off southern Brazil in November-April is associated with the appearance of large-sized fish, which, with the appearance of postlarvae and juveniles, assures that this is an immigration of spawning adults. They note that "White marlin seem to spawn rarely in the equatorial waters", but that "matured adults appear in the northern Caribbean Sea and off Florida during

[4] The material in Hayasi et al. (1970) is a summary of English of material which was presented by Ueyanagi et al. (1970) in anese, except for a synopsis, table and figure captions, names of fishes, and English language references, which were presented in English (S. Hayasi, pers. commun.).

Figure 5.—Distribution and apparent relative abundance of white marlin in the Atlantic Ocean. Data are from records of the Japanese longline fishery, 1956-67. The catch per unit of effort (CUE) for each month in rectangle is the arithmetic mean of the CUE of each month that the area in the rectangle was fished in the 12-yr period. CUE is the number of fish caught per 100 hooks. (From Mather et al., 1972, Fig. 6 and caption.)



Figure 6.—Distribution of larvae of white marlin, November-April. Large dots - larvae present; small dots - larvae absent. (From Ueyanagi, 1970, Fig. 21 and caption.)

il through June. Collection of juveniles in January he adjacent waters, Lat. 10°N and 20°N, Long. W and 60°W, suggests long lasting spawning on of the species."

2.22 Adults

dult white marlin are very widely distributed over warm parts of the Atlantic Ocean and the adja- seas, but the degree of concentration, or apparent ndance, varies greatly, both geographically and onally.

he catch and effort statistics of the Japanese line fishery comprise the most comprehensive s of data on the distribution and apparent relative ndance of adult white marlin available. The effort the catches of each species tabulated by months 5° (latitude and longitude) quadrangles, and dis- utional charts of catch and effort, for the first s of the fishery, 1956-62, were published by hama, Myojin, and Sakamoto (1965). Similar and charts for each of the years 1962-70 were lished by the Research Division of the Fisheries ncy of Japan (1965, 1966, 1967a, 1967b, 1968, 1969, , 1971, 1972) in its annual reports of effort and h statistics of the Japanese tuna longline fishery. eral authors have organized this enormous mass of , including that for white marlin, into more com- form. Wise and Le Guen (1969) presented catch s by years, months, and areas for 1956-63. Wise 8) and Wise and Fox (1969) made similar yses of the data for 1964 and 1965. Wise and Davis 3) show the average catch rates for the years 1956- r each quarter of the year by contours (Fig. 7). We e added indications of seasonal coastal occurrences to these charts. Ueyanagi et al. (1970) have also presented a thorough analysis, showing catch rates by half-year periods (Fig. 4) and by months. Mather et al. (1972) presented the average catches for white marlin for the years 1956-67 by months (Fig. 5). Our discussion of the oceanic distribution of adult white marlin is based mainly on these sources.

In the South Atlantic, white marlin concentrate in the eastern side off Angola, in June-July, and on the western side off Brazil and sometimes in the center, for most of the rest of the year (Figs. 5, 7). The concentration off Brazil centers off Recife in September and October, but extends southward to the vicinity of Rio de Janeiro. In November the concentration moves to the south and also well to the east. In December-February the catch rates are highest off Rio de Janeiro, but isolated areas of concentration occur to the eastward, in the central and eastern parts of the ocean. March-May is a period of generally low catch rates, except for isolated areas off easternmost and southernmost Brazil. Concurrently with the June-July concentration of white marlin off Angola there are small areas of high catch rate off easternmost Brazil, and in July in mid-ocean between these two areas. Perhaps because of the limited development of sport fisheries in the South Atlantic, no important coastal concentrations of white marlin have been found in that ocean.

A few isolated records, however, supplement the data provided by the Japanese longline fishery on the distribution of white marlin in the South Atlantic. LaMonte (1955) reported that the species occurs off Cabo Frio, Brazil, December-February. During a visit to the Yacht Club at Rio de Janeiro in January 1969, the senior author was informed that sport fishermen



Figure 7.—Distribution of catches of white marlin (per 1,000 hooks) in the four quarters of the year, 1956–68. (From Wise and Davis

casionally caught white marlin off that city. Talbot nd Penrith (1962, 1963) recorded single captures of hite marlin off Cape Town, South Africa, in March 961 and February 1962; 110 miles north-northwest of Walvis Bay, South West Africa in April 1961; and off ne island of St. Helena (no date reported). These uthors also noted that in the summer of 1961, three riped marlin, *T. audax*, were caught in the same ea as the white marlin. Records of the Game Fish nion of Africa (Anon., 1972b) show two additional hite marlin, 63 pounds (28.6 kg) and 111 pounds 50.5 kg), caught off St. Helena on 17 February 1958 nd 16 February 1961, respectively.

The information on the distribution of white marlin n the North Atlantic obtained from the oceanic isheries is supplemented by studies of important easonal coastal concentrations and records of oc-asional occurrences in various localities. Peak in-hore sport fishing and offshore longline fishing do not ecessarily coincide seasonally in each area, probably ecause of differences in distribution relative to depth f water, and changes in swimming depth and feeding abits.

In summer, the peak longline catches are in the Gulf of Mexico and the northwestern Caribbean. Low atch rates (Figs. 5, 7) over extensive areas indicate hat white marlin are widely but thinly distributed ver much of the deep waters of the open North Atlan-ic during this season.

The most important inshore concentrations in ummer and early fall are on and near the edges of the ontinental shelf in the northern Gulf of Mexico, from Cape Hatteras, N.C., to Cape Cod, Mass., and off La Guaira, Venezuela. A less known concentration in late ummer and early fall occurs outside the Strait of Gibraltar.

Gibbs (1957) studied seasonal variations of longline atches in relation to water temperature in the Gulf of Mexico. He noted a summer concentration off the Mississippi Delta, which dispersed as the water cooled n fall. This concentration and another at the De Soto Canyon in the northeastern Gulf contribute impor-antly to sport fisheries based at New Orleans, La.; Mobile, Ala.; and Pensacola, Panama City, and Destin, Fla. (Mather et al., 1972; Nakamura and Rivas, 1972). Some successful sport fishing for white narlin has also occurred at the edge of the continental helf off Port Aransas, Tex. (M. H. Weil, pers. com-mun.). White marlin are available to the sport isheries in the Gulf from early June into October, vith peak abundance off the delta in July and at the De Soto Canyon in August.

The concentration of white marlin which supports he largest sport fishery for this species in the world ccurs in summer between Cape Hatteras and Cape Cod (Farrington, 1937, 1949a, 1949b; Earle, 1940; June nd Reintjes, 1957; Migdalski, 1958; de Sylva and Davis, 1963; Brooks, 1968). The greatest concentra-ion of fishing is between Oregon Inlet, N.C., and Atlantic City, N.J. In the earlier years, most of this fishing was within 30 or 40 miles of shore, on shoals such as the famous "Jack Spot" off Ocean City, Md., and other productive areas. In recent years, however, most of the successful fishing has been nearly 80 miles offshore, concentrating around the canyons at the edge of the continental shelf, from Norfolk Canyon off the Chesapeake Capes to Block Canyon off eastern Long Island.

The northeastern limit of the usual summer coastal occurrence of white marlin is off Nantucket Island, south of eastern Cape Cod, Mass. The species occurs further east and north on Georges Bank and the Nova Scotia Banks, especially along their oceanic edges (Farrington, 1949b; J. S. Beckett, pers. commun.) but rarely enters the Gulf of Maine. We know of only two records from coastal waters in the Gulf. Both were from traps at Provincetown, Mass. (John A. Worthington, pers. commun.). The first of these specimens, taken in August 1960, was deposited in the Museum of Comparative Zoology, Harvard University, Cambridge, Mass., as the first positive record of this species from the Gulf of Maine. Farrington (1949b), however, reports occurrences off the coast of Nova Scotia.

A concentration of white marlin in late August, September, and October off La Guaira, Venezuela, furnishes some of the world's best sport fishing for this species (Migdalski, 1958; Jaen, 1964; Brooks, 1968).

In late September and October after the conclusion of the sport fishing season between Cape Hatteras and Cape Cod, longline catch rates are high from the edge of the adjacent continental shelf eastward to long. 55°W. They are also good in the northern Gulf of Mexico in this period. Longline catch rates are low everywhere in the North Atlantic in November, but improve in the southeastern Caribbean in December. January has also been a poor month throughout the North Atlantic, but in February and March there have been good catches off the coast of South America from Venezuela to the equator. In March, this area of good fishing extends northward around the Lesser Antilles, the Virgin Islands, and Puerto Rico up to lat. 25°N. In April, the concentration of catches has been between lat. 15° and 25°N and long. 55° and 75°W, but there was little fishing effort further west in these latitudes. In May, the area of successful fishing extends further to the south (off western Venezuela and Colombia and northeast of Cuba and the Bahamas). June finds good longline fishing in the western Caribbean, the northern Gulf of Mexico, and east of the Bahamas and northern Florida.

Spring is the peak season for inshore sport fishing for white marlin in the Straits of Florida and among the Bahamas. Areas which support important sport fisheries for this species are Havana; Bimini and Cat Cay on the northwestern edge of the Great Bahama Bank; the Berry Islands at the northern end of the Tongue of the Ocean; and Walkers Cay at the

northern tip of the Bahamas (Farrington, 1937, 1949a, 1949b; Hemingway, 1949; Migdalski, 1958; Brooks, 1968). This period is also the most productive for sport fishing for white marlin off southeastern Florida, Puerto Rico (Erdman, 1957), and the Virgin Islands.

White marlin apparently are less abundant in the eastern North Atlantic and the Mediterranean, but have been recorded occasionally in those waters. LaMonte (1955) reports catches off "Portugal, Azores Islands, Madeira, and the Italian Riviera." Legendre (1928) and Desbrosses (1938) reported single captures of *T. belone* and *T. Lessonae*, respectively, both taken in September from tuna fishing boats well off Brittany (west of the entrance to the English Channel and "200 miles from Groix," respectively). Both records have been placed in the synonymy of *T. albidus* (LaMonte, 1955). Rodríguez-Roda and Howard (1962) reported on 19 specimens of white marlin captured off the southern coasts of Portugal and Spain in 1961—16 in August, 1 in September, and 2 in October. Thirteen of these captures were in the Atlantic, 2 in the Strait of Gibraltar, and 4 at La Linea, Spain, just inside the Mediterranean Sea. Robins (1974) lists 37 specimens caught off southern Portugal in late July and August 1961, and 2, observed by Mather, which were caught off Cadiz, Spain, on 6 October 1969. Twenty-three of these fish were caught by hook and line (most, if not all, by longline), and 35 by tuna traps (almadrabas). Furnestin et al. (1958) describe two specimens taken off Morocco, one by longline off Mazagan, August 1950, and one by trawl net off Casablanca, July 1953, and state that the species is taken very frequently by longline off Tangier and in the vicinity of Casablanca. A photograph (Lozano Cabo, 1958, Fig. 93) of a fish caught in a tuna trap at Ceuta, Spain (African coast, just inside the Mediterranean) is captioned and referred to as *T. belone*, but we concur with Robins and de Sylva, 1960, in re-identifying this fish as *T. albidus*. This information indicates that white marlin are fairly abundant in the Atlantic approaches to the Mediterranean in August and September, and occasionally enter the western part of the Alboran Sea.

Records of *T. albidus* in the remainder of the Mediterranean, however, are very scarce. Tortonese (1940, 1961, 1962, 1970) re-identified two mounted specimens, which had been caught near Genoa, Italy, in the nineteenth century, as *T. albidus* and reported the capture of two others in a small tuna trap 23 km east of that city in September 1970. In his opinion, the species occurs only in the western part of the Mediterranean, and is very rare there. Cesareo (1967) recorded the harpooning of a white marlin in September 1967, near the Island of Gallinara in the Ligurian Sea. Sarà (1968) noted that *T. albidus* contributed to the catches of some Sicilian tuna traps, and Bini (1968) (not seen by us, but quoted by Tortonese, 1970), also stated that this species occurs in Sicilian waters.

Annual variations in distribution are most notable in areas where the distribution is seasonal, especia near the limits of the range of the species. Th variations are usually attributable to one or more the determinants of distribution changes (see 2.3

2.3 Determinants of Distributional Changes

The distribution of white marlin is control primarily by the necessities of feeding and spawni and secondarily by conditions of the environmen

The spawning areas of white marlin have not b completely defined, but it appears that some coas concentrations (Bahamas [de Sylva and Davis, 19 Stephens, 1965], Cuba, and the Greater Anti [Erdman, 1956]) are related to spawning, while oth (northern Gulf of Mexico [Gibbs, 1957], Ca Hatteras-Cape Cod [de Sylva and Davis, 196 Venezuela [Jaen, 1964], and outside the Strait Gibraltar [Robins, 1974]) are related to feeding. T latter are all in areas of relatively high product ity.

It also appears that the oceanic concentration southern Brazil is a spawning assembly (Hayasi et a 1970).

Water temperature appears to exert an import influence on the distribution of white marlin. Gi (1957) related the monthly changes in the distribut of white marlin catches in the Gulf of Mexico to position of the 75°F (23.9°C) isotherm. Ovchinnik (1970) stated that a temperature of 24°C is optir for white marlin. Squire (1962) showed that at ploratory longline stations where white marlin w caught by MV *Delaware* and RV *Crawford* in western North Atlantic, surface temperatures rang from 70.0° to 83.0°F (21.1° to 28.3°C), with weighted average of 76.6°F (24.8°C), a temperatures at the estimated fishing dep (estimated as 173 feet or 52.7 m) ranged from 50.0° 80.5°F (10° to 27°C), with a weighted average 68.8°F (20.5°C). De Sylva and Davis (1963) showe from extensive observations taken of southern N Jersey and Maryland during the summer of 1959, th the white marlin grounds were generally characteriz by surface water temperatures between 78° and 80 (25.6° and 26.7°C). Most of the longline catches white marlin have been on the warm side of the 20 surface temperature isotherm (Fig. 4.)

Ovchinnikov (1970) reported that commercial co centrations of white marlin occur near shores in are where less saline coastal waters mix with more sali oceanic waters. De Sylva and Davis (1963) noted th in the summer of 1959 the white marlin grounds southern New Jersey and Maryland were of high salinity than the surrounding inshore waters, a were identifiable by an abrupt increase from lower higher salinity values. Those off Ocean City, M were further characterized by low oxygen values s rounded by high values. Important white marlin are showed distinctly high plankton volumes.

ıe color of water appears to be an indicator of able abundance of white marlin. Nakamura l) and Nakamura and Rivas (1972) found that ng success for white marlin in the northern Gulf exico was greater in proportion to the blueness of vater, and poorer in proportion to its greenness. ng the 1972 season, there was an unusual oc- ɔnce of green water at the normally productive e marlin fishing areas at the edge of the continen- helf off Maryland and southern New Jersey, and ıes in this water were very poor (M. Maiorana, commun.). In similar areas off Virginia and h Carolina, the water was the normal blue color, the fishing for white marlin was excellent, in- ting that the poor fishing further north was not to a scarcity of fish.

some areas, white marlin are concentrated near (usually occurring at interfaces between different ses of water), or weed lines.

fferential distribution of white marlin is also in- ıced by bottom topography. Steep drop-offs, sub- ne canyons, and shoals, when located in areas suitable water conditions, are often the scene of ɔrtant feeding concentrations of white marlin and ptionally productive fishing. Shoals of this nature ıde the Placer de la Guaira, off the Venezuelan of that name (Jaen, 1964), the "Cigar" off the inia Capes, the "Jack Spot" off Maryland rington, 1937, 1949a, 1949b; de Sylva and Davis,), and the Five Fathom Shoal of southern New ey (de Sylva and Davis, 1963). Drop-offs produc- ood white marlin fishing are found in many areas, ıding the Bahamas, Cuba, Puerto Rico, and the in Islands. In recent years, excellent fishing for e marlin and other oceanic game fishes has loped at many of the canyons along the edges of continental shelf. Among the more important are De Soto Canyon in the northeastern Gulf of Mex- Norfolk Canyon off the Virginia Capes; hington, Baltimore, and Wilmington canyons off Delmarva Peninsula; and Hudson Canyon off York City.

.4 Hybridization

o information was found in the literature.

BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

ıe white marlin is heterosexual. No apparent ex- al sexual diamorphism exists, except that the ıles attain larger sizes than the males (de Sylva Davis, 1963; Ueyanagi et al., 1970; Nakamura Rivas, 1972), and may be somewhat heavier than ɛs of the same length (de Sylva and Davis, 1963).

3.12 Maturity

Ueyanagi et al. (1970) state that the species attains sexual maturity at a length from orbit to fork of tail of about 130 cm. The relationship of age to size of white marlin is not known.

3.13 Mating

The only reference we have found to mating of white marlin is by Hemingway (1935, not seen by us, but quoted by LaMonte and Marcy, 1941). He reports paired breeding in the current off Cuba in May.

3.14 Fertilization

Fertilization is external. Eggs and sperm are probably discharged separately by adjacent fish and come together in the ambient water.

3.15 Gonads

White marlin gonads consist of two sausage-shaped organs tapering at both ends which lie ventral to each side of the stomach. The ovary is yellowish to orange red, circular in cross section and often covered with thick layers of connective tissue whereas the testis is white to pinkish, triangular in cross section, and is lobular in appearance, without connective tissue covering (LaMonte, 1958b).

LaMonte (1958b) found that in one male the gonads were about 280 mm in length and 80 mm in circumference and very flabby; but in two others they were the same length but about 170 mm in circumference and very firm. In a male specimen weighing 17.7 kg the gonads measured 178 mm in length and about 38 mm in circumference.

Krumholz (1958) recorded the ratio of gonad weight to body weight for 20 male and 22 female white marlin taken in the Florida Current in late April 1956. The gonads contributed from 0.097 to 1.266% of male body weight with an average of 0.422 and from 0.882 to 9.762% of female body weight with an average of 4.556. The male for which the testes was 1.266% of the body weight was a 36.4-kg fish, the largest examined. The relative weight of gonads was also much greater in large females than in small ones. Since these gonads were not yet ripe, these figures do not indicate the maximum relative weight of the gonads. The average gonad weights, in percent of weight of fish by 10-pound (4.5-kg) groups (Krumholz, 1958), were as follows:

	Weight	*Number of fish*	*Gonads*
Male:	20-29	2	0.216
	30-39	5	0.224
	40-49	3	0.360
	50-59	6	0.451
	60-69	2	0.750
	70-79	1	0.333
	80-89	1	1.266

	Weight	*Number of fish*	*Gonads*
Female:	50-59	3	1.521
	60-69	7	2.641
	70-79	5	5.007
	80-89	6	5.467
	90-99	1	8.338

Ueyanagi et al. (1970, Fig. 27) showed that the percentage group maturity of female white marlin caught in the area east of Brazil in November-March increased with size of fish.

No information on the fecundity or number of eggs produced in a year was found.

There is no direct information on the number of broods produced during a lifetime, but two white marlin tagged as adults have been recaptured about 6 yr after their release [unpublished WHOI (Woods Hole Oceanographic Institution) data]. This indicates that a white marlin might spawn at least six times in its lifetime.

No information was found on variation in fecundity with health or condition, or on correlation between the number of spawn or young produced and the nature of the environment.

3.16 Spawning

The available information indicates that white marlin spawn once a year (de Sylva and Davis, 1963; Hayasi et al., 1970; Ueyanagi et al., 1970). Knowledge of spawning seasons, areas, and behavior is incomplete, because of difficulties in identifying larvae, and the lack of continuous and comprehensive gonad studies. The known distribution of larvae and juveniles is described in 2.21. Ueyanagi et al. state that spawning occurs in subtropical waters and is at its peak in early summer. In their plot of percentage group maturity of female white marlin by areas and quarters of the year, Ueyanagi et al. (1970, Fig. 20) show that the only area in the South Atlantic with a concentration of maturing fish is the one between lat. 20°S and 30°S and long. 20°W and the South American coast. In the fourth quarter of the year, 77% of 51 fish examined in the part of this area west of long. 40°W and 95% of 736 examined in the part east of that meridian were maturing. In the first quarter, 89% of those examined in both parts of the area (94 in the west and 77 in the east) were maturing. Their studies of maturity of females in the area east of Brazil showed a decrease in the number of maturing fish and an increase in the number of spawned-out fish from November through January (Ueyanagi et al., 1970, Fig. 23). Hayasi et al. (1970) note that the rise in hooking rate off southern Brazil in November-April is associated with the appearance of large fish, presumably spawners, in the area, and that the accuracy of this assumption was demonstrated by the appearance of postlarvae and juveniles in the area. They also state that the white marlin taken wes long. 25°W were spawners, whereas those taken of there were feeding fish.

The same authors mention that white ma caught in the Caribbean in winter had derdeveloped gonads, but that adults with matu gonads were caught in the northern Caribbean April-June. These findings are in accord with E man's (1956) statements that the ovaries of w marlin taken off Puerto Rico were enlarged in A and that the best formed eggs he had seen were fish taken there on 9 June. They also fit with de Sy and Davis' (1963) report of nearly ripe females in eastern Straits of Florida in March and May Krumholz's (1958) finding of nearly ripe males in same area in late April. De Sylva and Davis fo that the ovaries of females taken off Cape Hatteras June and off Maryland and southern New Jerse summer were in postspawning or resting condit although many of the males were still ripe. I generally believed that summer concentrations white marlin in the Gulf of Mexico and between C Hatteras and Cape Cod are of spent fish which are tent on feeding. These findings suggest that spawn in the western North Atlantic occurs almost entir in spring, in the areas noted here and under 2.21

The spawning period and areas coincide at leas part, with those of the western Atlantic bluefin tu *Thunnus thynnus thynnus* (Rivas, 1955; Potthoff Richards, 1970). The young of this species share t summering area between Cape Hatteras and C Cod with the white marlin.

The spawning areas of white marlin are in deep blue oceanic waters with generally high surf temperatures (20°-29°C, except in the Atla southern gyrals, reg. 5.9.1), and high surface salini (>35‰). Except off Cabo Frio, Brazil, the prod tivity of these waters is considered to be low.

Nothing was found in the literature about variations of spawning grounds.

Little is known of the ratios and distribution of s es on the spawning grounds. Ueyanagi et al. (19 found 2,037 males and 1,051 females in the a between lat. 5°S and 30°S, and long. 15°W and coast of South America, during November-April. T area includes the South Atlantic spawning grou described above (lat. 20°S-30°S, long. 20°W to coast), and the period coincides with the spawn season as indicated by Hayasi et al. (1970). In April, Krumholz (1958) found 20 males and females in the eastern Straits of Florida.

Hemingway (1935) described the breeding of wh marlin as follows: "White marlins breed off Cuba May. They breed in the same way that the grou does, except that as they are a fish of the current, t breed in the current instead of on the reef. The fem marlin heads into the current while the male head the opposite direction, and while they are side by s the female expels the eggs and the male the milt;

ıle then catches the eggs in the basket-like opening his gill covers and lets them pass out through his ıuth." LaMonte and Marcy (1941) question the ısibility and utility of the latter operation, but the servation of paired spawning may nonetheless be sically accurate.

3.17 Spawn

White marlin eggs have yet to be accurately iden-ied according to Morrow (1965), but are free-ating in the water column. De Sylva and Davis 963) assume that ripe eggs would appear translu-nt.

3.2 Preadult Phase

3.21 Embryonic phase

Embryonic development of the white marlin has t been observed and information concerning this ıase of life is not available.

3.22 Larvae phase

Other than observations made on planktonic mples of larval white marlin, little is known about eir development and activities during this stage of e. The larval white marlin has heavy, pointed oper-lar spines and lacks the characteristic long bill of e adult (Scotton and de Sylva, 1972) (Fig. 8).

The postlarvae of white marlin are also rare. De lva (1963) gives a detailed description of a 125-mm ecimen collected off Wilmington, N.C. This fish is table in that it bears a saillike dorsal fin with four stinct ocelli near the base. A second very similar ecimen (191 mm in total length) was taken off the rthwest coast of Cuba (Anon., 1968).

Information on the behavior and continuous velopment during this period in the life cycle, wever, is not available.

3.23 Adolescent phase

No material in the range between these postlarvae ıd the 904-mm and 964-mm specimens mentioned in 1.31 are available. As noted above, the latter specimens have attained all the adult characteristics except that the middle of the first dorsal fin is somewhat higher, the pectorals very slightly shorter, and the pelvics longer than in adults. Maturity occurs at a length (from eye to fork of tail) of about 130 cm (145-cm body length) (Ueyanagi et al., 1970). This is equivalent to a weight of about 37 pounds (17 kg) for females and 31 pounds (14 kg) for males (de Sylva and Davis, 1963). These authors and Nakamura and Rivas (1972) indicate that very few smaller (immature) white marlin are taken in the U.S. sport fisheries. Ueyanagi et al. (1970), however, show that many white marlin in this immature size range are taken by the Japanese longline fishery in equatorial waters lat. 10°S to 10°N, and north of lat. 10°N in May-October. Most of the small fish taken from lat. 10°S to 30°N are males. Most of the available biometric data, however, are from mature fish.



gure 8.—Baby white marlin, about ⅛ inch (3.2 mm) long, th heavy, pointed spines and lacking the characteristic bill. rawing by Joy Godfrey Alexander, from Scotton and de lva, 1972, Figure p. 200.)

3.3 Adult Phase

3.31 Longevity

Weight frequency distributions and tag returns suggest that the white marlin may be longer lived than was once believed. Each of two weight frequency polygons for fish taken off Maryland and southern New Jersey (de Sylva and Davis, 1963, Figs. 4, 5) show several peaks. If these peaks actually represent year classes, as they seem to, this is a good indication of longevity. In addition several tagged white marlin have been recaptured after over 4 yr at large (Mather et al., 1974) including two recent recaptures of fish which had been out over 71 mo (unpublished WHOI data). Since these fish were not remarkably small when tagged, the life span must sometimes exceed 6 yr.

3.32 Hardiness

White marlin seem to be of a reasonably hardy nature. Mortality rates for fish which had been subjected to the rigors of capture on hook and line and tagging and had been at large for from less than 1 mo to more than 48 were 30% per year with 95% confidence limits of 23% and 36% and a coefficient of instantaneous total mortality of 0.36 ± 0.10 (Mather et al., 1974).

These fish do seem, however, to be somewhat temperature sensitive. Earle (1940) reported that in the Ocean City, Md. fishery, white marlin vanished after a sudden drop in water temperature of a few degrees and returned when water temperature returned to normal.

There is no information on survival of white marlin in confined environments, or on variations in hardiness with age, size, or physiological states.

Table 2.—Trophic relationships between epipelagic predatory fishes of the tropical waters.

Main group of predators	1	2	3	4	5	6	7
1. Small tuna (*K. pelamis*, young *Thunnus* sp.)	-	E, C	E,(C)	C	V, C	(V)	E
2. Large tuna (*Thunnus* sp.) and small marlins (*Tetrapturus* sp.)	V, (C)	-	E, (C)	(V)	V	V, C	E, C
3. Large marlins (*Makaira* sp.) and swordfish (*Xiphius gladius*)	V, (C)	V, (C)	-	(V)	V	V	C, E
4. *Coryphaena* sp.	C	(E)	(E)	-	C	-	(E)
5. Snake mackerel (*Gempylus serpens*)	E, C	E	E	C	-	E	(E)
6. Lancetfishes (*Alepisaurus* sp.)	(E)	E, C	E	-	V	-	E
7. Sharks	V	V, C	C, L	(V)	(V)	V	-

NOTE: Reading across the table, the relationships between each group of predators are seen (C - competitor; E - enemy; V - victim). The groups are numbered uniformly in the rows and columns.
Source: Parin, 1968, Table 13.

3.33 Competitors

White marlin compete with the tunas, large marlins, sharks, and to some degree with lancetfishes and are victimized by larger billfish and sharks (Parin, 1968) (Table 2).

Fox (1971) indicates from the Japanese longline data that white marlin distributions overlap strongly with those of yellowfin tuna, *Thunnus albacares*, and blue marlin, *Makaira nigricans*. Wise and Davis (1973) show that during certain seasons of the year centers of white marlin abundance coincide with those of blue marlin and sailfish, *Istiophorus platypterus*, abundance. During the summer, adult white marlin and small (2-50 kg) bluefin tuna, are both abundant off the U.S. coast between Cape Hatteras and Cape Cod (Farrington, 1949a). Since both of these fishes feed largely on small schooling fishes and squid, they must compete with each other considerably.

3.34 Predators

Very little is known about the predation on white marlin. It must be presumed that members of the shark families prey on this species at one time or another. When fighting white marlin on rod and reel the senior author has noted the presence of mako sharks, *Isurus oxyrinchus*. No documented evidence of actual attacks on white marlin are available however, although Poey (1861) described marlin on a fisherman's line as becoming furious at the approach of its natural enemy the shark and receiving frightful wounds from its adversary.

Maéda (1967) observed the killer whale, *Orcinus orca*, attacking marlin but added that such instances were probably rare since meetings of the two species were most likely infrequent.

3.35 Parasites, diseases, injuries, and abnormalities

Parasites and diseases: Records of parasite infesta and disease problems in white marlin are minim Although both problems are likely to occur in species, few exact classifications of them have b made.

Nikolayeva and Ezpeleta (1966) examined white marlin from the Gulf of Mexico and found fi 597 to 6,833 specimens of parasites per fish. Of the examined, all had members of the parasite gro Cestoidea, Trematoda, Nematoda, and Crustac three had members of Monogenoidea; and only had representatives of Sporozoa and Acanthocephs Silas (1967) and Silas and Ummerkutty (1967) su marized the known species of trematodes, cestoc and parasitic copepods found in or on white ma (Table 3). Jones (1971) mentioned reports of cra wounds caused by the squaloid shark, *Isist brasiliensis*, on various species of marlins.

Injuries and abnormalities: A common injury to white marlin is the loss of part or most of its bill.

Table 3.—Parasites found on white marlin.

Locality	Parasite	Location on host
Monogenetic trematodes (Silas, 1967):		
N.W. Atlantic	*Capsaloides cornuatus* (Verrill)	On body
N.W. Atlantic	*Capsala laevis* (Verrill)	
Cestodes (Silas, 1967):		
N.W. Atlantic	*Bothriocephalus manubriformis* (Linton)	Intestine
Copepods (Silas and Ummerkutty, 1967):		
W. Atlantic	*Gloiopotes ornatus* (Wilson)	On body

ıct problems this raises for the white marlin are not ɔwn but we have seen hearty, active fish which ve been caught with this condition.

3.4 Nutrition and Growth

3.41 Feeding

Most indications given by the degree of digestion of ious food items in the stomachs of white marlin ɔw them to be daytime feeders. De Sylva and Davis 63) found that nearly all round herring, *Etrumeus lina*, taken from marlin caught between Ocean y, Md., and Atlantic City, N.J., were only slightly ested and appeared quite fresh, indicating recent estion (i.e., from 0800 to 1700). At Bimini, umholz and de Sylva (1958) likewise observed that ne of the fish caught before 1000 still contained bait n that were only slightly digested, and fish caught 1130 and 1330 contained a small octopus and a fish respectively, neither of which was in any adnced stage of decomposition. The marlin may feed night as well, for squid which showed signs of hav- been in the stomachs for a considerable period of ıe were found in fish taken in the morning (de lva and Davis, 1963). This does not, however, pear to be the major feeding time.

The general areas of feeding may be of different tures from deep to shallow water and from near the ɔre to out in the open sea. Nakamura (1971) felt ıt sargassum lines are attractive as feeding areas for h in the Gulf. Off Bimini, Krumholz and de Sylva 58) inferred that the marlins are not primarily sure feeders but that they probably obtain a large porn of their food at depths as great as several hundred t.

The method used by the white marlin for capture of y is not certain. Earle (1940) reported that they kill stun their food by spearing it or hitting it with their l. This may not always be true, however, as whole cimens found in the stomach have been unscathed. such cases the marlin may have simply overtaken prey.

3.42 Food

Squid, it seems, play one of the most important rts as food of the white marlin in the different areas its abundance. Erdman (1958) examined seven fish d found squid in four marlins, snake mackerel in ɔ, doctorfish in two, filefish in one, triggerfish in e, and one each of a blue runner, young tuna, and ama in Puerto Rican samples. In the Bahamas, he ımined eight and found five with squid, four with eudoscopelus, five with snake mackerels and one h with octopods, doctorfish, bigeyes, and round ɔins (*Selar* sp. or *Decapturus* sp.). Krumholz and de lva (1958) also working in the Bahamas, found food nine of 50 stomachs examined, including three with ıid, three with octopods, two with unidentifiable fish remains, and one each with a crab, filefish, and balao.

In the Gulf of Mexico, Nakamura (1971) and Nakamura and Rivas (1972) found that the most consistently important food items of the white marlin from 1966 to 1971 were squid, dolphin, *Coryphaena hippurus*, and hardtail jacks, *Caranx crysos*. Mackerel were next in importance and flying fish and bonito also played a big part. Other items found were cutlassfish, swellfish, herring, barracuda, moonfish, triggerfish, remora, hammerhead sharks, and crabs but to a much lesser and more inconsistent degree.

Along the middle Atlantic coast the favorite food items for the white marlin appear to be the round herring and squid, *Loligo pealei* (de Sylva and Davis, 1963). Carangids were also well represented in addition to several other species as shown in Table 4 by number of times occuring.

This same area was studied by Wallace and Wallace (1942) (not seen by us; quoted by de Sylva

Table 4.—Frequency of occurrence of different organisms in stomachs of 55 white marlin, *Tetrapturus albidus*, taken by anglers between Ocean City, Md., and Atlantic City, N. J., 29 July to 12 September 1959, and in 18 white marlin taken at Atlantic City, 3 to 5 August 1960.

	Number of times occurring	
	1959	1960
VERTEBRATES		
Fishes		
Clupeiformes (unidentified)	1	
Clupeidae (Dussumieriinae)		
Atlantic round herring, *Etrumeus sadina*	22	1
Hemiramphidae		
Halfbeak, *Hyporpamphus unifasciatus*[1]	2	1
Carangidae		
Banded rudderfish, *Seriola zonata*	4	1
Round scad, *Decapterus punctatus*	1	1
Jack, *Caranx* sp.	1	4
Unidentified carangids	3	
Scombridae		
Mackerel, *Scomber* sp.	2	
Xiphiidae		
Swordfish, *Xiphias gladius*		1
Dactylopteridae		
Flying gurnard, *Dactylopterus volitans*	1	
Fish remains (unidentified)	9	
INVERTEBRATES		
Mollusca (Cephalopoda)		
Squid, *Loligo pealei*	13	2
OTHER		
Rockweed, *Fucus* sp.		1
EMPTY	10	4

[1]Not a bait.

Source: de Sylva and Davis, 1963, Table 5.

and Davis 1963), with similar results as far as prey species and relative abundance are concerned. They also found, however, dolphin, sand lances, *Ammodytes tobianus*, and anchovies, *Anchoa browni*.

3.43 Growth rate

Ages of white marlin have not been determined, but some estimates of growth in time intervals have been attempted. De Sylva and Davis (1963, Figs. 1-5) presented histograms of size frequencies for white marlin caught in various years and in different areas in the western North Atlantic. They believe that these fish school according to size and return to certain fishing areas in these groups. A strong year class apparently occurred from 1957 to 1959 at Ocean City, Md., peaking at 20 kg in 1957, 24.1 kg in 1958, and 28.6 kg in 1959. The difference between the modal sizes in consecutive years presumably represents the annual growth for these hypothetical year classes.

Nakamura (1971) calculated the constants a and b in the equation, $W = a L^b$ (W = weight in pounds and L = total length in inches), describing the length-weight relationships for white marlin taken in the Gulf of Mexico. He presented the length-weight-girth calculations for fish from 70 to 110 inches (177.8 to 279.4 cm) in total length which resulted from measurements taken on 162 individuals ranging from 75 to 100 inches (195.6 to 254 cm) in total length (Table 5).

3.44 Metabolism

Metabolic rates of the white marlin have not been determined.

3.5 Behavior

3.51 Migrations and local movements

Seasonal changes in distribution led to suppositions about the migrations of white marlin, but tagging has provided more positive information. In the first tagging efforts at Ocean City, Md., in the summer of 1939, Earle (1940) marked 84 white marlin, but none of these tags were returned. Tagging by the Coopers Game Fish Tagging Program of the Woods Hole Oceanographic Institution, U.S.A., (Mather, 1971; Mather et al., 1972; Mather et al., 1974) has produced considerable insights into the movements of white marlin in the western North Atlantic. A

Table 5.—Calculated weights and girths for white marlin, 70-110 inches total length, based on 162 specimens, New Orleans Big Game Fishing Club, 1966, 1968, 1970, with size range and constants shown.

Total length (inches)	Weight (pounds)	Girth (inches)
70	23	19.3
72	25½	19.8
74	28½	20.5
76	31½	21.1
78	35½	21.7
80	39	22.3
82	43	22.9
84	47	23.5
86	51½	24.1
88	56½	24.7
90	62	25.4
92	67½	26.0
94	73½	26.6
96	79½	27.2
98	86	27.9
100	93½	28.5
102	101½	29.2
104	109	29.8
106	117	30.4
108	126	31.0
110	135½	31.6

Size range			Weight constants		Girth constants	
Total length (inches)	Weight (pounds)	Girth (inches)	Log (a)	b	Log (a^1)	b^1
75-100	29-100	20-32	-5.85911	3.91484	-0.73896	1.09713

Source: Nakamura, 1971, Tables 27 and 29.

nuary 1973, some 9,000 of these fish had been mark- in that area, and 144 tags had been returned. The times of arrival in the various fishing grounds, id observations of the behavior of the fish emselves (Farrington, 1937) indicate that the white arlin which concentrate in summer in the Cape atteras-Cape Cod area first enter its southern por- ons then move northward and eastward along the ast as the season advances. Tag returns (Mather et ., 1974, Fig. 2), however, show that the fish may ove in various directions within this area during the summer. The fish leave these waters and move eastward (Fig. 9), sometimes as far as long. 40°W, in September and October. In late fall and winter, they gather off the northern coast of South America, from the Guianas to Colombia. In spring they migrate northward, some passing through the Yucatan Channel and the Straits of Florida, others through the other Passages of the Greater Antilles, and still others moving northward with the Antilles Current east of the West Indies and the Bahamas. Their migration from around the Bahamas back to their summering



Figure 9.—Tag returns from white marlin released in coastal waters between Cape Hatteras, N.C. and Cape Cod, Mass.

area apparently passes well east of the American coast. Fewer than 5% of the 124 returns from fish tagged in the Cape Hatteras-Cape Cod area show substantial deviations from this pattern. The release and recapture data for one fish released off Venezuela in the fall and three released among the Bahamas in spring (Fig. 10) also fit with the above pattern.

Results (Fig. 10) from tagging two groups of white marlin, presumably different from each other and from the one just discussed, which concentrate in the northern Gulf of Mexico in summer and off Venezuela in late summer and early fall, respectively, are clear-cut. Seasonal interchanges between the Gu Mexico in the warm season and the Straits of Flo and the adjacent Bahamas in the cold season been demonstrated, but fish marked in the latter have also been recaptured in the northern Caribb and well east of Brazil. Individuals marked of Guaira, Venezuela, in August-September have recaptured off the Guianas in November-Decem but also in the release area in November and Janu The monthly distribution of longline catches indic



Figure 10.—Tag returns from white marlin released in waters south of lat. 32°N. (Migration routes are diagrammatic.)

it a group of white marlin winters off Venezuela
l summers in the Gulf of Mexico (Mather et al.,
'2). The distribution of the group which summers
the middle Atlantic coast of the United States co-
ides in winter (off northern South America) with
it of white marlin which summer off Venezuela and
the Gulf of Mexico, and in spring (in the Straits of
orida) with other fish which summer in the Gulf of
exico. Thus interchanges between these groups,
ch as have been indicated by a few tag returns, are
obable.

The fact that not one of the white marlin tagged in
e western North Atlantic has been recaptured in
ner areas, and the low longline catch rates in the
uatorial area (Ueyanagi et al., 1970; Hayasi et al.,
'70; Mather et al., 1972), as well as the widely
parated spawning areas (see 3.16), indicate that the
arlin stocks of the North and South Atlantic are dis-
ict. The lack of tag returns showing migrations from
e western to the eastern North Atlantic is less con-
usive as an indication of stock separation, since the
ngline and sport fishing efforts in the latter area are
ght.

Robins (1974) cites Ueyanagi et al. (1970, appendix,
gs. 2j, k, l) as an indication that the white marlin
nich concentrate off Gibraltar in late summer may
ove progressively south along the coast of Africa in
e fall to about lat. 5°N.

The monthly distribution of longline catches in the
outh Atlantic (Fig. 5) shows a heavy concentration of
hite marlin off Brazil in September-February and a
sser one off western Africa in June-July, with con-
derable areas of high catch rate in various mid-
ean areas in July and October-February (Hayasi et
., 1970; Mather et al., 1972), but it is not certain
at these occurrences represent the migrations of a
ngle group of fish.

Most fish migrations relate to spawning or feeding.
ne summer concentrations of white marlin in the
orth Atlantic (Venezuela, Gulf of Mexico, Cape
atteras-Cape Cod area, and off Gibraltar) are for
eding (see 2.3), and occur in areas which are highly
oductive, at least in comparison with the surround-
g waters. The spring concentration in the vicinity of
e Greater Antilles, the Bahamas, and Florida is for
awning (see 2.3 and 3.16). Tagging has shown that
e group of white marlin migrates from a summer
eding area (Cape Hatteras-Cape Cod) eastward and
uthward to a wintering area (off northern South
nerica) whose attractive properties are not known,
en northward to a spring spawning area (Greater
ntilles-Bahamas) and thence further north to the
mmering area. Tagging also indicates that another
oup of white marlin may migrate from a summer
eding ground in the Gulf of Mexico to winter among
e northwestern Bahamas, spawn in spring in the
hamas-Cuba area, and return to summer in the
lf. The group, which, on the basis of longline catch
stribution (Mather et al., 1972), appears to summer
in the Gulf and winter in the southeastern Caribbean, may also spawn in spring near the Greater Antilles.

In the South Atlantic, the summer concentration off southern Brazil is for spawning (see 2.3 and 3.16). It appears that the marlins move to this area from off northeastern Brazil in October-November and disperse northward in March-May (Hayasi et al., 1970). These authors also note a feeding concentration of large females off Africa around lat. 10°S from June through August.

3.52 Schooling

Generally not considered a schooling fish, white marlin are most often seen as individuals or in pairs "tailing" with only the dorsal lobe of their caudal fins showing. Small schools (5-12 fish), however, are occasionally seen feeding on schools of bait, or tailing, but loose aggregations of numerous fish scattered over fairly large areas are more typical.

At some point in their life cycle the white marlin may come together in schools. De Sylva and Davis (1963) proposed that they may school according to size or sex at various seasons of the year. Little concrete evidence is available, however, for determining the extent of time a school may remain together or the size such schools may attain.

3.53 Responses to stimuli

As discussed previously, temperature seems to play a crucial part in the response of white marlin. Their disappearance from the Ocean City fishery during a temperature drop of a few degrees and return in numbers similar to catches before the drop as soon as the temperature returned to normal (Earle, 1940) is one example. Similarly the white marlin arrive along the east coast of the United States during the warm summer season and move offshore and to the southward as fall and cold water approach (Mather et al., 1972). Gibbs (1957) noted the relationships between the distribution of white marlin in the Gulf of Mexico and the 75°F (23.9°C) isotherm (see 2.3).

Hayasi et al. (1970) note that the two separate areas of high hooking rates for white marlin in the North and South Atlantic both shift northward during May-October and southward during November-April. In other words, they both move to higher latitudes as the water warms and to lower latitudes as it cools.

De Sylva and Davis (1963) performed a most exhaustive environmental study of white marlin grounds from Atlantic City, N.J. to Ocean City, Md. They found that temperature, salinity, oxygen content, and plankton volumes all played a role in the presence of the white marlin as explained in 2.3.

If marlin seek certain optimum environmental conditions at one time or another, they may swim in deeper layers when surface conditions are not favorable. In such a case types of fishing gear used will

play a role in determination of their presence. For example, longline gear may be necessary to catch the deeper strata of fish while trolling gear (baits, artificial lures, and teasers) attract those near the surface.



Figure 11.—Size composition of white marlin by season and area. (From Ueyanagi et al., 1970, Fig. 19 and caption.) (Area indications and legend added.)

The behavior of white marlin toward trolling b is very indicative of his intentions according to J (1964). If the fish is pale green and placidly follow the bait, he will probably not attack. However, if h dark-colored and his fins become a shiny bright b a reaction known to fishermen as "lighting up," will strike immediately. The raising of the first do fin is also considered to be an indication of exc ment.

4 POPULATION

4.1 Structure

4.11 Sex ratio

Extensive sampling of oceanic catches (Ueyanag al., 1970) shows that the sex ratio of white ma varies considerably with season and area (Fig. Males appear to be predominant in most areas exc in three cases: north of lat. 30°N in May-Octo from lat. 10°N to 30°N in November-April and fr lat. 5°S to 30°S, and east of long. 15°W in M October. The proportion of 2,037 males to 1, females in the area from lat. 10°S to 30°S and wes long. 15°W during November-April, the area period in which the major spawning of South Atla white marlin evidently occurs (Ueyanagi et al., 19 Hayasi et al., 1970), is interesting.

Sex ratios of white marlin collected in insh waters also varied considerably according to areas seasons. De Sylva and Davis (1963) examined caught between Atlantic City, N.J., and Ocean C Md., during the summers of 1959 and 1960. T found 17 males and 41 females in 1959, and 50 ma and 58 females in 1960. "The deviation from the pected 1:1 ratio is highly significant in 1 ($X^2 = 6.63$; $P < 0.001$), non-significant in 1 ($X^2 = 0.44$), and probably significant for the combi years ($X^2 = 5.02$; $P = 0.02$)," Krumholz (1958) fou 20 males and 22 females in a sample caught Bimini, Bahamas, in April 1956. Nakamura and Ri (1972) reported that females definitely outnumbe males in samples examined at four ports in the Gul Mexico during May-October:

	1967 M:F	1968 M:F	1969 M:F	1970 M:F	1971 M:F
uth Pass, La.	20:46	15:59	4:25	4:16	3:12
nsacola, Fla.					2:16
nama City, Fla.					7:13
stin, Fla.					19:91

ien (1964), however, stated that the majority of the hite marlin taken off Venezuela, mainly in August-ctober, were males.

4.12 Age composition

No age determinations were found in the literature.

4.13 Size composition

The length composition of the white marlin popula-on as a whole is best illustrated by Figure 11, which ows the length frequencies for samples of longline itches in various seasons and oceanic areas.

De Sylva and Davis (1963, Figs. 1-5) present weight equencies for white marlin caught in the sport sheries off Ocean City in 1940, 1941, and 1945-59; f Atlantic City in 1958 and 1959; between Ocean ity and Atlantic City in 1959 and 1960; off Puerto ico in 1950-57; and off the Bahamas in 1956. They ate that if these marlin belong to the same popula-on, it seems that marlin may school together accord-g to size or sex at various seasons.

The maximum documented size which we have ound for white marlin was 161 pounds (73.2 kg) and 8 et 8 inches (264 cm)—the present world record rod nd reel catch from off Miami Beach, Fla., in 1938 Anon., 1972a).

We found no information on the density of size roups, beyond the size composition data and catch ates shown above.

De Sylva and Davis (1963) showed that, at small zes [less than 68 inches (173 cm) fork length] female hite marlin tended to be heavier than males of the ame length (Fig. 12), according to the length-weight rmula (see 3.43):

$$\text{Males:} \quad W = 6.0 \times 10^{-4} L^{3.6}$$
$$\text{Females:} \quad W = 4.6 \times 10^{-3} L^{3.0}$$

The constants were derived from measurements of ostspawning fish taken from June through eptember between Atlantic City and Ocean City.

4.2 Abundance and Density (of Population)

4.21 Average abundance

Since no estimates of population size are available, the actual average abundance is unknown.

4.22 Changes in abundance

Local changes in abundance caused by hydrographic conditions were discussed in 2.3.

The apparent relative abundance of white marlin, as indicated by catch rates of the Japanese longline fishery in 1958-66 (Ueyanagi et al., 1970, Fig. 26), declined slightly after reaching a peak in 1962. More recent information (Table 6), however, shows that this net downward trend has continued through 1970, resulting in a decline from a maximum of 2.06 fish per 1,000 hooks in 1962 to 0.80 fish per 1,000 hooks in 1970. Important declines have occurred in the areas (Fig. 13) in which the largest catches were taken: from 4.34 in 1962 to 1.86 in 1970 in BAH; from 10.77 in 1967 to 1.20 in 1970 in RIO; from 2.31 in 1966 to 1.00 in 1970 in NOW; and from 2.44 in 1966 to 0.62 in 1970 in GUI. S. Ueyanagi (pers. commun.) has stated that the average size of the white marlin taken in the fishery has also declined with the catch rates. This is an additional indication that the fisheries have actually caused a decline in the abundance of the white marlin stocks.



gure 12.—Length-weight relationship of white marlin, *etrapturus albidus*, caught between Atlantic City, N.J. and cean City, Md., July-September 1959 and 1960. Males are presented by dotted line and closed circles; females are presented by solid line and open circles. (From de Sylva and avis, 1963, Fig. 6 and caption.)

Table 6.—Catch rates (numbers of fish/1,000 hooks) for white marlin of the Japanese Atlantic longline fishery, 1956-70, by and areas. Note: 0.00 =<0.005 but>0; 0 = effort but no catch; - = no effort. Data furnished by J. P. Wise (pers. commun. Figure 13 for areas.

Year	GM	NOW	NOE	CAR	GUI	CV	GG	BAH	BEN	RIO	To
1956	-	-	-	-	0.04	-	-	0	-	-	0.
1957	0	-	-	0	0.57	0.24	0.05	0	-	-	0.
1958	-	-	-	0	0.19	0.31	0.01	0.15	-	-	0.
1959	-	0	0	0.24	0.13	0.35	0.09	1.73	0	3.00	0.
1960	-	0	0.09	0.06	0.33	0.09	0.03	2.06	2.70	0	0.
1961	-	0.40	0.11	0.43	0.70	0.26	0.70	4.24	2.21	0	1.
1962	-	0.66	3.37	3.49	1.10	0.61	1.13	4.34	3.40	2.50	2.
1963	1.90	1.55	1.05	2.00	1.11	0.85	0.25	4.09	2.91	1.99	1.
1964	3.63	2.10	0.65	2.94	1.32	0.78	0.20	2.75	0.64	7.29	1.
1965	8.02	1.66	0.41	2.07	1.35	0.77	0.08	3.86	0.35	6.22	1.
1966	5.02	2.31	0.40	5.10	2.44	0.59	0.05	2.58	0.21	4.72	1.
1967	4.50	1.06	0.23	6.44	0.94	0.63	0.08	2.46	0.34	10.77	1.
1968	2.29	1.23	0.05	3.68	0.83	0.43	0.08	3.76	0.04	7.19	1.
1969	3.74	1.64	0.07	5.75	0.90	0.49	0.05	2.92	0.15	1.32	0.
1970	5.01	1.00	0.41	1.77	0.62	0.32	0.06	1.86	0.03	1.20	0.
Fish/1,000 hooks	4.14	1.61	0.45	3.33	1.07	0.58	0.30	3.34	0.71	4.33	1.
N fish × 10^3	26.5	113.7	4.9	56.6	97.3	50.6	26.6	193.9	61.4	154.1	784.
N hooks × 10^6	6.4	70.8	10.9	17.0	90.8	86.8	89.9	58.1	86.9	35.6	553.



Figure 13.—Division of the Atlantic Ocean. (From Wise and Davis, 1973, Fig. 1 and caption.)

No information was found on the effects of food competition, predation, and natural fluctuations on the abundance of the species.

4.23 Average density

No information was found in the literature.

4.24 Changes in density

Seasonal and areal variations in density, as indicated by longline catch rates, are shown in Figures 4, 5, and 7.

Some indications of seasonal changes in den with depth are given by the relative success of long and rod and reel fishing, and also by the availab of white marlin to longline fishing in situation which they are not seen on the surface. Off the coast between Cape Hatteras and Cape Cod, sur trolling is effective in summer, when white marlin concentrated on or at the edge of the continental sl and longline fishing is poor. In late September October, the fish have left the shelf (Mather et 1972) and longline catches in the adjacent deep wa are at their peak (Fig. 5). A similar situation exist Venezuela, where surface (sport) fishing is at its p from August-October, while longline results relatively poor. Longline fishing, on the other han good in the area in December and February when white marlin are taken on rod and reel (Fig. 5, longline catch rates; Jaen, 1964, for sport fish success).

A generally deeper distribution in oceanic tha coastal waters is indicated by the lack of visual ob vations of white marlin in deep, oceanic waters, e in areas and seasons in which longline fish produces good catches.

4.3 Natality and Recruitment

4.31 Reproduction rates

Nothing found in the literature.

4.32 Factors affecting reproduction

Nothing in the literature.

4.33 Recruitment

No information on rates, seasonal patterns, or annual variations of recruitment was found in the literature, nor was any information on the relation of recruitment to stock size and reproductive rates.

The overlapping migratory patterns found for different groups of white marlin (Mather et al., 1972) suggest that some groups may recruit adult fish from others, in addition to their recruitment from spawning.

The length (from orbit to fork of tail) of white marlin at first capture in the Japanese longline fishery varies with season and area (Fig. 11), but is usually in the 90- to 120-cm range. The length (from tip of lower jaw to fork of tail) at first capture in the sport fishery off the middle Atlantic coast of the United States is about 140 cm for males and 155 cm for females (de Sylva and Davis, 1963). Erdman (1956) reported that the smallest white marlin that he had observed at Puerto Rico was 22 pounds (10 kg). Among the white marlin examined by Krumholz (1958) at Bimini, Bahamas, the smallest male was 20 pounds (9 kg) and the smallest female 45 pounds (20 kg). Nakamura and Uchiyama (1972) reported that the smallest white marlin taken in each of three areas off northwestern Florida in 1971 weighed 31-42 pounds (14-19 kg), and that the smallest taken off South Pass, La., in each year from 1965-66 weighed 29-39 pounds (13-18 kg).

4.4 Mortality and Morbidity

4.41 Mortality rates

See 3.32.

4.42 Factors causing or affecting mortality

Little is known about the effect of natural factors such as predators, food abundance, and environmental conditions (see 3) on the mortality of the white marlin. The changes in its apparent relative abundance (see 4.22) which have occurred since the development of extensive oceanic longline fisheries, however, indicate that fishing is an important cause of mortality.

4.43 Factors affecting morbidity

See 3.35.

4.44 Relation of morbidity to mortality rates

No information appears in the literature.

4.5 Dynamics of Population (as a Whole)

De Sylva and Davis (1963) studied the dynamics of the Ocean City-Atlantic City sport fishery up until 1960. Assuming similar effort over the years and similar median weights for all fish caught whether boated or released, they found no apparent regular decrease in median weight as the number of marlin caught increased over the years observed. Thus, they concluded there was no reason to believe that angling had affected the stocks.

The present indications that the apparent abundance of the stocks and the average size of the fish taken are decreasing, however, show that studies of the dynamics of the population should be undertaken.

4.6 The Population in the Community and the Ecosystem

As the white marlin is an oceanic species, no specific study has dealt with the ecology of its wide range of distribution. De Sylva and Davis (1963) discuss the relationship of the white marlin with the round herring. On the basis of frequency of round herring in the marlin stomachs, they determined that areas of subsurface plankton concentration may be an important factor of the white marlin community. Squid, too, are a big item in the marlin's diet.

Certain physical features of the oceanic environment are also pointed out by these authors. Salinity, temperature, and oxygen patterns seem to play a role in the location of marlin grounds. Submarine ridges also seem to identify marlin feeding areas. This may be due to an upwelling of plankton attracting the fish and squid on which the marlin feed.

Being members of the holoepipelagic community (Parin, 1968) they perfer the isothermic surface layer of the North subtropical, tropical, and South subtropical regions of the Atlantic and adjacent seas. They reside above the main thermocline but during feeding excursions may pass into its upper horizons.

In the epipelagic environment the white marlin is located on the sixth of seven trophic levels (Fig. 14) beneath only the most powerful and swiftest sharks, *Carcharodon* and makos (Parin, 1968).

Fluctuations in populations occur in a cyclical manner according to the seasonal migration and distribution discussed in 2.

5 EXPLOITATION

5.1 Fishing Equipment

5.11 Gears

The types of fishing gears used for the white marlin vary somewhat from one area to another but all employ the basic hook and line technique. The major classifications are rod and reel, handline, and longline.

In the early days of sport fishing for white marlin the equipment (depending on the experience of the angler) consisted of linen line, varying between 9 and 18 thread (27- to 54-pound [12- to 25-kg] test), a rod with a 6- to 12-ounce (170- to 340-g) split bamboo tip,



Figure 14.—Diagram of the trophic links among epipelagic fishes (from Parin, 1968, Fig. 49 and caption).

A—subarctic waters; level I—phytoplankton (1); level II—phytophagous zooplankton: euphausiids (2), copepods (3), Pteropoda (4); level III—predatory plankton and planktophagous fishes: fish juveniles (5), saury (6), hyperiids (7), myctophids (8); level IV—nektonic predators: Pacific salmon (9), squid (10), lancetfishes (11); level V—large nektonic predators: mackerel shark (12).

B—antarctic waters: level I—phytoplankton (1); level II—copepods (2), euphausiids (3); level III—hyperiids (4), planktophagous fishes (5); level IV—predatory fishes (6).

C—tropical waters: level I—phytoplankton (1); level II—euphausiids (2), copepods (3), shrimp (4); level III—fishes of the "shifting layers" (5), flyingfishes (6), hyperiids (7), "subsurface" lanternfishes (8), moals (9); level IV—small deepwater ichthyophages (Chiasmodon, etc.) (10), nyctoepipelagic predators (snake mackerel) (11), squid (12), Corphaena (13); level V—tuna (14), lancetfishes (15); level VI—marlins (16), medium-sized sharks (17); level VII—large sharks (18).

nd a 4/0- or 6/0-star drag reel (Farrington, 1937). The tandard hooks were 7/0 or 8/0. Today's equipment ermits greater enjoyment through the use of lighter ackle. Experienced anglers usually use 20- or 30- ound (9- or 14-kg) test synthetic (nylon monofila- ıent or braided dacron) line and a 2½/0, 4/0, or 6/0 eel. The more expensive modern reels with lever drag ontrols are preferred, but the older and less costly tar drag type is still adequate. Leaders are at least 3 ı long and of about 60- to 100-pound (27- to 45-kg) est. They may consist of nylon monofilament, tainless steel wire, or nylon coated stainless steel able. A combination of long length of monofilament r coated cable followed by a short length of wire or oated cable ("Venezuelan rig") is very popular. here is also a trend toward using smaller hooks.

Surface trolling baits for white marlin take on many ɔrms (Brooks, 1968). Squid is generally regarded as he best bait off Ocean City with balao, mullet (15-18 m in length), eels and strip baits cut from fish also eing used. Balao and mullet are the most popular in outhern waters. White marlin will also hit artificial ures such as squids, feathers, pork rind strips, cedar nd bone jigs, and spoons.

An important, but not essential, accessory for troll- ng is the outrigger or tangon, a long bamboo, fiber lass, or aluminum pole which is swung outboard to n angle of about 45° from the vertical when in use. he line, after the bait has been let out to the desired istance, is set in a clip which is hauled to the end of he outrigger. This gives an attractive action to the aits in an area outside the direct wake, as well as eeping them separated. It also serves to give the fish "drop back" when the initial strike pulls the line out f the clip, perhaps making the fish believe that he as stunned his prey and certainly giving him time to ake it back into his mouth so that the chances of ooking him are better. When fishing "directly" without an outrigger) the angler must be constantly lert and ready to give the necessary "drop back" y releasing the drag on his reel at the critical mo- ıent.

Handline fishing is commonly practiced by com- ıercial fishermen in the Caribbean. Called cordeles" by the Venezuelans, the handlines basical- y consist of one or more hooks fastened to a strong ire leader which is in turn attached to the handline ıade of hemp or other suitable material (Fiedler, obel, and Lucas, 1947). The fishermen drift in cer- ain areas with their lines at various depths waiting or a fish to strike.

The Cuban criollo line (palangre criollo) is a odified version of the handline. It consists of three ines of different lengths, each attached to a float and onnected to each other and to the boat or an iden- ifying buoy by a horizontal line. The interval etween the floats is about 12-15 m. The standard set onsists of 10 of the 3-hook outfits placed about 90 or 80 m apart in a straight line (Farrington, 1971).

The most widely used and productive commercial method for the capture of white marlin is the Japanese longline. Known as "palangre" or "palangre japonais" to Cuban and Venezuelan fishermen, it consists of a horizontal mainline to which vertical branch lines, each consisting of a short length of line, a leader (often of two components), and a hook, are attached at regular intervals. The basic unit is a "basket," which consists of a specified length of mainline with a specified number of branch lines attached. Enough baskets are fastened together to form a set of the desired length. The fishing depth is determined mainly by the length of the dropper lines which connect the mainline, at each junction between "baskets," to buoys (Captiva, 1955; Yoshida, 1966; Gottschalk, 1972). In a modernized variation of the longline, designed to increase the speed of operation and reduce manpower requirements, a continuous mainline is stored on a power-driven reel or drum, and branch and float lines are attached as the mainline is set and removed as it is retrieved (Captiva, 1955; Anon., 1966). The physical characteristics of the longline gear vary according to the sizes and species of the intended catch. The fishing depth is also selected to take advantage of the observed hydrographic conditions and the known habits and preferences of the fish whose capture is desired.

Since the white marlin is very sensitive to water temperature, surface thermometers are useful to fishermen who troll on the surface, whereas bathythermographs are more helpful to longline and handline fishermen.

5.12 Boats

Various types of boats are used for white marlin fishing ranging from the large oceangoing longliners and some very expensive sport fishing craft down to outboard motorboats and small 4.5- to 6-m wooden sailboats and rowboats of the Caribbean area.

Power, too, ranges from a pair of oars or lateen sail, through outboard motors of all sizes, to the large expensive engines needed to power high-speed sport fishermen or 60-m longliners.

The sport fisherman needs only a roomy cockpit and enough power and range to take him where he wants to go and troll at the required speeds (Jaen, 1964). Everything else is surplus equipment depending on how comfortable he wants to be and how fast he wants to travel. Appropriate models are available from the 5.5-m open outboard to the luxurious, expensive, twin engine sport fishermen.

Commercial vessels range from small rowboats or sailboats to converted trawlers and snapper schooners rerigged for longlining (Bullis, 1955) to the specialized Japanese fishing vessels described by Yoshida (1966).

5.2 Fishing Areas

5.21 General geographic distribution

See 2 and 5.22.

5.22 Geographic ranges

White marlin fisheries range from the coastal waters out to mid-ocean areas. The longline fishery, however, is the only one operating much beyond the continental shelves. Most of the sport fishermen and the smaller commercial boats fish from 1 km off the coast to the edge of the continental shelf.

In the area from Cape Hatteras, N.C., to Cape Cod, Mass., white marlin are sought by sport fishermen traveling as far as 130 km offshore to the canyons along the edge of the continental shelf. Fish have been captured as far north as Nova Scotia but these are probably strays that wandered further to the north than usual in especially warm summers (Morrow, 1965). The major center of abundance in this region seems to be from off Cape Hatteras to off Atlantic City (Mather et al., 1972). The Jack Spot, a shoal 36 km southeast by south of Ocean City has become internationally known (Farrington, 1937; Migdalski, 1958).

Havana, Cuba, and the northwestern Bahamas are other areas of intense fishing. The Cuban commercial fishermen travel only from 1 to 6 km offshore before setting their lines in the Straits of Florida (Farrington, 1971) and sport fishermen at Walkers Cay, Cat Cay, and Bimini in the Bahamas, and at Miami can be putting out lines within minutes of leaving the dock.

The sport and small-boat commercial fisheries off Venezuela are centered about 24 km offshore. Farrington (1949a) noted the abundance of white marlin off Caracas but realized the need for improved equipment and crews with greater maturity and experience before it would be utilized. By 1958, Migdalski predicted that the area north of Venezuela would be another Ocean City and Jaen (1964) calls it the paradise for white marlin anglers.

A newly developing area of interest which shows a promise of abundance is in the northern Gulf of Mexico (Mather et al., 1972). White marlin also occur along the eastern coast of South America as far south as Rio de Janeiro, off the Azores, Madeira, Portugal, southern Spain, and South Africa.

5.23 Depth ranges

Parin (1968) reports that during feeding excursions marlin may pass even into the upper levels of the main thermocline 200-250 m in depth. Their usual range, however, varies between the surface and 20-30 m. Studies of the results of the Japanese longline fishery also indicate that almost all billfish catches are made while the line is moving near the su (Watanabe, 1961).

5.24 Conditions of the grounds

De Sylva and Davis (1963) give the most deta description of conditions on white marlin groun See 2 and 4.6.

5.3 Fishing Seasons

5.31 General pattern of seasons

See 2 and 3.5.

The white marlin ranges farther into the tempe zones during the warm seasons and congreg seasonally in certain coastal areas. Along the eas United States, white marlin are abundant during warm season from Cape Hatteras north to Cape as well as in the Gulf of Mexico. During the co seasons the marlin congregate in the more soutl waters of the Bahamas—peak in spring— Venezuela—peak August-October—where they be found at all seasons of the year (Mather et 1972).

The offshore fishery reflects the same gen pattern with longline catches being greater in n and south temperate waters during their respec warm seasons and higher in tropical areas in the ones (Ueyanagi et al., 1970; Wise and Davis, 19 The middle of fall marks the beginning of the m southward movement from North Atlantic fishe after the marlin have moved offshore from the c (Mather et al., 1972).

5.32 Dates of beginning peak and end season

See 2 and 3.5.

In the middle Atlantic coast sport fishery, w marlin arrive at Cape Hatteras in June and m north all the way to Cape Cod with fishing lasting til October in some areas. The peak usually progre northward starting off Oregon Inlet, N.C., in July moving up to Ocean City in late July and early Au and on up to Montauk, Long Island, in late Au before the fish move offshore in September (Mathe al., 1972).

Although some white marlin are available to s fishermen off the Bahamas throughout the year, t appear off the northwestern Bahamas, north Cuba, and southeastern Florida in greater numbe January and stay until June, with the best fis from mid-April to mid-June (Migdalski, 1958).

The sport fishing season for white marlin in northern Gulf of Mexico extends from early June October with peaks off the Mississippi Delta in and in the northeastern Gulf in August (Nakam and Rivas, 1972).

The Venezuelan sport fishery also produces s

te marlin on a year around basis, but the really in-sive fishing occurs from August through October. fish start moving into the western coast in July. ools may be seen off Puerto La Cruz (Distrito eral), moving east progressively until they reach Guaira (the port of Caracas) in September (Jaen, 4).

he seasonal trends in the oceanic longline catches white marlin differ considerably from those of the nore sport fisheries. The former are illustrated in ure 5, on which the following discussion is based. he North Atlantic, most of the successful fishing in ter occurs off the north coast of South America. In ing, the most productive areas are around the West ies and the Bahamas, and in the western Carib-n and the Gulf of Mexico. The latter two areas are only productive ones in the summer. In early fall, ch rates are high in oceanic waters off the middle antic coast of the United States out to long. 55°W, l also in the Gulf of Mexico. Otherwise there is lit-longline fishing success in the fall until catch rates prove in the southeastern Caribbean in Decem-

n the South Atlantic, the most important longline ches occur from September through February off zil. The location and extent of this fishery varies n month to month, gradually shifting southward. ere are also scattered areas of high catch rates in ious months, expecially spring, in mid-ocean, and ther concentration occurs in June-July between 5° and 15°S, long. 5°E, and the African coast.

5.33 Variations in date or duration of seasons.

Nothing appears in the literature dealing with iations in the seasons of offshore longline fish-es.

Several factors have been noted which contribute to iations in the seasons of some of the sport fisheries. there were not an accumulation of small forage ies on which the white marlin could thrive or if the ter temperature were not satisfactory, Farrington 49a) noted white marlin may not occur in the ean City area in quantities. Earle (1940) also show-that records from the Ocean City fishery show a ect relationship between catches and water nperatures. Nakamura and Rivas (1972) reported t the bluer the water in the South Pass, La., and rthwest, Fla., areas the greater the abundance of ite marlin. Current also plays an important role off uth Pass. The Loop Current, which comes through Yucatan Channel up toward South Pass and loops tward to the De Soto Canyon and down through Straits of Florida, holds the fish together and ngs them right up to South Pass when it is strong, when it is weak and failing, the fish scatter all r the Gulf of Mexico (H. B. Howcott, pers. com-n.) The dates and durations of seasons are in-enced by the times when these factors become orable or unfavorable.

5.4 Fishing Operations and Results

5.41 Effort and intensity

The effort of longline fisheries is easily measured by the number of hooks set, which is usually recorded in thousands. The catch per unit of effort is usually recorded in terms of fish per 100 hooks or per 1,000 hooks. In handline or troll fisheries, effort is usually recorded by boat-days, although greater refinement may be achieved by considering boat- or line-hours. The catch per unit of effort is recorded in the terms in which the data are available.

The effort of Japanese longline vessels in the Atlantic, 1956-70, in numbers of hooks fished by years and areas (Fig. 13) is shown in Table 7. These correspond with the catch rates for white marlin shown in Table 6. Logbook data for annual fishing effort for the years 1956-70 by months and 5° (lat. and long.) quadrangles have been published by Shiohama et al (1965) and the Research Division of the Fisheries Agency of Japan (1965, 1966, 1967a, 1967b, 1968, 1969, 1970, 1971, 1972). Factors to adjust these data to estimates for the whole fleet are supplied by Wise and Davis (1973).

Since comparable data are not available for other longline fisheries, it is difficult to estimate total fishing effort. The Japanese effort reached a peak of nearly 100 million hooks in 1965 and has ranged between about 30 million and 42 million from 1967 through 1970. The numbers of Canadian, Japanese, Korean, and Taiwanese vessels which fished in the Atlantic in various years from 1960 through 1971 may give some idea of the relative effort of these fisheries (Miyake and Tibbo, 1972). While the Japanese effort from 1966 through 1970 was well below the 1965 peak, the increased participation of Korean and Taiwanese vessels kept the total effort high in most of those years and the number of Japanese vessels rose sharply in 1971. The Canadian fishery, which was for swordfish, *Xiphias gladius*, and a similar United States fishery, for which we have no effort data, were virtually terminated in 1970 because of the discovery of heavy metals in swordfish. Likewise, we have no effort data for Scandinavian longline fisheries, which were primarily directed toward the porbeagle shark, *Lamna nasus*, but also caught white marlin like the Canadian and United States fisheries (neither country of which marketed this species).

There are several Latin American commercial fisheries directed toward tunas and billfishes—all of which are marketed in the respective countries. In addition to its local small-boat fishery, Cuba has 4 longliners of about 400 gross tons each and 19 of about 700 gross tons each (Ferrer Guzmán, Carrillo Cárdenas, and Jimenez Guerras, 1972). These vessels fish in the tropical Atlantic, the Gulf of Mexico, and the Caribbean Sea. The Venezuelan longline fishery based at Cumaná originated in 1959 and leveled off at about 40 vessels of from 20 to 300 gross tons which fish

Table 7.—Numbers of hooks (in thousands) fished by the Japanese longline fishery in the Atlantic, by years, 1956-70, and areas (see Fig. 13). Data furnished by J. P. Wise (pers. commun.).

Year	Fishing effort										
	GM(0)	NOW(1)	NOE(2)	CAR(3)	GUI(4)	CU(5)	GG(6)	BAH(7)	BEN(8)	RIO(9)	Tota
1956	0	0	0	0	129	0	0	2	0	0	13
1957	0	0	0	5	867	1,133	1,328	42	0	0	3,37
1958	0	0	0	290	3,534	2,019	1,806	353	0	0	8,00
1959	0	4	6	258	4,300	5,419	3,009	2,294	4	17	15,31
1960	0	2	127	330	4,704	5,649	5,609	3,918	379	7	20,72
1961	0	55	275	92	1,616	5,416	10,942	4,833	3,346	84	26,66
1962	0	4,177	161	1,324	12,260	8,076	9,884	12,028	6,012	997	54,92
1963	433	9,478	233	2,560	10,100	9,163	9,182	6,003	5,049	2,803	55,00
1964	2,556	17,098	1,830	2,628	17,172	12,956	5,191	11,387	8,912	5,269	84,99
1965	519	14,231	3,535	1,909	17,192	14,127	18,273	5,761	16,775	5,258	97,58
1966	263	5,997	1,244	4,538	4,168	7,585	6,905	7,446	12,774	2,871	53,79
1967	345	4,495	774	175	3,824	5,930	6,230	1,814	5,638	1,928	31,15
1968	652	3,459	645	806	3,523	3,769	4,252	1,169	8,515	3,411	30,20
1969	158	3,382	268	1,061	3,994	3,784	4,262	451	7,254	5,063	29,67
1970	1,460	8,382	1,796	977	3,413	1,809	3,032	592	12,251	7,842	41,55
Total	6,388	70,761	10,894	16,952	90,796	86,835	89,906	58,092	86,909	35,551	553,08

mainly in the eastern Caribbean and adjacent Atlantic waters (Mihara and Griffiths, 1971). Five Brazilian longliners set 598,000 hooks in 554 fishing days in 1971 (Paiva, 1972). Miyake and Tibbo (1972) show only small (less than 50 ton) annual catches of Atlantic billfishes for Mexico (1964-70) and the Union of Soviet Socialist Republics (1967-71), but we found no further information on these fisheries. Moroccan, Portuguese, and Spanish trap and hook and line fisheries also take some white marlin (Furnestin et al., 1958; Rodriguez-Roda and Howard, 1962; Robins, 1974).

The average geographical distribution of the Japanese longline fishery in terms of intensity is shown by quarters of the year by Wise and Davis (1973, Fig. 3). Most of the effort is in tropical waters. Extensions of intensive effort to latitudes higher than 20° occur in the central North Atlantic during each quarter and in the western North Atlantic, including the Gulf of Mexico, in the second and third quarters. In the South Atlantic, such extensions occur mainly on the western side in the first quarter and on the eastern side in the second and third quarters. In the fourth quarter, the southward extension is nearly oceanwide.

Seasonal changes in effort are probably caused mainly by the seasonal distributional cycles of the species being fished, which are in turn affected by various factors (see 2.2, 2.3).

Year-to-year changes in effort are probably caused, for the most part, by the interaction of economic and biological factors. The basic determinants are, for capitalistic countries at least, the market value of the fish and the costs of catching them. Naturally, costs depend on wage scales and the costs of boats, equipment, and bait, as well as the availability of the fish. Since the white marlin is usually an incidental catch, however, the commercial fishing effort to which it is subjected is usually determined by the economics the fisheries for more important species, such as yellowfin tuna, the albacore, *Thunnus alalunga*, a the bigeye tuna, *T. obesus*.

The effort of sport fisheries is much more difficult determine. Hundreds of charter and thousands private sport fishing boats fish for white marlin fr Cape Hatteras to Cape Cod, off southeastern Flori in the Gulf of Mexico, off the Bahamas and the W Indies, and off Venezuela. The number of boats is d ficult to estimate, as many move from place to pl seasonally. The actual effort for white marlin is ev more difficult to estimate, since other species usually being fished for concurrently, and boats oft divide their effort, over the season or the year, amo various types of fishing.

Nakamura and Rivas (1972) listed the effort pended by members of the New Orleans Big Ga Fishing Club off South Pass, La., 1967-71, and anglers from the Florida ports of Destin, Pana City, and Pensacola in 1971. The effort var between 2,339 and 5,801 boat-hours per season South Pass and totalled 7,890 boat-hours in d season off the three Florida ports.

When the white marlin sport fishery at Ocean C began in 1936, there were 12 charter boats and 1939, there were 39. After the lull during World W II, there were about 40 boats during the period fr 1946 through 1953 (June and Reintjes, 1957). Sin then the number increased to the present level about 70.

The number of boats in the Venezuelan int national fishing tournaments (in most of wh sailfish and blue marlin are taken, as well as wh marlin) increased from the low to middle 20's in earlier years to 43 in 1970 (Anon., 1971).

The number of boats in the Atlantic City Ma

urnament has been as high as 200 in 1964 and 1965, t has declined to 120 to 130 in recent years (Atlantic y Public Relations, pers. commun.).

5.42 Selectivity

There appears to be little difference between the rious hook and line gears as to the sizes of white rlin taken (see 4.13). Full selectivity in regard to e species of fish taken by the various gears is also ficult. With deep fishing methods (longline and ndlines), selection may be attempted by the areas, asons, depths, and environmental conditions in ich fishing is carried out. In surface fishing, the eas, seasons, and environmental conditions are also strumental in determining the species caught. This ay also be influenced, however, by fishing tactics, ch as type and size of bait or line and hook, position bait or lure relative to boat, and trolling speed.

5.43 Catches

The total annual yield of white marlin as recorded the FAO Yearbook of Fisheries Statistics (FAO, 71) declined from about 5,000 tons in 1965 to about 00 tons in 1967-70. These totals, however, include gures for Japan and China (Taiwan) only (Table 8); e total catches must actually be considerably eater.

The Japanese white marlin catch attained highs of 3,415 fish and 4,600 tons in 1965, and has varied tween about 27,000 and about 43,000 fish weighing m 700 to 1,000 tons in the years 1967-70 (Table 8; J. Wise, pers. commun.). The Taiwanese fishery took m 100 to 500 tons of white marlin annually in 1966- (Table 8). No data are available for white marlin tches of Korea, Cuba, Venezuela, and Brazil, but gures for the yearly catches of "others" (fishes other an albacore, yellowfin, bigeye, and bluefin tunas) by Korea, and of "billfishes" (marlins, sailfish, and spearfishes) by Cuba, Venezuela, and Brazil are provided by Miyake and Tibbo (1972). These authors' data indicate that white marlin comprised an average of 1.9% (by weight) of the total Japanese longline catches in 1967-71. Assuming that the tonnage of white marlin formed a similar percentage of the Korean catches, the estimated yearly totals increased from about 210 tons in 1967 to 705 tons in 1971. The Cuban catches of "billfishes" in 1961-71 ranged from 300 to 1,700 tons; those of Venezuela in 1968-71, from 300 to 500 tons; and those of Brazil in 1962-71, from 100 to 200 tons. In the Japanese Atlantic longline fishery, 1964-70, catches of white marlin constituted from 29 to 36%, by weight, of the total catches of marlins and sailfish, with an average of 33% (FAO, 1971). Catches of the Canadian and United States fisheries are not known, since the white marlin caught were discarded as no market exists in those nations. The considerable effort expanded by these fisheries in waters frequented by white marlin, however, must have resulted in numerous captures. The same is true of the Scandinavian longline fisheries which operated in the western North Atlantic in the early 1960's. The white marlin catches of Moroccan, Spanish, and Portuguese fishermen are likewide unknown. The total yearly commercial catches of white marlin in the years 1962-70 probably ranged between 2,000 and 6,000 tons (see 4.22).

Total catches for the numerous sport fisheries could only be estimated, but certain results from some of these areas are available to give indications. The Atlantic City Marlin Tournament catches have ranged from 7 fish in the first 3-day tournament in 1955 to 223 in 1963. Table 9 shows the catches, number of days fished, and number of boats fishing (Atlantic City Public Relations, pers. commun.). The numbers of white marlin caught in the Gulf of Mexico by the

Table 8.—Nominal catches of white marlin by fishing areas and by countries, as reported to the FAO (FAO, 1971).

Atlantic white marlin	**4.0**	**5.0**	**3.0**	**1.0**	**1.0**	**1.0**	**1.0**
Tetrapturus albidus (= *Makaira albida*)							
Atlantic, Northwest	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	--	(0.0)
Japan (. . . D)	0.0	0.0	0.0	0.0	0.0	--	0.0
Atlantic, Northeast	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	--	(0.0)
Japan (. . . D)	0.0	0.0	0.0	0.0	0.0	--	0.0
Atlantic, Western Central	(1.3)	(1.6)	(1.5)	(0.2)	(0.1)	(0.4)	(0.4)
China (Taiwan) (. . . D)	--	--	--	0.1	0.0	0.1	0.1
Japan (. . . D)	1.3	1.6	1.5	0.1	0.1	0.3	0.3
Venezuela (Aguja blanca)	---	---	---	---	---	---	---
Atlantic, Eastern Central	(0.3)	(0.6)	(0.3)	(0.1)	(0.3)	(0.2)	(0.1)
China (Taiwan) (. . . D)	--	--	0.1	0.0	0.0	0.1	0.0
Japan (. . . D)	0.3	0.6	0.2	0.1	0.2	0.1	0.1
Atlantic, Southwest	(1.6)	(2.1)	(1.1)	(0.4)	(0.6)	(0.5)	(0.3)
China (Taiwan) (. . . D)	--	--	--	--	--	0.2	0.1
Japan (. . . D)	1.6	2.1	1.1	0.4	0.6	0.3	0.2
Atlantic, Southeast	(0.3)	(0.3)	(0.2)	(0.2)	(0.3)	(0.2)	(0.3)
China (Taiwan) (. . . D)	--	--	--	0.1	0.2	0.1	0.2
Japan (. . . D)	0.3	0.3	0.2	0.1	0.1	0.1	0.1

Table 9.—Atlantic City, New Jersey, Marlin Tournament, results 1955-70, from the Atlantic City Public Relations Department.

Year	White marlin caught	Number of boats	Number of days fished
1955	7		2
1956	13	97	3
1957	36	100	3
1958	117	110	3
1959	104	180	3
1960	20	228	3
1961	5	182	4
1962	144	135	4
1963	223	165	3
1964	118	200	4
1965	85	200	4
1966	17	140	2
1969		120	3
1970	70	130	4
1971			4
1972		110	4

New Orleans Big Game Fishing Club from 1966 to 1970 were 151, 113, 95, 38, and 22 respectively (Nakamura, 1971), and in Venezuela in 1970, 190 white marlin were caught in 3 days of September from 45 sport fishing boats (Anon., 1970). The Ocean City catches for the first active season of marlin fishing in 1936 were 175 but by 1939 the figure had jumped to 1,343 (Earle, 1940). Table 10 shows the white marlin landings in Ocean City from 1936 to 1971 (June and Reintjes, 1957; Reintjes and Roithmayr, 1960; Ocean City Marlin Club, pers. commun.) In 1972 Ocean City Public Relations reports 837 marlin recorded by the 2nd of October; while boats based at Hatteras and Oregon Inlet, N.C., reported catching 1,212 white marlin in 1971 (Oregon Inlet Charter Boat Association, pers. commun.).

6 PROTECTION AND MANAGEMENT

6.1 Regulatory (Legislative) Measures

6.11 Limitation or reduction of total catch

At the present time no limitations have been put on the white marlin fishery by any legislative body. The need for such measures is being studied and may be considered in the future by the International Commission for the Conservation of Atlantic Tunas. This body has been given regulatory authority over both tunas and billfisheries to maintain maximum sustainable yield.

6.12 Protection of portions of population

Although no official protection measures are presently in force, conservation of this species is being carried on by sport fishermen in many areas. Ocean City, Md., is one major area where conservation become the rule rather than the exception. In only 84 out of 1,343 fish caught were released into the ocean even though they would serve no tical purpose, but up to July in the 1956 season approximately 800 fish caught were rele (Migdalski, 1958). Continuing this practice, m 60% of the Ocean City catch is released yearly, with tags from the Woods Hole Oceanogra Institution to aid in tracing the migrations o white marlin (Brooks, 1968).

Oregon Inlet, N.C., and southern New J fishermen, too, are deeply involved in this impo conservation effort. Of the 1,212 white marlin ca by Hatteras and Oregon Inlet-based boats in 197 were reported released and many of these were ta (Oregon Inlet Charter Boat Association, pers. mun.).

Other areas where tagging and releasing of fis practiced include Long Island, Cape Cod, Bahamas, Florida, the Gulf of Mexico, Virgin Isl and off Venezuela (Mather et al., 1972). Tab shows the number of white marlin tagged by yea these various areas by cooperating sport fishe and the numbers of tags returned.

6.2 Control or Alteration of Physical Featur the Environment

Nothing found in the literature.

6.3 Control or Alteration of Chemical Featur the Environment

Nothing found in the literature.

6.4 Control or Alteration of the Biolo Features of the Environment

Nothing found in the literature.

6.5 Artificial Stocking

Nothing found in the literature.

7 FISH POND CULTURE

Not applicable.

ACKNOWLEDGMENTS

The preparation of this synopsis was supporte the Office of Sea Grant, National Oceanic and mospheric Administration (NOAA), U.S. De ment of Commerce Grant Nos. GH-82 and 04-3-15 and the National Marine Fisheries Service, NO U.S. Department of Commerce, Contract No. 043-29.

Table 10.—Ocean City marlin history 1936 through 1971, recorded by the Ocean City Marlin Club.

Year	Total caught	Total released	Largest caught	Date first caught	Date last caught
1936	175	NR	112 lb.	NR	NR
1937	200	NR	130 lb.	NR	NR
1938	781	NR	118 lb.	23 June	5 Sept.
1939	1,259		113 lb.	16 June	26 Aug.
1940	134	19	114 lb.	20 July	23 Sept.
1941	107	3	111 lb.	28 June	15 Sept.
1942	No records kept during war years				
1943	" "	" "	" "		
1944	" "	" "	" "		
1945	227	2	92 lb.	13 July	14 Sept.
1946	374	50	96 lb.	6 July	1 Sept.
1947	509	60	110 lb.	27 June	15 Sept.
1948	260	18	100 lb.	25 June	20 Sept.
1949	108	1	90 lb.	22 June	18 Sept.
1950	317	11	115 lb.	20 June	29 Sept.
1951	283	33	103 lb.	25 June	1 Oct.
1952	349	88	123 lb.	29 June	23 Sept.
1953	824	282	114 lb.	28 June	1 Oct.
1954	1,022	525	129 lb.	24 June	8 Sept.
1955	520	279	94 lb.	23 June	24 Sept.
1956	1,616	947	130½ lb.	22 June	19 Sept.
1957	1,062	622	106 lb.	19 June	21 Sept.
1958	616	322	121 lb.	21 June	22 Sept.
1959	1,082	569	100 lb.	24 June	13 Sept.
1960	503	228	115 lb.	20 June	16 Sept.
1961	545	317	95 lb.	23 June	15 Sept.
1962	926	545	115 lb.	15 June	13 Sept.
1963	1,267	688	109 lb.	16 June	18 Sept.
1964	721	370	82 lb.	20 June	18 Sept.
1965	654	325	117 lb.	22 June	24 Sept.
1966	544	256	80½ lb.	1 July	18 Sept.
1967	1,146	721	106 lb.	4 July	10 Sept.
1968	1,735	925	105 lb.	26 June	29 Sept.
1969	2,507	1,696	94 lb.	17 June	1 Oct.
1970	2,098	1,520	91 lb.	12 June	14 Oct.
1971	2,206	1,635	108½ lb.	19 June	10 Oct.

nportant additional support has been received
n the Sport Fishing Institute as well as from many
rt fishing organizations and individual sportsmen.

LITERATURE CITED

SSIZ, L.
843. Recherches sur les poissons fossiles. Poissons cycloides. Neuchâtel 5:160 p.

NYMOUS.
966. Power reel to save labor on tuna vessels (from *Suisan Tsushin*, February 22, 1966). Commer. Fish. Rev. 28(5):52.
968. Relentless sea relinquishes rarity to resourceful researchers. Fla. Conserv. News 3(11):5.
970. Bulletin 29th ILTTA Sept. 24-27 Venezuela. International Light Tackle Association (results of tournament), 3 p. [Processed.]
971. Statistic of thirteen international fishing tournaments at La Guaira-Venezuela. *In* XI Torneo Internacional de Pesca Caraballeda, Venezuela, Sept. 16-19, 1971 (announcement) Caracas, Venezuela, Federacion Venezolana de Pesca. [Processed.]

1972a. World record marine game fishes. Int. Game Fish Assoc., Fort Lauderdale, Fla., 18 p.
1972b. List of open records. Game Fish Union of Africa, Salisbury, Rhodesia.

BINI, G.
1968. Atlante dei pesci delle coste Italiane. Vol. 6. Mondo Sommerso Editrice, Milano, 177 p.

BLOCH, M. E., and J. G. SCHNEIDER.
1801. Systema ichthyologiae, iconibus CX illustratum, post obitum auctoris opus inchoatum absolvit, correxit, interpolavit Io. Gottlob Schneider, Berl., 584 p.

BONAPARTE, C. J. L.
1841. Iconografia della fauna Italica . . . 3, Part 1, Pesci. Rome, Salviucci, n.p.

BROOKS, J.
1968. Salt water game fishing. Harper and Row, N.Y., 340 p.

BULLIS, H. R., JR.
1955. Preliminary report on exploratory long-line fishing for tuna in the Gulf of Mexico and the Caribbean Sea. Part I - Exploratory fishing by the *Oregon*. Commer. Fish. Rev. 17(10):1-15.

Table 11.—Releases (after slash) and returns (before slash) for white marlin, *Tetrapturus albidus*, grouped by major tagging localities. Nearly all were caught on rod and reel (trolling) and tagged by sport fishermen cooperating with Woods Hole Oceanographic Institution.

Year	Hatteras to Chesapeake	Chesapeake to Barnegat	Barnegat to Cape Cod	Oceanic North Atlantic	S.E. Florida and W. Bahamas	West Indies and vicinity	Gulf of Mexico	Venezuela and vicinity	Cozumel and Yucatan	Totals
1954			0/4							0/4
1955		1/116				0/8	0/21			1/145
1956		1/402				0/3	0/8			1/413
1957	0/3	0/140	0/1	0/1						0/145
1958	0/1	0/39	0/1							0/41
1959		0/190	0/10					0/2		0/202
1960		0/96	0/2		0/4	0/1	0/4	0/4		0/111
1961	0/2	2/187	0/10		0/13	0/9	0/11	0/30		2/262
1962	0/30	4/294	0/18		0/41		0/4			4/387
1963	0/75	4/533	0/4	0/3	0/35		0/10			4/660
1964	4/182	8/258	0/1	0/5	1/67		0/13			13/526
1965	0/15	6/258	0/5		0/67	0/5	0/10	2/25		8/385
1966	1/36	9/172	1/64	0/6	1/54	0/4	0/23	4/149		16/508
1967	0/37	6/234	0/6		0/88	0/7	1/46	0/103		7/521
1968	2/100	15/569	1/32		1/95	0/16	0/56	0/16		19/884
1969	8/360	12/829	0/27		2/86	0/18	2/35	2/46		26/1,401
1970	12/320	8/463	1/55		2/49	0/15	0/24	0/17	0/4	23/947
1971	2/247	11/559	0/17		1/57	0/20	0/18	0/95	0/4	14/1,017
1972	0/167	0/164	0/14		0/36	0/10	0/62	0/21	0/1	0/475
Unknown	2/2	4/4					1/1			7/7
Totals	31/1,577	91/5,507	3/271	0/15	8/692	0/116	4/346	8/508	0/9	145/9,041

LDWELL, D. K.
1962. Postlarvae of the blue marlin, *Makaira nigricans*, from off Jamaica. Los Ang. Cty. Mus. Nat. Hist., Contrib. Sci. 53:1-11.
NESTRINI, G.
1861. Sopra una nuova specie di *Tetrapturus*. Arch. Zool. Anat. Fisiol. Genoa 1:259-261.
1872. Pesci d'Italia. *In* E. Cornalia, Fauna d'Italia. Part 3. Vallardi, Milano, 208 p.
PTIVA, F. J.
1955. Preliminary report on exploratory long-line fishing for tuna in the Gulf of Mexico and the Caribbean Sea. Part II - Long-line gear used in yellowfin tuna exploration. Commer. Fish. Rev. 17(10):16-20.
RPENTER, E. J., S. J. ANDERSON, G. R. HARVEY, H. P. HKLAS, and B. B. PECK.
1972. Polystyrene spherules in coastal waters. Science (Wash., D.C.) 178:749-750.
SAREO, A.
1967. Pesca d'altura: bilancio d'una stagione movimentata da due eccezzionali catture. Mondo Sommerso 9:950-952.
BUEN, F.
1950. Contribucion a la ictiologia. 3. La familia Istiophoridae y descripción de una especie uruguaya (*Makaira perezi* de Buen). Publ. Cient. Serv. Oceanogr., Montevideo 5:163-178.
SBROSSES, P.
1938. Sur les poissons-épées du genre *Tetrapturus* Rafin. 1810 rencontrés près des cotes d'Europe. Bull. Soc. Zool. Fr. 63:48-58.
SYLVA, D. P.
1958. Juvenile blue marlin, *Makaira ampla* (Poey), from Miami, Florida, and West End, Bahamas. Bull. Am. Mus. Nat. Hist. 114:412-415.
1963. Postlarva of the white marlin, *Tetrapturus albidus*, from the Florida Current off the Carolinas. Bull. Mar. Sci. Gulf Caribb. 13:123-132.
In press. Family istiophoridae. *In* J. C. Hureau, Check-list of the fishes of the north-east Atlantic and of the Mediterranean.
SYLVA, D. P., and W. P. DAVIS
1963. White marlin, *Tetrapturus albidus*, in the middle Atlantic bight, with observations on the hydrography of the fishing grounds. Copeia 1963:81-99.
RLE, S.
1940. The white marlin fishery of Ocean City, Maryland. U.S. Fish Wildl. Serv., Spec. Sci. Rep. 6, 15 p.
MUNDS, P. H.
1972. Genic polymorphism of blood proteins from white marlin. U.S. Fish Wildl. Serv., Res. Rep. 77, 15 p.
DMAN, D. S.
1956. Recent fish records from Puerto Rico. Bull. Mar. Sci. Gulf Caribb. 6:315-340.
1957. Marlin fishing in Puerto Rican waters. Proc. Second Int. Game Fish Conf., 13 p. Univ. Miami Mar. Lab., Miami.
1958. Foods of pelagic fishes from Puerto Rico and the Bahama Islands. Proc. Third Int. Game Fish Conf., 12 p. Univ. Miami Mar. Lab., Miami.
) (Food and Agricultural Organization of the United Nations).
1971. Yearbook of fishery statistics. Catches and landings, 1970. Vol. 30, 476 p.
RRINGTON, S. K., JR.
1937. Atlantic game fishing. Kennedy Brothers, N.Y., 298 p.
1949a. Fishing the Atlantic, offshore and on. Coward-McCann, N.Y., 312 p.
1949b. Spearfish of the world. *In* B. Vesey-Fitzgerald and F. LaMonte (editors), Game fish of the world, p. 144-155. Harper & Brothers, N.Y.
1971. Fishing with Hemingway and Glassell. David McKay Co. Inc., 118 p.
FERRER GUZMAN, R., E. CARRILLO CARDENAS, and E. JIMENEZ GUERRAS.
1972. Sintesis de las pesqueras Cubanas en el Oceano Atlantico durante el ano 1971. Int. Comm. Conserv. Atl. Tunas, SCRS/72/68.
FIEDLER, R. H., M. J. LOBELL, and C. R. LUCAS.
1947. The fisheries and fishery resources of the Caribbean area. U.S. Fish Wildl. Serv., Fish. Leafl. 259, 211 p.
FISHERIES AGENCY OF JAPAN, RESEARCH DIVISION.
1965. Annual report of effort and catch statistics by area on Japanese tuna longline fishery, 1962. 183 p.
1966. Annual report of effort and catch statistics by area on Japanese tuna longline fishery, 1963. 322 p.
1967a. Annual report of effort and catch statistics by area on Japanese tuna long line fishery, 1964. 379 p.
1967b. Annual report of effort and catch statistics by area on Japanese tuna long line fishery, 1965. 375 p.
1968. Annual report of effort and catch statistics by area on Japanese tuna longline fishery, 1966. 299 p.
1969. Annual report of effort and catch statistics by area on Japanese tuna longline fishery, 1967. 293 p.
1970. Annual report of effort and catch statistics by area on Japanese tuna longline fishery, 1968. 283 p.
1971. Annual report of effort and catch statistics by area on Japanese tuna longline fishery, 1969. 299 p.
1972. Annual report of effort and catch statistics by area on Japanese tune longline fishery, 1970. 326 p.
FOX, W. W.
1971. Temporal-spatial relationships among tunas and billfishes based on the Japanese longline fishery in the Atlantic Ocean, 1956-1965. Univ. Miami, Sea Grant Tech. Bull. 12, 78 p.
FURNESTIN, J., J. DARDIGNAC, C. MAURIN, A. VINCENT, R. COUPÉ, and H. BOUTIÈRE.
1958. Dounées nouvelles sur les poissons du Maroc Atlantique. Rev. Trav. Inst. Pêches Marit. 22(4):379-493.
GEHRINGER, J. W.
1957. Observations on the development of the Atlantic Sailfish *Istiophorus americanus* (Cuvier), with notes on an unidentified species of istiophroid. U.S. Fish Wildl. Serv., Fish. Bull. 57:139-171.
GIBBS, R. H., JR.
1957. Preliminary analysis of the distribution of white marlin, *Makaira albida* (Poey), in the Gulf of Mexico. Bull. Mar. Sci. Gulf Caribb. 7:360-369.
GOODE, G. B.
1881. The taxonomic relations and geographical distribution of the members of the sword-fish family, Xiphiidae. Proc. U.S. Natl. Mus. 4:415-433.
GOTTSCHALK, J. S.
1972. Longlines and billfish. Mar. Fish. Rev. 34(11-12):20-25.
GREY, Z.
1928. Big game fishing in New Zealand seas. Nat. Hist. 28:46-52.
HARVEY, G. R., V. T. BOWEN, R. H. BACKUS, and G. D. GRICE.
1972. Chlorinated hydrocarbons in open-ocean Atlantic organisms. *In* D. Dyrssen and D. Jagner (editors), Nobel Symp. 20, The changing chemistry of the oceans, p. 177-186. John Wiley Sons, Inc., N.Y.

HAYASI, S., T. KOTO, C. SHINGU, S. KUME, and Y. MORITA.
1970. Status of tuna fisheries resources in the Atlantic Ocean, 1956-1967. [In Jap. and Engl.] *In* Resources and fisheries of tunas and related fishes in the Atlantic Ocean, p. 1-72. Far Seas Fish. Res. Lab., S Ser. 3.
HEMINGWAY, E.
1935 Marlin off Cuba. *In* E. V. Connett (editor), American big game fishing, p. 55-82. Derrydale Press, N.Y.
1949. Cuban fishing. *In* B. Vesey-Fitzgerald and F. LaMonte (editors), Game fish of the world, p. 156-160. Harper & Brothers, N.Y.
HIRASAKA, K., and H. NAKAMURA.
1947. On the Formosan spearfishes. Bull. Oceanogr. Inst. Taiwan 3:9-24.
HORN, M. H., J. M. TEAL, and R. H. BACKUS.
1970. Petroleum lumps on the surface of the sea. Science (Wash., D.C.) 168:245-246.
JAEN, R.
1964. Fishing in the Caribbean (version inglesa-Ana Teresa Arcaya de Arcaya). Editorial Arte, Caracas, Venezuela, 228 p.
JONES, E. C.
1971. *Isistius brasiliensis,* a squaloid shark, the probable cause of crater wounds on fishes and cetaceans. Fish. Bull., U.S. 69:791-798.
JORDAN, D. S., and B. W. EVERMANN.
1896. The fishes of North and Middle America. U.S. Natl. Mus. Bull. 47:1-1240.
1926. A review of the giant mackerel-like fishes, tunnies, spearfishes and swordfish. Calif. Acad. Sci., Occas. Pap. 12:1-113.
JUNE, F. C., and J. W. REINTJES.
1957. Survey of the ocean fisheries off Delaware Bay. U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 222, 55 p.
KRUMHOLZ, L. A.
1958. Relative weights of some viscera in the Atlantic marlins. Bull. Am. Mus. Nat. Hist 114:402-405.
KRUMHOLZ, L. A., and D. P. DE SYLVA.
1958. Some foods of marlins near Bimini, Bahamas. Bull. Am. Mus. Nat. Hist. 114:406-411.
LAMONTE, F. R.
1955. A review and revision of the marlins, genus *Makaira.* Bull. Am. Mus. Nat. Hist. 107:323-358.
1958a. Scales of the Atlantic species of *Makaira.* Bull. Am. Mus. Nat. Hist. 114:381-395.
1958b. Notes on the alimentary, excretory, and reproductive organs of Atlantic Makaira. Bull. Am. Mus. Nat. Hist. 114:396-401.
LAMONTE, F., and D. E. MARCY.
1941. Swordfish, sailfish, marlin and spearfish. Ichthyol. Contrib. Int. Game Fish Assoc. 1(2):1-24.
LEGENDRE, R.
1928. Présence du *Tetrapturus belone* au large de la Bretagne. Bull. Soc. Zool. Fr. 53:391-392.
LOWE, R. T.
1840. On new species of fishes from Madeira. Proc. Zool. Soc. Lond. Part VIII, p. 36-39.
LOZANO CABO, F.
1958. Los escómbridos de las aguas españolas y marroquíes y su pesca. Primera parte. Trab. Inst. Esp. Oceanogr. 25, 254 p.
MAÉDA, H.
1967. Distribution pattern of fish in relation to fishing method, with special reference to that of tuna along long-line. Proc. Symp. Scombroid Fishes, Part 3, p. 1025-1041. Mar. Biol. Assoc. India, Mandapam Camp.
MATHER, F. J., III.
1960. Recaptures of tuna, marlin and sailfish tagged i western North Atlantic. Copeia 1960:149-151.
MATHER, F. J., III, A. C. JONES, and G. L. BEARDSLEY
1972. Migration and distribution of white marlin and marlin in the Atlantic Ocean. Fish. Bull., U.S. 70:28
MATHER, F. J., III, J. M. MASON, JR., and H. L. CLAR
1974. Migration of white marlin and blue marlin in the ern North Atlantic Ocean—tagging results since May *In* R. S. Shomura and F. Williams (editors), Proceedi the International Billfish Symposium, Kailua-K Hawaii, 9-12 August 1972, Part 2. Review and Contri Papers, p. 211-225. U.S. Dep. Commer., NOAA Tech. NMFS SSRF-675.
MIGDALSKI, E. C.
1958. Angler's guide to the salt water game fishes—Atl and Pacific. Ronald Press, N.Y., 506 p.
MIHARA, T., and R. C. GRIFFITHS.
1971. La flota atunera Venezolana. Projecto Investig Desarrollo Pesquero MAC-PNUD-FAO. Inf. Tech. 26,
MIYAKE, M., and S. HAYASI.
1972. Field manual for statistics and sampling of Atl tunas and tuna-like fishes. Int. Comm. Conserv. Tunas, 79 p.
MIYAKE, M., and C. G. TIBBO.
1972. Statistical bulletin 2. Int. Comm. Conserv. Tunas.
MONOD, T.
1968. Le complex urophore des poissons téléostéens. M Inst. Fondam. d'Afr. Noire 81, 705 p.
MORRIS, B. F.
1971. Petroleum: Tar quantities floating in the northwes Atlantic taken with a new quantitative neuston Science (Wash., D.C.) 173:430-432.
MORROW, J. E.
1957. On the morphology of the pectoral girdle in the g *Makaira.* Bull. Bingham Oceanogr. Collect., Yale U 16:88-105.
1965. White marlin. *In* A. J. McClane (editor), McCl standard fishing encyclopedia and international an guide, p. 997-999. Holt, Rinehart, Winston, N.Y.
NAKAMURA, E. L.
1971. An analysis of the catches and biology of big g fishes caught by the New Orleans Big Game Fishing C 1966-1970. East Gulf Sport Fish. Mar. Lab. Rep., 38
NAKAMURA, E. L., and L. R. RIVAS.
1972. Big game fishing in the northeastern Gulf of Me during 1971. Natl. Mar. Fish. Serv., Panama City, 20 p.
NAKAMURA, H.
1938. Report of an investigation of the spearfishes Formosan waters. [In Jap.] Reports of the Tai Government-General Fishery Experiment Station, No. [Translated by W. G. Van Campen, 1955, U.S. Dep. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 153, 46
1949. The tunas and their fisheries. [In Jap.] Take Shobō, Tokyo. [Translated by W. G. Van Campen, U.S. Dep. Int., Fish Wildl. Serv., Spec. Sci. Rep. Fish 115 p.]
NAKAMURA, I., T. IWAI, and K. MATSUBARA.
1968. A review of the sailfish, spearfish, marlin and sword of the world. Kyoto Univ., Misaki Mar. Biol. Inst., S Rep. 4, 95 p.
NARDO, G. D.
1832. (*Skeponopodus*). Ber. ü. Verein. Dtsch. Natur Aerzte:99.

KOLAYEVA, V. M., and K. EZPELETA.
1966. On parasitic fauna of fishes in the Gulf of Mexico. [In Russ.] Naukova Dumka, Kiev, p. 194-204.
VCHINNIKOV, V. V.
1970. Mech-ryba i parusnikovye (Atlanticheskii okean. Ekologiya i funktsional'naya morfologiya) (Swordfishes and billfishes in the Atlantic Ocean. Ecology and functional morphology). Atl. Nauchno-Issled. Inst. Rybn. Khoz. Kaliningrad, 106 p. (Translated by H. Mills, Israel Program Sci. Transl., 1971, 77 p.; available U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA, as TT 71-50011.)
AIVA, M. P.
1972. Pesquerias de tunidos y especies afines de Brasil, en al año 1971. Int Comm. Conserv. Atl. Tunas, SCRS/72/57, 6 p.
ARIN, N. V.
1968. Ikhtiofauna okeanoskoi epipelagiali (Ichthyofauna of the epipelagic zone). Akad. Nauk SSSR, Tr. Inst. Okeanol., Moskva. (Translated by M. Raveh, 1970, Israel Program Sci. Transl., 206 p.)
OEY, F.
1861. Memorias sobre la historia natural de la Isla de Cuba. acompañadas de sumarios latinos y extractos en frances. Vol. 2. Viuda de Barcina, Havana, 442 p.
OSTEL, E.
1959. Liste commentée des poissons signalés dans l'Atlantique tropico-oriental nord, du Cap Spartel au Cap Roxo, suivie d'un bref apercu sur leur répartition bathymétrique et géographique. Bull. Soc. Sci. Bretagne 34(1+2) (Premiere partie), (3+4) (Deuxieme partie):129-170, 241-281.
OTTHOFF, T., and W. J. RICHARDS.
1970. Juvenile bluefin tuna, *Thunnus thynnus* (Linnaeus), and other scombrids taken by terns in the Dry Tortugas, Florida. Bull. Mar. Sci. 20:389-413.
ADCLIFFE, L.
1926. "Opah" and "Skilligallee" landed at Boston fish pier. Copeia 151;112.
AFINESQUE, C. S.
1810. Caratteri di alcuni nuovi generi e nuove specie di animali e piante della Sicilia, con varie osservazioni sopri i medesimi. Palermo, 105 p.
EINTJES, J. W., and C. M. ROITHMAYR.
1960. Survey of the ocean fisheries off Delaware Bay. Supplement Report, 1954-57. U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 347, 18 p.
VAS, L. R.
1954. A preliminary report on the spawning of the western North Atlantic bluefin tuna (*Thunnus thynnus*) in the Straits of Florida. Bull. Mar. Sci. Gulf Caribb. 4:302-322.
1956. Definitions and methods of measuring and counting in billfishes (Istiophoridae, Xiphiidae). Bull. Mar. Sci. Gulf. Caribb. 6:18-27.
OBINS, C. R.
1974. Summer concentration of white marlin, *Tetrapturus albidus*, west of the Straits of Gibraltar. *In* R. S. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972, Part 2. Review and Contributed Papers, p. 164-174. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-675.
OBINS, C. R., and D. P. DE SYLVA.
1960. Description and relationships of the longbill spearfish, *Tetrapturus belone*, based on the Western North Atlantic specimens. Bull. Mar. Sci. Gulf Caribb. 10:383-413.
1963. A new western Atlantic spearfish, *Tetrapturus pfluegeri*, with a redescription of the Mediterranean spearfish *Tetrapturus belone*. Bull. Mar. Sci. Gulf Caribb. 13:84-122.
RODRÍGUEZ-RODA, J., and J. K. HOWARD.
1962. Presence of Istiophoridae along the south Atlantic and Mediterranean coasts of Spain. Nature (Lond.) 196:495-496.
ROSA, H., JR.
1950. Scientific and common names applied to tunas, mackerels and spearfish of the world with notes on their geographic distribution. FAO (Food Agric. Organ. U.N.), Wash., D.C., 235 p.
1965. Preparation of synopses on the biology of species of living aquatic organisms. FAO (Food Agric. Organ. U.N.) Fish. Synop. (1), Rev. 1, 75 p.
SARÀ, R.
1968 La evoluzione della pesca del tonno nel basso Tirreno, negli ultimi anni. Boll. Pesca Piscic. Idrobiol., New Ser. 23:33-46.
SASSI, A.
1846. De pesci del mare di Genova. Nuovi Ann. Sci. Nat. Rend. Sess. Soc. Agr. Accad. Sci. Ist Bologna, Ser. 2, 6:386-393.
SCOTTON, L. N., and D. P. DE SYLVA.
1972. Fish babies. Sea Front. 18:194-201.
SHIOHAMA, T., M. MYOJIN, and H. SAKAMOTO.
1965. The catch statistic data for the Japanese tuna long-line fishery in the Atlantic Ocean and some simple considerations on it. Rep. Nankai Reg. Fish. Res. Lab. 21, 131 p.
SILAS, E. G.
1967 Parasites of scombroid fishes. Part I. Monogenetic trematodes, digentic trematodes, and cestodes. Proc. Symp. Scombroid Fishes, Part 3, p. 799-875. Mar. Biol. Assoc. India, Mandapam Camp.
SILAS, E. G., and A. N. P. UMMERKUTTY.
1967. Parasites of scombroid fishes. Part II. Parasitic copepoda. Proc. Symp. Scombroid Fishes, Part 3, p. 876-993. Mar. Biol. Assoc. India, Mandapam Camp.
SMITH, J. L. B.
1956. Swordfish, marlin and sailfish in South and East Africa. Rhodes Univ., Dep. Ichthyol., Ichthyol Bull. 2:25-33.
SQUIRE, J. L., JR.
1962. Marlin and swordfish in oceanic waters of the western North Atlantic. Copeia 1962:216-219.
STEPHENS, W. M.
1965. Summer cruise to the Sargasso Sea. Sea Front. 11:108-123.
TALBOT, F. H., and M. J. PENRITH.
1962. Tunnies and marlins of South Africa. Nature (Lond.) 193:558-559.
1963. The white marlin *Makaira albida* (Poey) from the seas around South Africa. S. Afr. J. Sci. 59:330-331.
TORTONESE, E.
1940. Sugli scomberoidei mediterranei del genere *Tetrapturus*. Boll. Mus. Zool. Anat. Comp. Torino 48, Ser. 3(115):173-178.
1961. Mediterranean fishes of the family Istiophoridae. Nature (Lond.) 192:80.
1962. Osservazioni comparative intorno alla ittiofauna del Mediterraneo e dell'Atlantico occidentale (Florida e Isole Bahamas). Natura (Milano) 53:1-20.
1970. Comparsa di *Tetrapturus albidus* Poey (Pesci S gombroidi) nel Golfo di Genova. Boll. Pesca Piscic. Idrobiol., New Ser. 25(1):81-83.
UEYANAGI, S.
1959. Larvae of the striped marlin, *Makaira mitsukurii* (Jordan et Snyder). [In Jap., Engl. summ.] Rep. Nankai Reg. Fish. Res. Lab. 11:130-146. (Translated by W. G. Van

Campen, 1961, 20 p.; available at Southwest Fisheries Center, National Marine Fisheries Service, NOAA, Honolulu, Hawaii.)

UEYANAGI, S., S. KIKAWA, M. UTO, and Y. NISHIKAWA.
1970. Distribution, spawning, and relative abundance of billfishes in the Atlantic Ocean. [In Jap., Engl. synop.] Bull. Far Seas Fish. Res. Lab. (Shimizu) 3:15-55.

UEYANAGI, S., and H. WATANABE.
1965. On certain specific differences in the vertebrae of the istiophorids. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 22:1-8.

VÉRANY, J. B.
1847. Aggiunto al catalago de' pesci della Liguria. Atti Ottava Riunione Sci. Ital. Genoa 1846:492-494.

VOSS, G. L.
1953. A contribution to the life history and biology of the sailfish, *Istiophorus americanus* Cuv. and Val., in Florida waters. Bull. Mar. Sci. Gulf Caribb. 3:206-240.

WALLACE, D. H., and E. M. WALLACE.
1942. Observations on the feeding habits of the white marlin, *Tetrapturus albidus* Poey. Dep. Res. Educ., Board Nat. Resour., Publ. Chesapeake Biol. Lab. 50:3-10.

WATANABE, H.
1961. Study of tuna hooked layer by longline. [In Ja
Kanagawa Prefect. Fish. Exp. Stn. Rep. Ser. 4, 10 p.

WISE, J. P.
1968. The Japanese Atlantic longline fishery, 1964, and
status of the yellowfin tuna stocks. U.S. Fish Wildl. Se
Spec. Sci. Rep. Fish. 568, 5 p.

WISE, J. P., and C. W. DAVIS.
1973. Seasonal distribution of tunas and billfishes in
Atlantic. U.S. Dep. Commer., NOAA Tech. Rep. NM
SSRF 662, 24 p.

WISE, J. P., and W. W. FOX, JR.
1969. The Japanese Atlantic longline fishery, 1965, and
status of the yellowfin tuna and albacore stocks. U.S. F
Wildl. Serv., Spec. Sci. Rep. Fish. 582, 7 p.

WISE, J. P., and J. C. LE GUEN.
1969. The Japanese Atlantic long-line fishery, 1956-1
In Proc. Symp. Oceanogr. Fish. Resour., Trop. Atl.-F
Pap. Contrib. 26:317-347.

YOSHIDA, H. O.
1966. Tuna fishing vessels, gear, and techniques in the Pac
Ocean. *In* T. A. Manar (editor), Proc. Gov. Conf. Ce
Pac. Fish. Resour., p. 67-89. State of Hawaii.

Synopsis of the Biology of the Sailfish, *Istiophorus platypterus* (Shaw and Nodder, 1791)[1]

G. L. BEARDSLEY, JR.,[2] N. R. MERRETT,[3] and W. J. RICHARDS[2]

IDENTITY

1.1 Nomenclature

1.11 Valid name

Istiophorus platypterus (Shaw and Nodder, 1791) (Fig. 1).



Figure 1.—Sailfish, *Istiophorus platypterus* (Shaw and Nodder, 1791).

Originally described as *Xiphias platypterus* Shaw and Nodder, 1791. Type locality, Indian Ocean. Holotype, British Museum of Natural History (BMNH) Catalog No. 1964.7.2.1 (G. Shaw and F. P. Nodder. 1791. *Xiphias platypterus*. The broad-finned swordfish. Natural Misc. No. 28. pl. 88).

1.12 Objective synonymy (modified after Morrow and Harbo, 1969).

Xiphias platypterus Shaw and Nodder, 1791 (see above).

Scomber gladius Block, 1793. Type locality, East and West Indies. Type specimen, none.

Xiphias velifer Block and Schneider, 1801. Type locality, India. Replacement name for *S. gladius* Block, 1793.

Istiophorus gladifer Lacépède, 1802. Type locality, Indian Ocean. Type specimen, none (based on Broussonette's "Le Voilier").

Histiophorus indicus Cuvier, *in* Cuvier and Valenciennes, 1831. Type locality, Indian Ocean. Holotype, BMNH Catalog No. 1964.7.2.1.

Histiophorus americanus Cuvier, *in* Cuvier and Valenciennes, 1831. Type locality, Atlantic Ocean off Brazil. Type specimen, none (based on Marcgrave's description and figure).

Histiophorus pulchellus Cuvier, *in* Cuvier and Valenciennes, 1831. Type locality, South Atlantic. Holotype, a juvenile, current location unknown.

Histiophorus gracilirostris Cuvier, *in* Cuvier and Valenciennes, 1831. Type locality, Mauritius (?). Holotype, Masée National d'Histoire Natural, Paris (NMHN) Catalog No. A9462.

Histiophorus ancipitirostris Cuvier, *in* Cuvier and Valenciennes, 1831. Type locality, Red Sea. Holotype, NMHN A9463.

Makaira velifera Cuvier, 1832. Type locality, Haiti. Holotype, NMHN Catalog No. B1648 (?).

Histiophorus orientalis Temminck and Schlegel, 1844. Type locality, Japan. Type specimen, none extant, based on a fresh specimen.

Histiophorus immaculatus Rüppel, 1835. Type locality, Red Sea near Djedda. Holotype, an 18-inch juvenile, whereabouts unknown.

---

[1]Contribution No. 234, Southeast Fisheries Center, National Marine Fisheries Service, NOAA, Miami, Fl 33149.

[2]Southeast Fisheries Center, National Marine Fisheries Service, NOAA, Miami, FL 33149.
[3]Institute of Oceanographic Sciences, Surrey, England.

Histiophorus granulifer Castelnau, 1861. Type locality, Cape of Good Hope. Holotype, a mutilated skeleton, said to be in the Capetown Museum.

Istiophorus greyi Jordon and Evermann, 1926. Type locality, Cape San Lucas, lower California. Type specimen, California Academy of Sciences (CAS) Catalog No. 605, a photograph.

Istiophorus wrighti Jordan and Evermann, 1926. Type locality, off Miami, Fla. Type specimen, CAS Catalog No. 603, a photograph.

Istiophorus maguirei Jordan and Evermann, 1926. Type locality, off Long Key, Fla. Type specimen, CAS Catalog No. 604, a photograph.

Istiophorus volador Jordan and Evermann, 1926. Type locality, off Long Key, Fla. Type specimen, CAS Catalog No. 606, a photograph.

Istiophorus eriquius Jordan and Evermann, 1926.[4] Type locality, Hawaii. Holotype, Bernice P. Bishop Museum Catalog No. 3424.

Istiophorus brookei Fowler, 1933. Type locality, Tahiti. Holotype, Academy of Natural Sciences, Philadelphia Catalog No. 55923.

For further lists see Nakamura, Iwai, and Matsubara (1968).

Several authors have recently considered the senior synonym of the Atlantic form of the sailfish to be *Istiophorus albicans* (Latreille, 1804). Whitley (1939) first called attention to a series of volumes entitled "Nouveau Dictionaire d'Histoire Naturelle" published in Paris from 1802 to 1804. The fish sections in these volumes were by L. A. G. Bosc and are clearly designated with the letter (B) at the end of each fish account. Bosc (1803) in volume 13 described the Atlantic sailfish based on Marcgrave's description and figure. Bosc applied the common name 'le makaira blanchâtre' to this species. Latreille (1804) in volume 24, following his description of the genus *Makaira*, clearly applied a scientific name to Makaira blanchâtre, in the following manner: "Makaira blanchâtre, *makaira albicans* Bosc." Latreille wrote nothing else, thus under the International Rules of Zoological Nomenclature this name must be considered a nomen nudum. Had Latreille referred to Bosc's description in the earlier volume the name would be available.

1.2 Taxonomy

1.21 Affinities

Suprageneric

Phylum Chordata
Subphylum Vertebrata
Superclass Gnathostomata
Class Osteichthyes
Subclass Actinopterygii
Order Perciformes
Suborder Xiphioidei
Family Istiophoridae

Gosline (1968) in his review of perciform subord presented evidence that the relation of billfishes scombrids and their allies may be one of convergen He placed the family Istiophoridae along with the f ily Xiphiidae and provisionally, the Luvaridae, i separate suborder Xiphiodei.

Generic

Monotypic genus, see specific diagnosis.

Istiophorus Lacépède, 1802. (ref.). Type spe designated as *Istiophorus platypterus* (Shaw Nodder, 1791). Holotype, BMNH 1964.7. Designated by Morrow and Harbo (1969).

Specific

Diagnosis, first dorsal fin very high, sail shaped, middle rays decidedly the longest; the ventral rays very long and fin membrane well develop

1.22 Taxonomic status

Two recent reviews have treated the sailfish ge *Istiophorus*. Morrow and Harbo (1969) and Nakam et al. (1968). The latter recognized two species sailfishes—*I. platypterus* (Pacific sailfish, Basho jiki) from the Indo-Pacific and *I. albicans* (Atla sailfish, Nishibashokajiki) from the Atlantic. T based their separation on greater length of the p toral fin and the caudal fin on Atlantic specimens than 90 cm in length. Above 90 cm in length they indistinguishable. Morrow and Harbo disputed t contention in a footnote on p. 39 of their paper say that their data (though not presented) do not supp such a distinction. In this synopsis we follow Mor and Harbo, but emphasize that studies are needed determine exactly the specific status of the ge *Istiophorus*.

1.23 Subspecies

Various authors have recognized subspecies. Ba on Morrow and Harbo (1969) this does not seem to justified.

1.24 Standard common names and vernacu names.

The names capitalized are official or in more co mon use. Compiled mostly from FAO (1950).

Arabia — FÁRAS
Brazil — AGULHÃO DE VELA, Agulhão bandei Agulhão, Bicudo, Guebuçú.

[4]Jordan and Evermann (1926) attribute *I. eriquius* to Jordan and Ball, but under present interpretation of the International Code of Zoological Nomenclature, Jordan and Evermann must be given the responsibility for the name.

ritish West Indies — SAILFISH, Billfish, Ocean gar, squadron, Mère, Balahoo, Mammon balahoo (Patois names).
eylon — SAILFISH, Thalapatha (Sinhalese dialect), Myl min (Tamil dialect).
hina — TONG FANG CHIYII
omores Is. — MBASSI KOURI
uba — AGUJA VOLADORA, Aguja de abanico, Abanico, Prieta, Voladora, Bicuda.
rance — LE VOILIER
hana — AMERICAN SAILFISH, Onyankle (Ga), Adzietekwesi (Adanme), Fetiso (Fante).
idia — SAILFISH, Yemungolah (Tamil dialect), Plain sailfish, Tadmása (Marathi dialect).
apan — BASHOKAJIKI, Nishibashokajiki (Atlantic Ocean form).
enya — NSULI NSULI, Mbassi
Iadagascar — NDWARO
Ialaya — LAYER, Mersuji
Iexico — PEX VELA, Volador
hilippines — SAILFISH, Sailfin, Malasugi (Bikol), Kandayan, Kandelan (Marinao, Samal and Tao Sug), Dogso, Dugso, Malasugi, Liplipan (Visayan), Dumosok (Pangasinan).
ortugal — VELEIRO
enegal — ESPADON, Oumbajhe
umatra — IKAN-JEGAN, Johoo malags
aiwan — YU SAN YU, yu sanchiyu
anzania — NSULI NSULI, Mbassi
SSR — PARUSNIK-RYBA
nion of South Africa — SAILFISH
nited States — SAILFISH, Atlantic sailfish, Pacific sailfish, Mexican volador, Pacific volador, Florida sailfish, Spikefish, Boohoo, Woohoo, Voilier, Volador, Au-lepe (Hawaiian).
enezuela — AGUJA VELA
ietnam — CÁ CO

1.3 Morphology (condensed from Nakamura et al., 1968; Morrow and Harbo, 1969).

1.31 External morphology

The dorsal fin is extremely high, sail shaped with the middle rays decidedly the longest. The ventral fin rays are very long, almost reaching the vent, and the fin membrane is well developed. On the sides of the body there are 10 rows of striped crest patterns consisting of many light blue dots. There is a clear crest on the outer edge of the head between the preocular area and the base of the first dorsal fin. The body is extremely flat; the cranium long and narrow. The neural spines and hemal spines of the central vertebrae form a triangular shape. There are 24 vertebrae (12 + 12 = 24). The lateral apophysis is not well developed.

There are two dorsal fins and two anal fins (counts are given in Table 1). The pectoral fin has 17-20 soft rays; the ventral fin has 1 spine and 2 soft rays. The caudal fin has 17 principal rays (9/8) and 20-22 secondary rays (10/10 or 11/11).

The body is elongated and laterally compressed. The snout is long and round in cross section. Many of the scales have a single point which is not sharp. The lateral line is easily distinguishable, curving above the pectoral fin and then continuing in a straight line toward the tail. The caudal fin is deeply forked and two scutes precede it on each side of the caudal peduncle. The pectoral fin is low on the body, relatively long and pointed. The second dorsal and second anal fins are small. Internally, the nasal rosette is radially shaped with about 48 nasal laminae. The abdominal cavity is extremely long, extending past the vent to the middle of the base of the second anal fin. The visceral organs are also elongate.

Other than the differences noted above between the

Table 1.—Number of dorsal and anal rays. (Modified from Morrow and Harbo, 1969; Merrett, 1971.)

	Number of dorsal fin rays																
	First dorsal fin														Second dorsal fin		
Region	37	38	39	40	41	42	43	44	45	46	47	48	49	$\bar{x}$	6	7	8
Atlantic	1	2	-	3	2	12	13	29	21	5	1	1	-	43.66	25	55	3
Peru									2	3	4			46.22	-	17	-
Pacific							1	-	5	4	7	5	2	46.63	7	17	-
Indian								1	3	2	7	5	-	46.67	1	18	-
Africa			3	-	3	15	16	9	10	3	4	-	3	43.59	7	34	2

	Number of anal fin rays													
	First anal fin										Second anal fin			
	8	9	10	11	12	13	14	15	16	$\bar{x}$	5	6	7	8
Atlantic	1	1	3	3	14	39	19	-	1	12.80	-	39	44	-
Peru				5	1	2	1			11.89		1	8	
Pacific					1	8	12	3		13.71		6	18	
Indian					2	10	3	1		13.19		4	13	
Africa	4	2	6	22	14	12	7			11.55	2	11	23	2

Atlantic and Pacific populations (see 1.22) there are some indications of possible differentiation of certain populations, but according to Morrow and Harbo (1969), "there is little consistency in the direction of these indications."

Morrow and Harbo show some differentiation with age, and Vick (1963) examined allometric and isometric growth of adult *I. platypterus* in some detail, but extreme differences are noted in the young and will be covered in 3.2.

1.32 Cytomorphology

Nothing found in the literature.

1.33 Protein specificity

Nothing found in the literature.

2 DISTRIBUTION

2.1 Total area

Sailfish are circumtropical, occurring in all wa waters of the world. In the Atlantic Ocean sail range from lat. 30°S to 30°N on the western side w only occasional stragglers beyond these limits. W (1953) stated that individuals taken as far north the Gulf of Maine are apparently summer stragg and their distribution is extended during long, summers with prevailing southerly winds. In eastern Atlantic distribution is generally m restricted, from lat. 10°S to 20°N (Fig. 2). It may inferred that the densest concentrations also occu the western Atlantic close to land masses accordin data from the Japanese longline fishery with the ta realization that Japanese commercial vessels fai









Figure 2.—Distribution of catches of sailfish and spearfish (per 1,000 hooks) in the four quarters of the year, 1956-68 (from Wise and Davis, 1973).



Figure 3.—Distribution of the sailfish, *Istiophorus gladius*, in the Pacific Ocean (from Howard and Ueyanagi, 1965).

differentiate between sailfish and spearfish. Ueyanagi et al. (1970) indicate, however, that concentrations of Atlantic sailfish are greatest on the western side. Large concentrations of sailfish in the Gulf of Mexico, along the southeast coast of Florida, and throughout the Carribbean are heavily fished by sports fishermen.

In the eastern Atlantic abundance is apparently low based on longline catch records. This may be misleading since there are no established sport fishing centers in the eastern Atlantic and longline operations are often not close enough to shore to detect sailfish concentrations. However, a localized area of high concentration of sailfish has been noted by Ovchinnikov (1970) off West Africa (the Freetown-Conakry region). He indicated that a similar accumulation of sailfish also occurred in the western Atlantic in the region of eastern and northeastern Brazil.

In the Pacific, sailfish distribution is also more extensive in the western half than in the eastern (Fig. 3). Longline catch data show sailfish catches as far south as 27°S and as far north as lat. 40°N. Koto, Furukawa, and Kodama (1959) reported heaviest concentrations near New Guinea, the Caroline Islands and Solomon Islands, and in the Banda Sea, Timor Sea, and East China Sea. Howard and Ueyanagi (1965) suggested that there is a close relationship between the distribution of sailfish in the western Pacific and the Kuroshio Current since both adults and juveniles are found in the coastal waters of Japan in the Kuroshio Current. In the eastern Pacific, longline catch records show a rather restricted distribution from lat. 5°S to 25°N (Kume and Joseph, 1969a). High catch rates near the coast of Mexico indicate areas of high abundance. Sailfish are caught by sport anglers near shore from the Gulf of California to northern Peru.

There is little information on sailfish distribution in the Indian Ocean. Williams (1964) indicated that sailfish are distributed throughout the East African coastal area from lat. 1°30′N to 10°30′S with localized concentrations. Japanese longliners catch small numbers of sailfish throughout most of the Indian Ocean to at least lat. 29°S in the western Indian Ocean.

Sailfish have been reported from the Mediterranean; however, most of these reports were based on larval or juvenile specimens and in at least one case (Ben-Tuvia, 1966) have been subsequently identified as the Mediterranean spearfish, *Tetrapturus belone*. It is doubtful if sailfish occur in the Mediterranean.

2.2 Differential Distribution

2.21 Spawn, larvae, and juveniles

Gehringer (1956) reported on sailfish larvae captured in the western Atlantic and Gulf of Mexico from March through October. He concluded that d development of Atlantic specimens takes plac warm Gulf Stream waters. In a later p (Gehringer, 1970) he examined larvae from eastern Atlantic captured in April.

Ueyanagi (1959) found that sailfish larvae are c ly associated with the Kuroshio Current in western Pacific. In the eastern Pacific sailfish la and juveniles have been captured from Septe through April (Howard and Ueyanagi, 1965).

Jones (1959) reported on 16 sailfish larvae capt in February from the Laccadive Sea.

See also 3.16.

2.22 Adults

Sailfish are year-round residents over most of t range, however pronounced seasonal variation abundance and distribution are evident in most a Wise and Davis (1973) showed that sailfish are ca off Brazil by longliners all year long, but catches larger and over a broader area from October thr March than from April through September. Gre abundance in the Caribbean Sea and Gulf of Me is from April through September; few are caug the winter months. Voss (1953) stated that sailfis found off the lower Florida coast throughout the in considerable numbers, but during the sum months they are less concentrated than in wi Cadenat (1961) has shown that sailfish in the eas Atlantic, off the Ivory Coast, are present inshor greatest abundance during the winter months, practically disappear from the coastal waters du August to November.

Howard and Ueyanagi (1965) reported that in eastern Pacific sailfish are present all year seasonal changes in density are marked. Kume Joseph (1969a) stated that sailfish appear to be ab dant off central Mexico all year with appa seasonal movements. In the Gulf of Panama sai are most abundant from April-May thro November-December with a peak from June September. Near Acapulco, sailfish appear to be n abundant in winter. From the Gulf of Califo northward the period of greatest abundance is f May to October.

Nakamura (1949) stated that sailfish are n abundant around June in the Kuroshio Current of the Philippine Islands and Formosa, and Koto e (1959) indicated that sailfish are most abundan the East China Sea from November through A with low densities from May through August.

There is little information on seasonal distribut in the Indian Ocean; however, at Malindi, Ke Williams (1970) reported that sport catches restricted to the period October to March with p catches in December and January. That sailfish s a peak in abundance during the northeast monsoo

is and the broader area of the equatorial western Indian Ocean was confirmed by Merrett (1971).

2.3 Determinants of Distribution Changes

Voss (1953) stated that sailfish distributions along he east coast of Florida appear to be affected by wind nd temperature. In the summer there is a "diffusion" f sailfish to the northward correlated with a orthward extension of warm water. These same fish re driven southward to congregate in schools off the lorida coast with the beginning of cold weather and ortherly winds.

Evidence presented by Ovchinnikov (1966) indicates that the frontal zone of the Canaries Current nd the Equatorial Countercurrent is responsible for he aggregation of sailfish off the coast of West Africa. Ie stated that in spring sailfish move along the coast om south to north, and apparently move back again om north to south in the autumn, following the 28°C otherm. Similarly, Cadenat (1961) indicated that he period of increased abundance off the Ivory Coast oincides with the periods of maximum surface emperature, around 28°C.

In the western Pacific the distribution of postlarvae nd adult sailfish appears to be closely related to the uroshio Current (Nakamura, 1949; Ueyanagi, 1959). Iakamura also stated that the season of dense disribution in this area coincides with the spawning eason. Koto et al. (1959) found that sailfish over 160 m long migrate southward out of the East China Sea, resumably for spawning.

In the eastern Pacific, Kume and Joseph (1969a) ound that seasonal north-south movements of sailfish ff the coast of Mexico appear to coincide with the easonal movements of the 28°C isotherm.

Williams (1964, 1970) concluded that the abunance and distribution of sailfish in the Indian Ocean f East Africa is positively correlated with the onths of the northeast monsoons when the East frican Coastal Current reaches its maximum mperature (29°-30°C) and minimum salinity (35.2-5.3‰). This is also the time of highest biological roductivity in surface waters caused by a mixing due the junction of the southward flowing Somali urrent and the northward flowing East African oastal Current. The inshore penetration of nutrient ch Somali Current water fluctuates seasonally. Villiams (1970) summarized by stating that: "The nown fluctuations in overall productivity and the easonal abundance of shoals of pelagic fish in the orth Kenyan area during the northeast monsoon ight well be due to the fluctuations in the penetraon of the SC tongue."

2.4 Hybridization

Nothing found in the literature.

3 BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

Sailfish are heterosexual. There are no external characters known to distinguish males from females, although Jolley (1974) and Nakamura and Rivas (1972) indicated that large Atlantic sailfish were usually females.

3.12 Maturity

There is little information available on size at first maturity. Merrett (1971) examined gonads from 65 female sailfish from the equatorial western Indian Ocean but found only one immature specimen 130 cm long (measured from center of orbit to shortest caudal ray) and weighing 13.6 kg. According to de Sylva (1957) sailfish this size in the western Atlantic are approximately 18 mo old.

3.13 Mating

Mating in large pelagic fishes has often been assumed to be promiscuous with sexual products released indiscriminately and fertilization occurring by chance. Nakamura (1949) stated, however, that "Spawning (in sailfish) is carried on with a male and a female paired or two or three males chasing a single female, and this act can be seen often during the spawning season." Voss (1953) reported observing similar behavior in sailfish off the Florida coast. Sailfish were often sighted in groups of two or three swimming very slowly in shallow water. The ovaries of females captured in this area were all extremely large and very ripe. Voss also reported that another observer sighted two sailfish pressed tightly together and swimming slowly in shallow water. One was harpooned and was identified as a female containing ripe overies weighing about 3.2 kg.

3.14 Fertilization

Fertilization is probably external.

3.15 Gonads

According to Voss (1953) the number of eggs spawned by a female sailfish is large based on western Atlantic specimens. He determined that a 1.4-kg ovary contained 2.3 million eggs and obtained counts as high as 4.7 million eggs in later analyses. Voss also stated that gonads from female sailfish may be as long as 50.8 cm and weigh up to 3.6 kg. Jolley (1974) showed that ripe gonads of Atlantic sailfish weighed 8.1-12.5% (mean 9.9%) of total body weight. Fecundity estimates ranged from 0.75 to 1.56 million ova. Only the most advanced ova, 1.2- to 1.4-mm diameter, were counted. Ovchinnikov (1970) gave fecundity estimates of 1.6-11.5 million eggs from ovaries of 1.1-2.1 kg from Atlan-

tic sailfish. Merrett (1971) found as many as 19.5 million eggs in sailfish from East African waters and pointed out that fecundity increased sharply with fish size.

A definite relation between relative ovary weight (ovary weight × 10^3/fish weight) in *I. platypterus* and the state of gonad development has been established (Merrett, 1971). Having demonstrated that egg diameter is a reliable means of determining the state of maturity of an overy, he compared the relative ovary weight with this criterion. It was found that while this character did not provide staging evidence throughout the oogenic cycle, it could provide for accurate diagnosis during the growth phase. Comparable application of relative testis weight was found to be less meaningful due to the lack of seasonal variation in testis weight of the sailfish examined. From a total of 111 male sailfish examined over a period of 3½ yr, none was found to be in a running ripe condition, although spermatozoa were present in all testes examined due to differential maturation of the lobules within the gonad. It was suggested that the viability of the male was maintained at all times, owing to the possession of a muscular seminal vesicle in which spermatozoa are stored. The vesicle is thought to be capable of active expulsion of milt and subsequent passive replenishment, at all times, by suction.

3.16 Spawning

In the Atlantic, Voss (1953) summarizing early accounts, stated that, "From the literature it can be seen that it has previously been considered, mostly through capture of gravid females, that the sailfish breed near or off the Florida coast and the consensus of opinion gives the time as the early summer." Voss also reported that larvae from 3.9 to 8.0 mm standard length were captured on the east side of the Gulf Stream from 29 May to 2 July. Voss believed that sailfish move into shallow inshore waters to breed. Gehringer (1956) stated that, ". . . spawning appears to extend from April to September from south of Cuba north to Carolina waters, and beyond the 100 fathom line." Gehringer also indicated that there was a northward shift of size occurrences of larvae as the season progressed, indicating a corresponding northward shift in general spawning season. De Sylva (1957) indicated an extended spawning season in the western Atlantic and that very young larvae were captured from late April until mid-August. He also cited a personal communication with Al Pflueger who stated that he had found ripe females off southern Florida as late as October. Jolley (1974) found ripening females prominent among Atlantic sailfish examined during May through September.

Spawning in the eastern Atlantic has been observed all year long (Ovchinnokov, 1970), with peak intensity during the summer months on the West African shelf in the Conakry-St. Louis region. Ovchinnikov (1970) stated that sailfish in the Conakry-Freetown region become mature in February, March, and April, ar the end of April the fish spawn intensively. Ueya et al. (1970) reported taking sailfish larvae off An and Sierra Leone in the period November to A Gehringer (1970) reported on 34 specimens from to 238.0-mm standard length captured in the Gu Guinea in April, and Cadenat (1961) reported two vae from the same area in February and April, 35 44 mm total length.

In the Gulf of Mexico, Baughman (1941a) st that several ripe females were taken late in the se (probably late August) off the Texas coast, an males in all stages of maturity were captured in ly August. Based on the occurrence of larvae an surface current patterns in the Gulf of Me Gehringer (1956) concluded that two separate spa ing areas existed in the Gulf, one in the southeas Gulf from April through August and the other in western Gulf from June through August.

In the Pacific Ocean, sailfish apparently sp throughout the year in warm tropical wa (Nakamura, 1932, 1940; Yabe, 1953; Ueyanagi, 19 Based on the distribution and occurrence of lar Ueyanagi (1964) concluded that spawning took p in inshore waters near islands. The capture of adults from the western Pacific has been reporte Nakamura (1949) in May, July, August, December and by Ueyanagi (1964) in May, J November, and December. Ueyanagi stated although data were insufficient it appeared that spawning season in the northwest Pacific and South Pacific is in their respective summers.

Laurs and Nishimoto (1970) reported on juvenile sailfish captured in September in the eas Pacific off Central America. Surface temperature 29.1°C, and high standing stocks of zooplankton measured in the vicinity. Howard and Ueya (1965) stated that W. L. Klawe reported to them capture of juvenile sailfish in the equatorial wate the eastern Pacific in the months of February, A May, and December. Beebe (1941) reported the ture of two juveniles off the coast of Mexico, on March and the other in November, and Howard Ueyanagi reported on 11 juvenile sailfish taken in Gulf of Panama in September. Based on these counts, the spawning season in the eastern Pacif lengthy, and probably some spawning occurs all long.

Jones and Kumaran (1964) concluded from the tribution of larvae in the Indian Ocean that sailfis the eastern Indian Ocean spawn west of Sumatr September and October, and in the western In Ocean they spawn near Madagascar and Seychelles Islands in December and January. Mo (1964) examined sailfish at Taiwan and found run roe and milt in July. He stated that sailfish exam at Ceylon and the Maldive Islands in August September, the Seychelles in October and Novem and off the East African coast in January and M

50 and November 1957-January 1958 were not in a eeding condition. He concluded that most iophorids apparently spawn in the summer months, ughly May through September.

Williams (1963) stated that sailfish move into astal waters of Kenya in December, January, and bruary for spawning based on the presence of ma- re and spent adults in the area. Merrett (1971), wever, suggested that the abundance of sailfish in shore waters off Kenya was not for spawning but as a postspawning, feeding migration and reported pturing two running-ripe sailfish on longline gear proximately 111 km from the coast, one in January d the other in April. Williams (1970) revised his rlier spawning hypothesis and concluded that a mbination of the two hypotheses was most likely rrect. Jones (1959) indicated an additional spawn- g locality in the Laccadive Sea.

3.17 Spawn

A redescription of the egg of *I. platypterus* is given Nakamura (1949) who wrote: "The ripe ovarian gs are about 0.85 mm. in diameter and they have a ngle oil globule. Around the oil globule there is a le yellow indefinite nimbus. There are no structures the vitelline membrane and the egg as a whole is most colorless and clear." Merrett (1970) described genesis and spermatogenesis in this and other lfish species. He measured eggs shed from a cap- red sailfish and reported that the mean diameter as 1.304 mm. He also examined the morphology of e spermatozoa of this species and *Tetrapterus dax* and *Makaira indica*. Measurements showed the ads to be 1.3-1.9 μm in diameter; in length, the mid- e pieces measured 0.8-1.3 μm, and the tails up to μm. The only difference observed between the ecies was that the heads of the spermatozoa in lfish tended to be in the upper part of the diameter nge, and the middle pieces were found generally to shorter than in the other two species.

3.2 Preadult phase

3.21 Embryonic phase

(Defined as from fertilization to hatching, i.e., dur- g incubation period.) Nothing found in the erature.

3.22 Larvae phase

See 3.23.

3.23 Adolescent phase

Since young of these fishes gradually transform m larvae to adults, both the larval phase and olescent phase are discussed here in one unit.

Young billfishes (Istiophoridae) are all very similar their development, and consequently identification of the young stages is difficult. The first description of a young sailfish was by Cuvier (*in* Cuvier and Valenciennes, 1831). This specimen was 108 mm long and is the type specimen of *Histiophorus pulchellus* (see 1.12). Rüppell (1835) reported on a 45.7-cm juvenile from the Red Sea which he described as *H. immaculatus*. Günther (1873-74) published figures of young billfish 9, 14, and 60 mm long, and these were republished in one of his later works (1880). Lütken (1880) also republished these figures as well as a 5.5-mm specimen of his own. It is difficult to say positively that the specimens of Günther and Lütken are in fact sailfish. Goode (1883) also republished these figures and included an English translation of Lütken (1880). LaMonte and Marcy (1941) described a 14-mm specimen, and Beebe (1941) described two 12-cm specimens. Beebe also reviewed briefly the early work mentioned above and republished all of the figures mentioned. However, the fine detail presented in the earlier figures was not reproduced by Beebe and for detailed study we recommend that the original work be consulted. Descriptions of occasional specimens thought to be sailfish have appeared prior to papers describing complete series. Such descriptions are by Nakamura (1932, 1940, 1942, 1949) and Deraniyagala (1952).

Two studies describing complete series of larval through juvenile stages of sailfish were published in 1953, one by Voss based on Atlantic specimens and the other by Yabe based on Pacific specimens. Following these two papers, several papers also described sailfish based on complete series or else give important data on young forms. These studies are by Gehringer (1956, 1970), Jones (1959), Sun (1960), Ueyanagi and Watanabe (1962, 1964), de Sylva (1963), Ueyanagi (1963b) Mito (1966, 1967), and Strasburg (1970). We have reproduced here the figures of larval sailfish from Gehringer 1956 and 1970 (Fig. 4) that best delineate changes which take place during development. The most striking features are the early development of prominent pterotic and preopercular spines, the beaklike snout, and the very dark pigmentation which is acquired in early stages. At lengths greater than 10 mm the large dorsal fin begins to develop and with the long bill make this a most striking juvenile.

When seen in life between 4 and 15 mm in length the dark area is dark metallic blue (brown when preserved); the belly silvery (white when preserved); the head area, particularly the opercles, are flecked with a brilliant gold (this is lost soon after preservation); and the dorsal fin is a deep velvety black. The juvenile is colored very similar to the adult.

Ueyanagi (1963a, 1964) gave methods for distinguishing sailfish young from the other billfish species in the Indo-Pacific. His methods (1964 paper) are as follows:

> It is not easy to identify the larvae of different istiophorid species, because of their close resemblance with each other and



Figure 4.—Young sailfish: (a) 3.6 mm standard length (b) 4.7 mm SL; (c) 5.6 mm SL; (d) 8.1 mm SL; (e) 9.5 m (f) 11.3 mm SL; (g) 18.2 mm SL; (h) 20.9 mm SL; (i) 27



) 37.1 mm SL; (k) 55.1 mm SL; (l) 98.9 mm SL; (m) 155
L; (n) 216 mm SL (from Gehringer, 1956, 1970).

of marked differences from their respective adults, generally speaking, in their morphological characteristics. This is particularly true with those of very early stage before the snout develops its specific characteristics. However, the specific separation of the larvae is possible throughout their entire range mainly on the basis of their head profile.

Following are the criteria for identification:

(1) *Larvae under 5 mm in length*: The characters, as shown in Table A [Table 2], can be used for specific separation, although snout length does not provide a useful clue.

(2) *Larvae between 5 and 10 mm in length*: Besides the criteria given in Table A [Table 2], snout length and size of eyes can be used. Black marlin larvae are recognized by their snout. . . . The ratio of snout length to diameter of orbit is largest in sailfish, smallest in black marlin, and is between in shortnosed spearfish. More precisely, the ratio tends to be >1 in sailfish, <1 in black marlin, and $=1$ in shortnosed spearfish in specimens 7—8 mm length.

(3) *Larvae between 10 and 20 mm in length*: They are grouped into two on the basis of their snout length; the long snout group with shortnosed spearfish, sailfish and striped marlin, and the short snout group with black marlins. In the former, the snout length exceeds 1/5 of their body length, while in the latter, it does not. For the specific separation of the former group, Table A [Table 2]; shortnosed spearfish is distinguishable by black chromatophores on brachiostegal membrane, while sailfish is separated from striped marlin by the difference of their head profile: Unlike striped marlin with a straight snout, sailfish has a beak-like snout. And because of this difference in the shape of the snout, they are separable by the difference of the location of snout in terms of the center of eyes. In sailfish, the center of eyes is above the tip of snout, while in striped marlin, they are on a nearly same level.

Separation of white marlin from black marlin can be made on the basis of the form of the pectoral fin.

(4) *Larvae over 20 mm in length*: On top of the criteria of Table A [Table 2], the following characters, as listed in Table B [Table 3], can be applied.

In addition to the above, the distance between anus and the first anal fin insertion works as a reference to discriminate shortnosed spearfish. The distance of the said species is larger, as compared with other species.

The vertebral formulae are also useful tools for separation: the formulae of shortnosed spearfish, sailfish and striped marlin are all 12+12, while those of black and white marlin are 11+13.

Ueyanagi (1964) discussed some of the misidentifications of sailfish larvae made by Sun (1960). He stated that Sun's specimen No. 3 (Fig. 5c; 19.1 mm total length) and No. 4 (Fig. 5d; 30.5 mm total length) seem to be blue marlin. Ueyanagi (1959) pointed out that the larvae identified tentatively by Gehringer (1956) as *Tetrapturus belone* are probably blue marlin. He added that Gehringer's (1956) sailfish figures 6, 10, and 14 closely resemble striped marlin young and hence may be the larvae of the Atlantic cognate of striped marlin, namely white marlin, *T. albidus*.

Little is known on the biology of young sailfish.

Table 2.—Summary of the prominent diagnostic characters of istiophorid larvae less than 5 mm in length (from Ueyanagi,

Species characters	*Tetrapturus angustirostris*	*Istiophorus orientalis*	*Makaira mitsukurii*	*Eumakaira nigra*	*Marlina marlina*
Profile of head	Tip of snout is lower in level than center of eye.	Same to the left.	Tip of snout and center of eye are on a nearly equal level.	Tip of snout is lower in level than center of eye.	Same to the le
	Anterior edge of orbit does not project forward.	Same to the left.	Same to the left.	Anterior edge of orbit projects forward.	Anterior edge orbit does n project forw
Presence or absence of chromatophores on the branchiostegal membrane.	Present	Absent Chromatophores generally present on the peripheral zone of lower jaw membrane.	Absent	Absent	Absent
Pectoral fins	Fins extend along the lateral side of the body and can be readily folded against the side of the body.	Same to the left.	Same to the left.	Same to the left.	Fins stand out the lateral si the body at a angle and ca be folded ag the body wit breaking the

Table 3.—Diagnostic characters usable in distinguishing the istiophorid larvae more than 20 mm in standard lengt (from Ueyanagi, 1964).

Species characters	*Tetrapturus angustirostris*	*Istiophorus orientalis*	*Makaira mitsukurii*	*Eumakaira nigra*	*Marlina marlina*
Number of first dorsal fin-rays	More than 48	[1]43-47	Less than 45	Less than 45	Less than 45
Shape of first dorsal fin	Anterior-high type	Posterior-high type	Anterior-high type	Anterior-high type	Anterior-high type (presumed).
Lateral line	Single	Single	Single	Complex-having branches	Not single(2)[2] (obscure).

[1]This range is estimated from a small number of materials examined, so it seems to become somewhat wider with an increase o materials.

[2]Lateral line pattern is not yet ascertained.

Beebe (1941) reported that they primarily eat copepods. Voss (1953) identified the stomach contents of his specimens and found predominantly copepods and fish larvae. Gehringer (1956) also identified the stomach contents of his specimens and his results are shown in Table 4.

In summary, sailfish less than 10 mm long generally eat copepods, while fish larvae predominate in larger sizes.

3.3 Adult Phase

3.31 Longevity

De Sylva (1957) stated that natural mortality is undoubtedly high, pointing out that off Florida the sport fishery is dependent upon 1- and 2-yr-old fish. Mather et al. (1974) has recorded the recovery of a tagged sailfish that was at liberty for more than 4 yr. The fish was estimated to weigh 21.3 kg when tagged, w according to de Sylva would be about a 2-yr-old. specimen, therefore, was at least 6 yr old when re tured.

There is no information available for the Pacifi Indian oceans, although Koto and Kodama (1 found that the growth rate of sailfish in the China Sea is similar to that of sailfish in the Atla

3.32 Hardiness

Adult sailfish are probably adaptable to relati large changes in their environment and in their ab to utilize different food items.

3.33 Competitors

Although sailfish probably compete with man the large pelagic scombridlike fishes for food

ble 4.—Stomach contents of specimens from *Theodore N. Gill* collections (from Gehringer, 1956).

ze of fish (mm)	Contents
3.9	5 copepods
5.4	Parts of 9 copepods
6.0	3 copepods; parts of 7 copepods
6.0	7 copepods; parts of 1 fish larva
9.4	2 copepods; parts of 3 fish larvae (heads 1.4 mm)[2]
[1]10.1	1 fish larva,[2] 4.2 mm
10.1	1 copepod; part of 1 fish larva
[1]10.3	1 fish larva,[2] 4.8 mm
[1]10.3	1 fish larva, 6.0 mm; part of 1 fish larva[2]
[1]10.8	4 fish larvae,[2] approximately 5.4 mm
[1]11.3	Part of 1 fish larva (head, 3.0 mm)
[1]11.4	1 fish larva,[2] 6.6 mm; piece of another fish larva
[1]12.3	1 copepod; part of 1 unidentified crustacean; 1 fish larva,[2] 5.1 mm; part of 1 fish larva
[1]12.8	1 fish larva,[2] 7.2 mm
12.8	4 flying fish larvae, 4.8 to 6.0 mm; head of 1 fish larva
[1]13.0	2 copepods; 1 istiophorid larva, 6.0 mm
13.0	1 flying fish larva, 4.8 mm; parts of 2 other fish, one approximately 6.0 mm
[1]13.9	1 flying fish larva, 5.1 mm; 1 fish larva,[2] 7.2 mm
14.6	Part of 1 fish larva (head, 2.1 mm)
15.3	3 flying fish larvae, 3.6 mm; parts of 2 flying fish larvae
16.2	1 istiophorid larva (head, 2.4 mm)
16.9	Part of 1 fish larva (head, 1.8 mm)
16.9	1 fish larva, 8.4 mm
17.8	3 fish larvae, 3.0 mm, 3.6 mm, and 7.2 mm
18.2	1 thin fish larva, 11.4 mm; part of fish larva
20.9	Parts of fish larva
[1]21.0	1 istiophorid larva, 10.2 mm
27.4	Parts of fish larva
[1]34.5	Parts of fish larvae
38.8	8 flying fish larvae, 4.8 to 8.7 mm; 2 fish larvae, 6.3 mm. and 6.6 mm; parts of 3 fish larvae
[1]45.0	Parts of fish larvae
[1]45.0	Parts of fish larvae

[1]Unidentified species.

[2]All fish listed under footnote 2 appear to be identical, and unless otherwise noted, measurements given are of standard lengths.

awning areas, it is likely that they are relatively unfected by this competition. Among the billfishes, ilfish and black marlin, *Makaira indica*, are the minant inshore species in the Pacific (Howard and yanagi, 1965) and undoubtedly compete for food d living space, although black marlin probably ilize larger food items than do sailfish. Royce (1957) amined 11 black marlin captured in the Pacific and und fishes weighing up to 18 kg in the stomach con-nts as well as remains from other fishes which were obably too large for a sailfish to consume. ssimilarities in distribution are evident between ilfish and some of the other billfishes. Kume and seph (1969a), for example, showed that the centers of abundance of the shortbill spearfish, *Tetrapturus angustirostris*, and the sailfish in the eastern Pacific are widely separated. The shortbill spearfish did not occur in the catches of research vessels fishing within 370-550 km of the coast while the majority of the sailfish were caught within this limit.

Williams (1967) noted that, ". . . in East Africa longline catches of striped marlin were high and of sailfish low, whereas the sport fishery catches of striped marlin were low and of sailfish very high." Williams suggested that this was partly due to the different habits of the two species. Similar results were obtained by Merrett (1968a, 1971) from the same area, indicating that the sailfish was the most coastal of the five istiophorid species present in the equatorial western Indian Ocean.

In the Atlantic, however, Wise and Davis (1973) showed that during certain seasons of the year, centers of sailfish abundance coincide with centers of white marlin, *T. albidus*, and blue marlin, *M. nigricans*, abundance. Fox (1971) also indicated that sailfish and blue marlin distributions show a strong overlap.

3.34 Predators

There is little information about predation on adult sailfish. Baughman (1941a) reported on the sighting by another observer of a school of bottlenose porpoise, *Tursiops truncatus*, attacking and apparently consuming an adult sailfish in the Gulf of Mexico. Maéda (1967) stated that killer whales attack "marlins," a category which undoubedly includes sailfish. He added, however, that "marlins" are probably not preyed upon to any great extent. Merrett (1968a) commented that shark attacks on fishes hooked on longline gear are well known, but predation on free-swimming tunas and billfishes would be considerably less.

3.35 Parasites, diseases, injuries, and abnormalities

Silas (1967) and Silas and Ummerkutty (1967) summarized previous accounts of parasites of scombroid fishes (Table 5). Williams (1967) stated that he found thousands of euryphorids (copepods, Caligoidea) up to 10 mm in length on the skin of sailfish near the head and caudal areas. He also recorded a heavy infestation of ascarid nematodes in the stomach of one specimen. Ward (1954) listed the helminth parasites *Bothriocephalus manubriformis* (Linton), *Contracaecum histiophori* Yamaguti, and *Hirudinella marina* Garcin as occurring in the viscera of sailfish from Florida waters. Jones (1971) reported on crater wounds caused by the squaloid shark, *Isistius brasiliensis*, in istiophorids.

Billfish are often found with portions of their bill broken off. Marlins and swordfish have been reported as attacking vessels and burying their bills in wooden planking. Sailfish, however, are apparently not as

Table 5.—List of parasites found on sailfish (adapted from Silas, 1967, and Silas and Ummerkutty, 1967).

Locality	Parasite	Location on host
Monogenetic trematodes (Silas, 1967)		
Japan and Sri Lanka	*Capsala ovalis* (Goto)	body surface
West coast of Sri Lanka	*Capsala megacotyle* (von Linstow)	body surface
Japan	*Capsaloides sinuatus* (Goto)	gills
Digenetic trematodes (Silas, 1967)		
Japan	*Hemiurus sardiniae* (Yamaguti)	stomach
Cestodes (Silas, 1967)		
N.W. Atlantic	*Bothriocephalus manubriformis* (Linton)	intestines
Copepods Silas and Ummerkutty (1967)		
Japan	*Caligus quadratus* Shiino	?
?	*Caligus* sp. (Silas and Ummerkutty	behind eye
Pacific off Mexico	*Gloiopotes costatus* Wilson	body surface
Gulf of Mannar	*Gloiopotes longicaudatus* (Marukawa)	body surface and inside gill cover
Sri Lanka	*Gloiopotes watsoni* (Kirtisinghe)	body surface
Pacific off Mexico	*Pennella filosa* (Linnaeus)	buried in body
Sri Lanka	*Pennella instructa* (Wilson)	body surface
Formosa	*Pennella* sp. Ho	buried in body

guilty of aggressive behavior of this type as are marlins and swordfish.

Evans and Wares (1972) examined stomachs from 151 sailfish captured in the eastern Pacific and found 22% with stomach ulcers. They suggested that injuries from the spines of prey fishes were factors in the ulcerations.

3.4 Nutrition and Growth

3.41 Feeding

One of the most detailed accounts of sailfish feeding is found in Voss (1953):

> Apparently no feeding was done at night for in the early morning the fish were scattered. Around 9:00 the first schools began to appear as the sailfish, numbering from 6 or 8 to 25 or 30 began to mill about small groups of "pilchards," forcing them into compact schools by slowly circling about them at the surface, their sails half raised. While feeding the sailfish were oblivious to their own danger and a boat could be eased down into the mass of circling fish until often the sailfish would tually bump against the sides of the vessel. . . .
>
> At short intervals, while the "pilchards" were kept bun up by the circling sailfish, a single sailfish would break out o circle and swim rather slowly directly through the small sc of "pilchards," thrashing vigorously sideways with its bill, ting fairly large numbers of the small fish stunning or ki them. After thrashing through the school, the sailfish w then turn, swim slowly downward beneath the school whe would then swim about picking up the dead "pilchards" as sank downwards.

Voss added that in normal feeding, sailfish w relatively slow in their actions and bursts of sp were exceptions. He concluded that accounts sailfish being fast swimmers were exaggerated that speeds of over 13 knots were probably unusu

3.42 Food

It has been fairly well established that the foo very young sailfish consists primarily of copep (Beebe, 1941; Voss, 1953; Gehringer, 1956) (see 3. Voss noted that with only a few millimeters incre in size, the diet of young sailfish changed predominately fishes. Gehringer (1956) stated tha the 32 larvae he examined from the South Atla coast of the United States, fish were a major portio the diet in all larvae above 6.0 mm. He also noted t no copepods were found in sailfish above 13.0 mm l (15 specimens). The food of sailfish from the east and western Atlantic, off the West African shelf off northeast Brazil, was examined by Ovchinni (1970), who gave a list of food items taken from stomachs of sailfish from both areas. He stated t the composition of food from eastern Atlantic altered according to the season. He reported tha February the food was mainly cephalopo anchovies, and *Otoperca*, whereas in April sailfish on *Sardinella* and in May jack mackerel. He indica that forage resources were more scant in the o ocean with about half of the stomachs examined be found empty. The qualitative composition of the f was also different, squid and gempylids being main items.

The above confirms what appears to be a gen consensus that although fish and squid form the m jor portion of their diet, adult sailfish are fairly opp tunistic feeders and eat whatever happens to be p ent. Nakamura (1949) stated that billfishes in gen probably do not have any particular tastes in food probably feed on whatever was abundant in the a or what could be most easily seen or caught. He ad that billfish definitely did not feed on deme organisms although they did feed occasionally deep-sea fishes. He also stated that it is m reasonable to believe that deep-sea fishes are eate night when they approach the surface rather t billfishes diving deep to feed during the day. Ovc nikov (1970) presented data on the degree of stom fullness of sailfish off the West African shelf, which

icated that they fed most intensively during the vening hours. Again, judged on the degree of stomach llness, he found that they fed most intensively at ie beginning of February, in the second half of April, id in May. He observed that this sporadic intensity feeding was correlated inversely with periods of insified spawning.

Evans and Wares (1972) found mostly squid and shes in stomachs from sailfish captured in the astern Pacific. Voss (1953) stated, however, that ailfish often feed on typical bottom dwellers such as a robins (Triglidae) and gastropod mollusks (Table). He also added that one of the cephalopods, *Grimpteuthis* sp., found in the stomach contents, is a bottom dweller from rather great depths. Baughman 1941a, b) stated that sailfish examined from the estern Gulf of Mexico contained a large proportion shrimp in their stomachs.

3.43 Growth rate

Sailfish apparently have a very rapid growth rate. e Sylva (1957) studied the growth of sailfish in the lorida region based on length frequencies of almost ,000 young and adult sailfish. He stated that sailfish verage a total length of 17.8 cm at the end of their rst month, 50.8 cm by the end of the second month, 9 cm by the end of the third month, and 111.7 cm at ie end of 4 mo. He added that there was considerable ariation in these lengths due to variation in time of pawning and subsequent variation in availability nd abundance of planktonic food. By the end of their rst year of life sailfish average 183 cm in total length, 16 cm the second year, and 233.7 cm by the end of ie third year. De Sylva noted that determination of ge after 2 yr was difficult. Variation in weight is rge; a 3-yr-old fish with a modal length of 233.7 cm ay vary in weight from 19 to 49.4 kg. Jolley (1974) uggested the possibility of differential growth etween male and female Atlantic sailfish. This ypothesis partially explains the wide variation in eight and length at a given age described by de ylva. Jolley showed significant differences between ie length-weight relationship of males and females.

Koto and Kodama (1962) estimated growth of ailfish from the East China Sea. Their growth rates r categories "n, $n+1$, $n+2$" were similar to those esmates by de Sylva for 1-, 2-, and 3-yr-old fish. Koto nd Kodama also noted marked seasonal changes in owth rate with a maximum from June to December.

Williams (1970) discussed growth of sailfish off the ast of East Africa, and although his data were too w for a detailed analysis of growth, he noted that ased on de Sylva's work the majority of the sailfish ught off East Africa were 3 yr old or older. He stated at sailfish in the Indian and Pacific oceans unubtedly grow larger than Atlantic sailfish and this uld be due to a longer life span or some other feature growth.

Table 6.—Stomach contents of 241 adult specimens of the sailfish, *Istiophorus americanus* (from Voss, 1953).

Specimens	Number	Total
Scombridae (mackerels, etc.)		59
Euthynnus alletteratus (little tuna or false albacore)	35	
Acanthocybium solanderi (wahoo)	1	
Scomberomorus (*regalis* (?) painted mackerel)	1	
Hemiramphidae (halfbeaks)		42
Hemiramphus sp. (*brasiliensis*?) balao	16	
Exocoetidae (flying fish)		6
Cypselurus heterurus (Atlantic flying fish)	1	
Belonidae (needlefish)		25
Ablennes hians (flat needlefish)	2	
Strongylura notatus (common needlefish)	5	
Trichiuridae (cutlassfish)		14
Trichiurus lepturus (cutlassfish)	13	
Monocanthidae (filefish)		2
Monocanthus hispidus (common filefish)	2	
Tetradontidae (swell or blowfish)		1
Lagocephalus laevigatus (rabbitfish)	1	
Mugilidae (mullet, some undoubedly bait)		10
Mugil cephalus (black mullet)	1	
Mugil trichodon (silver mullet)	1	
Mugil curema (silver mullet)	3	
Priacanthidae (Bigeye)		1
Priacanthus sp. (catalufa)	1	
Clupeidae (herrings)		19
Sardinella anchovia (Spanish sardine)	11	
Engraulidae (anchovies)		2
Carangidae (jacks)		43
Decapterus macarellus (mackerel scad)	7	
Caranx ruber (blue runner)	9	
Oligoplites saurus (leather jacket)	3	
Sparidae (porgies)		12
Lagodon rhomboides (pinfish)	12	
Balistidae (triggerfish)		2
Balistes (*forcipatus*?) (spotted triggerfish)	1	
Balistes carolinensis (common triggerfish)	1	
Coryphaenidae (dolphins)		1
Coryphaena hippurus (dolphin)	1	
Triglidae (sea robins)		1
Pomatomidae (bluefish)		2
Pomatomus saltatrix (bluefish)	2	
Gerridae (mojarras)		1
Gadidae (cod)		1
Myrophidae (worm eel)		1
Unidentifiable fish		138
Cephalopoda (octopods and squid)		77
Octopoda (octopods)		
Argonautidae (paper nautilus)		
Argonauta argo (paper nautilus)	15	
Argonauta sp. (*hyans*?)	1	
Stauroteuthidae		
Grimpoteuthis (?)	11	
Decapoda (squid)		49
Ommastrephidae		
Sthonoteuthis bartrami (flying squid)	29	

3.5 Behavior

3.51 Migrations and local movements

Sailfish perform long-range migrations of thousands of miles as well as fairly extensive short-term movements most likely based on local environmental conditions.

Mather et al. (1974) reported on the results of 18 yr of billfish tagging under the Woods Hole Oceanographic Institution Cooperative Gamefish Tagging Program. Over 12,000 sailfish were tagged and 95 recoveries were recorded (Table 7). The longest distance between release and recovery was from off Cape Hatteras, N. C., to just north of Surinam, a distance of over 1,853 km (Fig. 5). The longest time at large was over 4 yr. Voss (1953) believed that there is a population of sailfish present off Florida all year long, but in summer with the extension of warm water northward there is a movement of sailfish northward along the inside edge of the Gulf Stream. With the arrival of the winter northerlies these fish are driven southward to regroup off the coast of Florida.

Squire (1974) reported on the results of a billfish tagging program in the eastern Pacific from 1954 through 1971. Over 4,700 sailfish were tagged; only 8 were recovered. Most recaptures were near the tagging locality and the longest period at large was 15 mo.

Nakamura et al. (1968) stated that sailfish in the western Pacific migrate from New Guinea, the Solomon Islands, and the Philippines along the Japanese Current to Japan and the Pacific coast of Mexico. In autumn sailfish enter the warm Tsushima Current and move into coastal waters where they a sometimes captured in traps. In the East China S Koto et al. (1959) reported that sailfish migra northward in summer and return southward autumn where they overwinter in the southernm area. They also reported that fish under 160 cm lo move into the area between May and July while fi over 160 cm move southward out of the area, probab for spawning.

In the Indian Ocean, Williams (1970) and Merr (1971) discussed the movements of sailfish off t coast of East Africa. Williams believed that there a two overlapping migratory patterns, one a localiz onshore-offshore migration with a superimpos migration from the north and northeast with the i trusion of the Somali Current. Merrett believed th the onshore migration was a postspawning, feedi migration.

3.52 Schooling

Sailfish occasionally form schools or groups of fro 3 to 25 or 30 individuals. More often, however, conce trations of sailfish occur in loose aggregations over broad area. Large numbers of sailfish occur off t southeast coast of Florida during the winter month Voss (1953) reported that in early morning sailfish the Florida coast were scattered, but by 0900 schoo of up to 30 fish began to form and feed on conce trations of small forage fish.

Furukawa (1961) presented a method of calculati the size of an average school or aggregation using t

Table 7.—Releases for Atlantic sailfish, *Istiophorus albicans*, in the western North Atlantic by years, and returns from these by months at liberty (from Mather et al. 1974).

Releases		Months at liberty								
Year	Number	0-0.9	1-1.9	2-5.9	6-11.9	12-17.9	18-23.9	24-35.9	36-47.9	Totals
1954	27									
1955	16				1					1
1956	0									
1957	24									
1958	28				1	1				2
1959	113									
1960	827		2	1	1	1				5
1961	1,157	1		1	2	2				6
1962	1,284	2	1	1	5	1				10
1963	1,162	3	1	4	1					9
1964	1,080	2	2	1	1					6
1965	1,093	2	1	2	3	1				9
1966	1,139	5		4	4		2	1	1	17
1967	828	2	2	7	1	1				13
1968	775	3	1	1	3	1	1			10
1969	763	1	1	2		1				5
1970	621	1								1
1971	1,068				1					1
Total	12,005	22	11	24	24	9	3	1	1	95



Figure 5.—Releases and recaptures of sailfish in the North Atlantic Ocean (from Mather et al., 1974).

number of sailfish caught by one longline. His estimates for the East China Sea and for the Okinawa region were 2,462 and 624, respectively, which was less than for striped marlin or black marlin but greater than for blue marlin or swordfish.

Ovchinnikov (1970) reported a high degree of schooling in sailfish off West Africa from the frequency of occurrence on neighboring hooks along a tuna longline. He found percentage catches on adjacent hooks extremely high; often sailfish were caught, or lost, from every hook for a length of several kilometers.

4 POPULATION

4.1 Structure

4.11 Sex ratio

Ueyanagi et al. (1970) examined 353 sailfish from the Atlantic and found 208 females and 145 males, and Nakamura (1971) also reported on predominance of females in the Gulf of Mexico sport fishery. He examined 81 sailfish over a period of 4 yr and found an overall ratio females to males of 2.4:1.

Nakamura (1949) stated that the sex ratio of sailfish in the East China Sea area was about even and did not change greatly throughout the year. Williams (1964) found a sex ratio males to females of 3.25:1 in 17 specimens captured by longline gear off East Africa. He later (Williams, 1970) examined sport catches off Malindi, Kenya, and found a sex ratio males to females of 2:1. Merrett (1971) examined 79 sailfish caught by longlines off East Africa and found males predominating, although not as greatly as in Williams' sample. In the inshore troll fishery, Merrett noticed no significant deviation from a 1:1 ratio.

4.12 Age composition

De Sylva (1957) concluded from his analysis of age and growth of sailfish off southern Florida that most of the population consisted of fish less than 3 yr old. He stated that the winter sport fishery is almost entirely dependent upon fish which are about 6 and 18 mo old. In the summer, the majority caught are 1- and 2-yr-old fish.

Williams (1970) stated that if growth rates for the Atlantic and Indian oceans were similar, then the majority of the sailfish caught off Kenya are over 3 yr old. He believed that the single mode in length frequencies of sailfish caught off East Africa undoubtedly included at least two successive year classes. Williams added that growth rates may not be similar, however, and the larger sizes of Pacific and Indian ocean sailfish may indicate much longer life or some other factor.

4.13 Size composition

De Sylva (1957) examined length frequencies of over 9,000 sailfish from off southern Florida (Fig The majority of the sport catch range from about to 140 cm trunk length (posterior edge of the orb the anterior insertion of the caudal keels) or 173 to cm total length (tip of upper jaw to vertical line dr between tips of caudal lobes) with a wide rang weight from 6.0 to 49.4 kg. The second largest grou from 61 to 94 cm trunk length averaging under 6 The small fish begin to appear in the catch in summer, becoming prominent in Novem December, and January. De Sylva stated that t fish are the incoming year class.

Ueyanagi et al. (1970) found a unimodal distri tion in length frequencies of sailfish caught longlines in the Atlantic. The majority were from to 165 cm body length (from posterior edge of orbi centralmost caudal rays). The largest sailfish cau by sport fishing gear from the Atlantic is a 55.8 specimen caught off the Bahamas in 1950.

Williams (1970) found a unimodal distributio length frequencies of sailfish caught in the sport f ery at Malindi, Kenya (Fig. 7). The majority ran from 203 to 254 cm fork length (tip of snout to c tral rays of the caudal fin) or from 224 to 279 cm t length (same as de Sylva's total length) with a wei range of from 18.1 to 47.2 kg. Williams reported t the Kenya record for 1957 weighed 62.1 kg. Mer (1971) examined 77 sailfish caught on longlines East Africa. He also found a unimodal distribut with the majority of the specimens between 160 185 cm body length (center of orbit to tip of shor caudal ray).

Koto et al. (1959) presented length frequencie sailfish caught by longlines in the East China (Fig. 8). Size ranged from 105 to 240 cm body le (tip of lower jaw to central rays of caudal fin). S percent were from 165 to 190 cm body length. The tribution is essentially unimodal except in June w a group of 125 to 150 cm fish suddenly appeared in catch. Koto et al. believed that these small fish e the East China Sea from other areas during month. The largest sailfish recorded on sport fish gear in the Pacific is a 100.2-kg specimen caught n the Galapagos Islands in 1947.

Length-weight relationships have been reporte de Sylva (1957), Merrett (1968b, 1971), Kume Joseph (1969b), Williams (1970), Nakamura (19 and Jolley (1974). These results are summarize Table 8. None but Jolley were able to detect differences between sexes. Nakamura (1949) st that sex differences, if any, were inconspicuous Pacific sailfish.

4.2 Abundance and Density (of Population)

4.21 Average abundance

Nothing found in the literature.



Figure 6.—Left, length-frequencies for Atlantic sailfish from southern Florida, 1953. Right, length-frequencies for Atlantic sailfish from southern Florida, 1954 (from de Sylva, 1957).



Figure 7.—Monthly length-frequencies of sailfish from Malindi, Kenya, pooled for the four seasons from 1958 through 1962. (Smoothed curves from three figure moving average.) (From Williams, 1970.)

Table 8.—Length-weight relationship of sailfish, *Istiophorus platypterus*.

Author	Measurement	Range	Log *a*	Log *b*
Merrett (1968b)	Eye-fork length in centimeters, weight in pounds	126-194 cm	-4.459	2.819
Kume and Joseph (1969b)	Eye-fork length in centimeters, weight in kilograms	134-205 cm	-3.9357	2.4156
Williams (1970)	Tip of snout to fork in inches, weight in pounds	60-100 inches	-2.1458	1.8896
Nakamura (1971)	Total length in inches, weight in pounds	70-110 inches	-4.68194	3.26326
Jolley (1974)	Trunk length in centimeters, weight in kilograms	70-144 cm (males)	-5.784	3.342
		42.5-151.5 cm (females)	-4.941	2.950



Figure 8.—Body length composition of sailfish in the East China Sea, by month from (Koto, Furukawa, and Kodama, 1959).

4.22 Changes in abundance

Changes caused by hydrographic conditions are eferred to in general terms in 2.3.

4.23 Average density

Howard and Ueyanagi (1965) indicated catches of *I. latypterus* by the number caught per tuna longline peration (approximately 2,000 hooks) for a com- osite year in the Pacific Ocean. In oceanic fishing perations the catches are shown to be generally 1.1- .0 fish per operation while closer to land masses the verage density was approximately 5.1-10.00 fish per peration in areas commonly fished by Japanese ongliners.

For comparison, an average hook-rate of 0.168 ailfish per 100 hooks was obtained from 154 longlin- ig operations in the equatorial western Indian Ocean uring 1964-67 (Merrett, 1971). This figure is similar o that (0.14 for 100 hooks) obtained by Williams 1967) during an earlier survey.

4.24 Changes in density

Kume and Joseph (1969a) showed that heavy long- ne fishing had affected the apparent abundance of ailfish in the waters off Central America. The initial ook-rate in 1964 was 10.6 per 100 hooks, 9.5 the llowing year, and 5.8 in 1966.

Williams (1970) examined the catches of a sport shery in the equatorial western Indian Ocean off the enya coast from 1958-68 and his results are shown in igure 9. He indicated that, with effort relatively able, the catch and catch per unit effort (CUE) aralleled one another over the 10-yr period. He ointed out, however, that while the CUE and catches ave increased since 1962, there was a sharp decline in ie mean (and median) weight of fish caught.

Kume and Joseph (1969a) indicated the seasonal ends in density reflected by the catches of sailfish



igure 9.—Sailfish effort, catch, and catch per unit effort for e 10 seasons from 1958 through 1968 at Malindi, Kenya (from illiams, 1970).

within 370-550 km of the coast in the tropical eastern Pacific. They showed that in the northern hemisphere the area of highest sailfish abundance is between Guatemala and the mouth of the Gulf of California. In the area around central Mexico sailfish appear abundant all year; during the first quarter this is the only area where they are. During the second quarter, there appears to be a southerly extension of the area of high abundance to as far south as lat. 10°N. This is maintained during the third quarter as well as a northward movement of fish to about lat. 20°-22°N. By the fourth quarter they reported a northward shift in the southern limit to about 15°N and in the northern limit to about lat. 25°N.

Evidence of seasonal changes in density of sailfish in the equatorial western Indian Ocean was given by Williams (1967, 1970) and Merrett (1968a, 1971). In the tuna longline survey off the East African coast during 1958-60, Williams (1967) found a mean hook-rate of 0.01 per 100 hooks during the southeast monsoon (April-October) while in the northeast monsoon (November-March) it rose to 0.31 per 100 hooks. Continuing this survey (1964-67) in a wider area of the equatorial western Indian Ocean, Merrett (1971) confirmed the earlier results by obtaining hook-rates of 0.11 per 100 hooks during the southeast monsoon and 0.35 per 100 hooks during the northeast monsoon. Williams (1970) indicated more specifically the seasonal variation in density off the Kenya coast. He showed that, with few exceptions, sailfish are restricted to the period October to March, with peak catches coming in December and January. The catches of the latter 2 mo constituted 47.6-77.3% of the seasonal totals.

4.3 Natality and Recruitment

4.31 Reproduction rates

Nothing found in the literature.

4.32 Factors affecting reproduction

Nothing found in the literature.

4.33 Recruitment

According to de Sylva (1957) some sailfish first enter the catches of the sport fishery in the southern Florida area during the winter at an age of about 6 mo. Based upon de Sylva's age determinations, the age of recruits to the East China Sea fishery reported upon by Koto and Kodama (1962) is likely to be approximately 1 yr old (modal size: 140-175 cm). In the sport fishery off Malindi, Kenya, Williams (1970) found a modal size of 249-cm total length (29.5-31.3 kg), which is slightly longer at the same weight than that of a 42-mo-old Atlantic fish. However, he pointed out that this mode may be expected to include at least two successive year classes and that a constant growth

rate over the range of the species is doubtful. Thus, factors affecting recruitment such as growth, predation, and migration act differentially in various stocks of sailfish throughout the world's oceans.

No estimates of recruitment have been found in the literature. Yet in the light of the high growth rate and high natural mortality suggested by de Sylva, it follows that the reproductive rate and recruitment rate should be higher than in slower growing, longer lived species of similar population size.

4.4 Mortality and Morbidity

4.41 Mortality rates

There have been no studies on the mortality of sailfish, although de Sylva (1957) stated that it seemed evident that natural mortality is high off southern Florida based on the fact that only two year classes support the sport fishery.

4.42 Factors causing or affecting mortality

See 3.34.

4.43 Factors affecting morbidity

See 3.35.

4.44 Relation of morbidity to mortality rates

Nothing found in the literature.

4.5 Dynamics of Population (as a Whole)

Nothing found in the literature.

4.6 The Population in the Community and the Ecosystem

There has been no detailed study of the ecology of *I. platypterus* over the whole range of its distribution. However, the direct relation between this species and some of the physical features of the biotope in certain localities have been discussed by Voss (1953), Gehringer (1956), Cadenat (1961), Ovchinnikov (1966, 1970), Williams (1967, 1970), Kume and Joseph (1969a), and Yurov and Gonzales (1971). On a broader basis these features are discussed fully by Parin (1968). The vertical zone of the community in which the sailfish lives is characterized by good illumination and is likely to be delimited below by temperature at the main thermocline (from 10-20 m to 200-250 m, depending on area). Temperature is apparently important also in the latitudinal distribution of the species and some authors, Ovchinnikov (1966, 1970) and Kume and Joseph (1969a), suggested that the 28°C isotherm is optional for the species, while Cadenat (1961) indicated that the increased abundance of sailfish in the inshore waters off the Ivory Coast coincided with approximately this temperature. The association of larval *I. platypter* with the warm waters of the Gulf Stream was dicated by Gehringer (1956) and of larvae and adu with the Kuroshio Current by Yabe (1953), Ueyana (1959), and Howard and Ueyanagi (1965). It is lik that the greatest effect salinity produces on t ecosystem in which sailfish live occurs at the bour aries of surface water masses and other regions which a rise in salinity occurs simultaneously with rise in temperature, which produces conditions cc ducive to high productivity. Aggregations of sailfish such areas have been shown off West Africa and the east coast of Brazil by Ovchinnikov (1966, 197 and by Ueyanagi et al. (1970).

5 EXPLOITATION

5.1 Fishing Equipment

5.11 Gear

The primary commercial fishing gear for sailfish the tuna longline. Basically, the tuna longline is drifting horizontal line of considerable length (up 120 km), from which single-hooked branch lines ha down at regular intervals. The gear and fishi methods are well known and have been described Yoshida (1966).

Trolling is the primary method used by spo fishermen for catching sailfish and in principal co sists of towing a baited hook or lure through the wat to simulate a swimming fish. The bait is trolled aste of a motor launch, and the line is loosely connected its passage inboard to a clip attached at the tip of laterally directed trolling pole (locally call "tangon" or "outrigger"). When a sailfish strikes t bait, the line snaps free from the clip. This mome tarily stops the bait's progress through the water a is supposed to resemble the effect of the prelimina strike on a live fish after which the sailfish turns a swallows the bait.

5.12 Boats

The types of fishing vassels used in tuna longlini have been described, again by Yoshida (1966). N special boat is required by sport fishermen. All that necessary is one which has an open stern where t angler may stand, a trolling pole on one or both side and the power to troll at up to 6 knots. Neverthele the design of many modern, sport fishing boats highly sophisticated, incorporating addition facilities for the fisherman.

5.2 Fishing Areas

5.21 General geographic distribution

See 5.22.

5.22 Geographic ranges

Although incidental in tuna longline catches
roughout the Atlantic and Indo-Pacific oceans,
ailfish are specifically sought around the edges of the
ontinental shelf in many tropical areas of the world.
he most important fisheries are concentrated in
outheastern Florida; the northern and northeastern
ulf of Mexico; the Bahamas; the Caribbean region;
enezuela; the eastern tropical Pacific between
outhern California and Chile; Hawaii; New Zealand;
astern Australia; Kenya to Cape Town; South Africa
nd Ghana to Senegal.

Since sport fishing is a leisure activity, the areas of
naximum fishing effort may or may not coincide with
ne areas of greatest density of sailfish. Within the
istributional range of sailfish, some areas of greatest
shing intensity may be correlated with the density of
uman population on the adjacent land mass. The
eaward extent of the fishery is similarly affected by
ne desires of the angler. Thus, in most cases, the
ange is governed by the optimum distance offshore
hich can be attained during a day trip.

5.23 Depth ranges

Merrett (1968a) showed that sailfish are caught
ith equal frequency on all hooks on a longline which
sh at depths from the surface down to about 160 m.

5.24 Conditions of the grounds

Nothing found in the literature.

5.3 Fishing Seasons

5.31 General pattern of seasons

Sailfish are caught during all seasons of the year.

5.32 Dates of beginning, peak, and end of seasons

Sailfish are most abundant in the Atlantic longline
atch in spring and summer (Wise and Davis, 1973).
he sport fishery in the western Atlantic captures
ailfish all year long with best fishing in winter and
arly spring.

Williams (1970) stated that the best sport fishing
r sailfish at Malindi, Kenya, is from October to
larch with peak catches in December and January.

Howard and Ueyanagi (1965) indicated that in the
astern Pacific sailfish are caught in good numbers off
capulco all year long, but peak fishing is during the
inter months.

5.33 Variation in date and duration of season

Williams (1970) stated that the start and finish of
ne period of peak abundance of sailfish in the sport
shery at Malindi, Kenya, shows a marked relationship to annual fluctuations in overall environmental conditions in the area.

5.4 Fishing Operations and Results

5.41 Effort and intensity

The oceanic longline fishery, initially a Japanese effort but recently joined by China, Cuba, Republic of Korea, Venezuela, and others, is the primary commercial fishery utilizing sailfish. The unit of effort commonly used in this fishery is number of fish caught per 100, 1,000, or 10,000 hooks fished.

In the Atlantic, peak Japanese fishing effort was in 1965 when more than 97 million hooks were fished, resulting in a catch of 118,000 "other marlins," a category which includes spearfish and sailfish. Catch per 10,000 hooks for this category rose from 39.0 in 1959 to a high of 189.0 in 1967.

Kume and Joseph (1969a) presented data on catches from the eastern Pacific that indicated a high abundance of sailfish in this area. Almost 330,000 sailfish were caught in 1965, and catch per 100 hooks was 9.4. Catch per effort declined from 10.6 per hundred hooks in 1964 to 5.8 in 1966. Total fishing effort in this area declined from a high of 62 million hooks in 1964 to 47 million in 1966.

5.42 Selectivity

Nothing found in the literature.

5.43 Catches

See 5.41.

6 PROTECTION AND MANAGEMENT

6.1 Regulatory (Legislative) Measures

6.11 Limitation or reduction of total catch

The State of Florida has a catch limit of two sailfish per day per person.

6.12 Protection of portions of population

Nothing found in the literature.

6.2 Control or Alteration of Physical Features of the Environment

Nothing found in the literature.

6.3 Control or Alteration of Chemical Features of the Environment

Nothing found in the literature.

6.4 Control or Alteration of Biological Features of the Environment

Nothing found in the literature.

6.5 Artificial Stocking

Nothing found in the literature.

7 POND FISH CULTURE

Not applicable.

LITERATURE CITED

BAUGHMAN, J. L.
1941a. Notes on the sailfish, *Istiophorus americanus* (Lacépède), in the western Gulf of Mexico. Copeia 1941: 33-37.
1941b. Scombriformes, new, rare, or little known in Texas waters with notes on their natural history or distribution. Trans. Tex. Acad. Sci. 24:14-26.

BEEBE, W.
1941. A study of young sailfish (*Istiophorus*). Zoologica (N.Y.) 26:209-227.

BEN-TUVIA, A.
1966. Red Sea fishes recently found in the Mediterranean. Copeia 1966:254-275.

BLOCH, M. E.
1793. Naturgeschichte der auslandischen Fische, Vol. 7, Berlin.

BLOCH, M. E., and J. G. SCHNEIDER.
1801. Systema ichthyologiae, inconibus CX illustratum, post obitum auctoris opus inchoatum absolvit, correxit, interpolavit Io. Gottlob Schneider, Berl., 584 p.

BOSC, L. A. G.
1803. Les poissons, les reptiles, les mollusques et les vers. Nouveau Dictionaire d'Histoire Naturelle (Paris) 13: 557-558.

CADENAT, J.
1961. Notes d'Ichthyologie ouest-africaine XXXIII,—Note sur les poissons-Épeé d'Côte d'Ivoire, et plus spécialement sur les "Voiliers". Bull. Inst. Fr. Afr. Noire, Ser. A, 23: 116-130.

CASTELNAU, F. L.
1861. Memoire sur les poissons de l'Afrique australe, Vol. 7, 78 p., Paris.

CUVIER, G.
1832. Sur le poisson appelé machaera. Nouv. Ann. Hist. Nat. (Paris) 1:43-49.

CUVIER, G., and A. VALENCIENNES.
1831. Histoire naturalle des poissons. Chez F. G. Levrault, Paris, Vol. 8, 509 p.

DERANIYAGALA, P. E. P.
1952. A colored atlas of some vertebrates from Ceylon. Ceylon Natl. Mus. Publ. 1:106-107.

DE SYLVA, D. P.
1957. Studies on the age and growth of the Atlantic sailfish, *Istiophorus americanus* (Cuvier), using length-frequency curves. Bull. Mar. Sci. Gulf Carribb. 7:1-20.
1963. Postlarva of the white marlin, *Tetrapturus albidus*, from the Florida Current off the Carolinas. Bull. Mar. Sci. Gulf Caribb. 13:123-132.

EVANS, D. H., and P. G. WARES.
1972. Food habits of striped marlin and sailfish off Mexico and southern California. U.S. Fish Wildl. Serv., Res. Rep. 76, 10 p.

FOOD AND AGRICULTURE ORGANIZATION OF TH UNITED NATIONS.
1950. Scientific and common names applied to tun mackerels and spear fishes of the world, with notes on th geographic distribution; a progress report on the compi tion of scientific and common names of important fo fishes. FAO (Food Agric. Organ. U.N.), Wash., D. 235 p.

FOWLER, H. W.
1932. A new species of sailfish, *Istiophorus brookei*, fr Tahiti. Proc. Acad. Nat. Sci. Phila. 84:403-404.

FOX, W. W.
1971. Temporal-spatial relationships among tunas a billfishes based on the Japanese longline fishery in t Atlantic Ocean, 1956-65. Univ. Miami, Sea Grant Tec Bull. 12, 78 p.

FURUKAWA, I.
1961. Studies on the tuna long line fishery in the east China Sea and Okinawa region — II. School compositio [In Jap., Engl. summ.] Bull. Jap. Soc. Sci. Fish. 566-577.

GEHRINGER, J. W.
1956. Observations on the development of the Atlant sailfish *Istiophorus americanus* (Cuvier) with notes on unidentified species of istiophorid. U.S. Fish Wild Serv., Fish. Bull. 57:139-171.
1970. Young of the Atlantic sailfish, *Istiophorus platypte us*. U.S. Fish Wildl. Serv., Fish. Bull. 68:177-190.

GOODE, G. B.
1883. Materials for a history of the sword-fish. Rep. U. Fish Comm. (1880). Part 8:289-394.

GOSLINE, W. A.
1968. The suborders of perciform fishes. Proc. U.S. Na Mus. 124:1-78.

GÜNTHER, A.
1873-74. Erster ichthyologischer Beitrage nach Exem plaren aus dem Museum Godeffroy. J. Mus. Godeffro Hamburg 1:169-173.
1880. An introduction to the study of fishes. Adam a Charles Black, Edinb., 720 p.

HOWARD, J. K., and S. UEYANAGI.
1965. Distribution and relative abundance of billfish (*Istiophoridae*) of the Pacific Ocean. Univ. Miami Ins Mar. Sci., Stud. Trop. Oceanogr. 2, 134 p.

JOLLEY, J. W., JR.
1974. On the biology of Florida east coast Atlantic sa fish (*Istiophorus platypterus*). *In* R. S. Shomura and Williams (editors), Proceedings of the International Bi fish Symposium, Kailua-Kona, Hawaii, 9-12 August 197 Part 2. Review and Contributed Papers, p. 81-88. U.S. De Commer., NOAA Tech. Rep. NMFS SSRF-675.

JONES, E. C.
1971. *Isistius brasiliensis*, a squaloid shark, the probab cause of crater wounds on fishes and cetaceans. Fis Bull., U.S. 69:791-798.

JONES, S.
1959. Notes on eggs, larvae and juveniles of fishes fr Indian waters. II. *Istiophorus gladius* (Broussone Indian J. Fish. 6:204-210.

JONES, S., and M. KUMARAN.
1964. Distribution of larval billfishes (Xiphiidae and Ist phoridae) in the Indo-Pacific with special reference to t collections made by the Danish Dana Expedition. Pr

Symp. Scombroid Fishes, Part 1, p. 483-498. Mar. Biol. Assoc. India, Mandapam Camp.

ORDAN, D. S., and B. W. EVERMANN.
1926. A review of the giant mackeral-like fishes, tunnies, spearfishes and swordfishes. Calif. Acad. Sci., Occas. Pap. 12:1-113.

KOTO, T., I. FURUKAWA, and K. KODAMA.
1959. Studies on the tuna longline fishery in the East China Sea - III. Ecological studies on the sailfish. [In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. Lab. 10:88-106.

KOTO, T., and K. KODAMA.
1962. Some considerations on the growth of marlins, using size-frequencies in commercial catches. I. Attempts to estimate the growth of sailfish. [In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. Lab. 15:97-107.

KUME, S., and J. JOSEPH.
1969a. The Japanese longline fishery for tunas and billfishes in the eastern Pacific Ocean east of 130°W, 1964-1966. [In Engl. and Span.] Bull. Inter-Am. Trop. Tuna Comm. 13:277-418.
1969b. Size composition and sexual maturity of billfish caught by the Japanese longline fishery in the Pacific Ocean east of 130°W. [In Engl., Jap. summ.] Bull. Far Seas Fish. Res. Lab. (Shimizu) 2:115-162.

LACEPEDE, B. G. E.
1802. Histoire naturelle des poissons. Plasson, Paris, Vol. 2, 558 p.

LAMONTE, F., and D. E. MARCY.
1941. Swordfish, sailfish, marlin and spearfish. Ichthyol. Contrib. Int. Game Fish Assoc. 1(2):1-24.

LATREILLE, P. A.
1804. Tableau méthodique des poissons. Nouveau Dictionaire d'Histoire Naturelle (Paris) 24:104.

LAURS, R. M., and R. N. NISHIMOTO.
1970. Five juvenile sailfish, *Istiophorus platypterus*, from the eastern tropical Pacific. Copeia 1970:590-594.

LÜKTEN, C. F.
1880. Spoilia Atlantica. Bidrag til kundskab om formforandringer hos fiske under deres vaext og unviklong, saerligt hos nogle af Atlanter havets hojsofiske. [In Dan., Fr. summ.] K. Danske Vidensk. Selsk. Skr. 5 ser 12:441-447, 592-593. (English translation of the French summary published as: Lütken, C. F. 1881. Spoilia Atlantica: Contributions to the knowledge of the changes of form in fishes during their growth and development, especially in the pelagic fishes of the Atlantic. Ann. Mag. Nat. Hist. Ser. 5, 7:1-14, 107-123.)

MAÉDA, H.
1967. Distribution pattern of fish in relation to fishing method, with special reference to that of tuna along longline. Proc. Symp. Scombroid Fishes, Part 3, p. 1025-1041. Mar. Biol. Assoc. India, Mandapam Camp.

MATHER, F. J., III, D. C. TABB, J. M. MASON, JR., and H. L. CLARK.
1974. Results of sailfish tagging in the western North Atlantic Ocean. *In* R. S. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972, Part 2. Review and Contributed Papers, p. 194-210. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-675.

MERRETT, N. R.
1968a. Tuna longline survey in the equatorial western Indian Ocean. East Afr. Agric. For. J. 34:17-65.
1968b. Weight-length relationships for certain scombroid fishes from the equatorial western Indian Ocean. East Afr. Agric. For. J. 34:165-169.
1970. Gonad development in billfish (Istiophoridae) from the Indian Ocean. J. Zool. (Lond.) 160:355-370.
1971. Aspects of the biology of billfish (Istiophoridae) from the equatorial western Indian Ocean. J. Zool. (Lond.) 163:351-395.

MITO, S.
1966. Nihon kaiyo plankton zukan—Dai 7 Kan—Gyoran—Chigyo (Japanese marine plankton picture book—Volume 7—Fish eggs—Larvae). [In Jap.] Soyosha, Tokyo, 75 p.
1967. Some ecological notes on the planktonic fish larvae. [In Jap., Engl. abstr.] Inf. Bull. Planktol. Japan 14: 33-49.

MORROW, J. E.
1964. Marlins, sailfish and spearfish of the Indian Ocean. Proc. Symp. Scombroid Fishes, Part 1, p. 429-440. Mar. Biol. Assoc. India, Mandapam Camp.

MORROW, J. E., and S. J. HARBO.
1969. A revision of the sailfish genus *Istiophorus*. Copeia 1969:34-44.

NAKAMURA, E. L.
1971. An analysis of the catches and biology of big game fishes caught by the New Orleans Big Game Fishing Club, 1966-1970. East Gulf Sport Fish. Mar. Lab. Rep., 38 p.

NAKAMURA, E. L., and L. R. RIVAS.
1972. Big game fishing in the northeastern Gulf of Mexico during 1971. Natl. Mar. Fish. Serv., Panama City, Fla., 20 p.

NAKAMURA, H.
1932. The ripe eggs of the sailfish, *Istiophorus orientalis* (Temminck and Schlegel). [In Jap.] Zool. Mag. 44: 244-245.
1940. On the spawning habits of sailfish. [In Jap.] Zool. Mag. 52:296-297.
1942. Habits of fishes of the family Istiophoridae in Formosan waters. [In Jap.] Suisan Gakkwai Ho 9:45-51.
1949. The tunas and their fisheries. [In Jap.] Takeuchi Shobō, Tokyo, 118 p. (Translated by W. G. Van Campen, 1952, U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 82, 115 p.)

NAKAMURA, I., T. IWAI, and K. MATSUBARA.
1968. A review of the sailfish, spearfish, marlin and swordfish of the world. [In Jap.] Kyoto Univ., Misaki Mar. Biol. Inst., Spec. Rep. 4, 95 p.

OVCHINNIKOV, V. V.
1966. The effect of oceanographic conditions on distribution of the sailfish, *Istiophorus americanus*, off the West African coast. Oceanology 6:566-567.
1970. Mech-ryba i parusnikovye (Atlanticheskii okean. Ekologiya i funktsional'naya morfologiya) (Swordfishes and billfishes in the Atlantic Ocean. Ecology and functional morphology). Atl. Naucho-Issled. Inst. Rybn. Khoz. Okeanogr., Kaliningrad, 106 p. (Translated by H. Mills, Israel Program Sci. Transl., 1971, 77 p.; available U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA, as TT 71-50011.)

PARIN, N. V.
1968. Ikhtiofauna okeanoskoi epipelagiali (Ichthyofauna of the epipelagic zone). Akad. Nauk SSSR, Tr. Inst. Okeanol. (Translated by M. Raven, Israel Program Sci. Transl., 1970, 206 p.)

ROYCE, W. F.
1957. Observations on the spearfishes of the central Pacific. U.S. Fish Wildl. Serv., Fish. Bull. 57:497-554.
RÜPPEL, M. E.
1835. Mémoire sur une nouvelle espèce de poisson du genre *Histiophore*, de la Mer Rouge. Trans. Zool. Soc. Lond. 2:71-74.
SHAW, G., and F. P. NODDER.
1791. *Xiphias platypterus*. The broadfinned swordfish. Natural Misc. 28.
SILAS, E. G.
1967. Parasites of scombroid fishes. Part I. Monogenetic trematodes, digenetic trematodes, and cestodes. Proc. Symp. Scombroid Fishes, Part 3, p. 799-875. Mar. Biol. Assoc. India, Mandapam Camp.
SILAS, E. G., and A. N. P. UMMERKUTTY.
1967. Parasites of scombroid fishes. Part II. Parasitic copepoda. Proc. Symp. Scombroid Fishes, Part 3, p. 876-993. Mar. Biol. Assoc. India, Mandapam Camp.
SQUIRE, J. L., JR.
1974. Migration patterns of Istiophoridae in the Pacific Ocean as determined by cooperative tagging programs. *In* R. S. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972. Part 2. Review and Contributed Papers, p. 226-237. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-675.
STRASBURG, D. W.
1970. A report on the billfishes of the central Pacific Ocean. Bull. Mar. Sci. 20:575-604.
SUN, T. G.
1960. Lichinki i mal'ki tuntsov, parusnikov i mech-ryby (Thunnidae, Istiophoridae, Xiphiidae) tsentral'noi i zapadnoi chasti Tikhogo Okeana [Larvae and fry of tuna, sailfish and sword-fish (Thunnidae, Istiophoridae, Xiphiidae) collected in western and central Pacific.] [In Russ.] Tr. Inst. Okeanol. Akad. Nauk SSSR 41:175-191. (Engl. transl. by W. L. Klawe, Inter-Am. Trop. Tuna Comm., 1960, 18 p.)
TEMMINCK, C. J., and H. SCHLEGEL.
1844. Pisces. *In* P. F. von Seibold, Fauna Japonica, 323 p.
UEYANAGI, S.
1959. Larvae of the striped marlin, *Makaira mitsukurii* (Jordan et Snyder). [In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. Lab. 11:130-146. (Translated by W. G. Van Campen, 1961, 20 p.; available at Southwest Fisheries Center, National Marine Fisheries Service, NOAA, Honolulu, HI 96812.)
1963a. Methods for identification and discrimination of the larvae of five istiophorid species distributing in the Indo-Pacific. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 17:137-150.
1963b. A study of the relationships of the Indo-Pacific istiophorida. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 17:151-165.
1964. Description and distribution of larvae of five istiophorid species in the Indo-Pacific. Proc. Symp. Scombroid Fishes, Part 1, p. 499-528. Mar. Biol. Assoc. India, Mandapam Camp.
UEYANAGI, S., and H. WATANABE.
1962. Methods of identification for larval billfishes a young tunas (I). [In Jap.] Prelim. Rep. Nankai R Fish. Res. Lab., 17 p.
1964. Methods of identification of larvae of tunas a young billfishes (II). [In Jap.] Materials for Tuna Fi eries Research Council, Nankai Reg. Fish. Res. Lab., H chi, 16 p. (English translation of sections pertaining to l val tunas available at National Marine Fisheries Servi Honolulu, HI 96812.)
UEYANAGI, S., S. KIKAWA, M. UTO, and Y. NISHIKAWA
1970. Distribution, spawning, and relative abundance billfishes in the Atlantic Ocean. [In Jap., Engl. syno Bull. Far Seas Fish. Res. Lab. (Shimizu) 3:15-55.
VICK, N. G.
1963. A morphometric study of seasonal concentrations the sailfish, *Istiophorus albicans*, in the northern Gulf Mexico, with notes on other Gulf istiophorids. Texas & M Univ., Dep. Oceanogr. Meteor. Rep., Proj. 286- Ref. 63-14T, 41 p.
VOSS, G. L.
1953. A contribution to the life history and biology of t sailfish, *Istiophorus americanus* Cuv. and Val., in Flori waters. Bull. Mar. Sci. Gulf Caribb. 3:206-240.
WARD, H. L.
1954. Parasites of marine fishes of the Miami regi Bull. Mar. Sci. Gulf Caribb. 4:244-261.
WHITLEY, G.
1939. Ichthyological genotypes: Some supplementary marks. Aust. Zool. 8:189-192.
WILLIAMS, F.
1964. The scombroid fishes of East Africa. Proc. Sym Scombroid Fishes, Part 1, p. 107-164. Mar. Biol. Ass India, Mandapam Camp.
1967. Longline fishing for tuna off the coast of East Afri 1958-1960. Indian J. Fish., Sect. A, 10:233-390.
1970. The sport fishery for sailfish at Malindi, Keny 1958-1968, with some biological notes. Bull. Mar. S 20:830-852.
WISE, J. P., and C. W. DAVIS.
1973. Seasonal distribution of tunas and billfishes in t Atlantic. U.S. Dep. Commer., NOAA Tech. Rep. NM SSRF-662, 24 p.
YABE, H.
1953. On the larvae of sailfish, *Istiophorus orientalis* c lected in the South-western Sea of Japan. [In Jap., En synop.] Contrib. Nankai Reg. Fish. Res. Lab. 6, 10 (Engl. transl. available at Southwest Fisheries Cent National Marine Fisheries Service, NOAA, Honolulu, 96812.)
YOSHIDA, H. O.
1966. Tuna fishing vessels, gear, and techniques in t Pacific Ocean. *In* T. A. Manar (editor), Proc. Gc Conf. Cent. Pac. Fish. Resour., p. 67-89. State of H waii.
YUROV, V. G., and J. C. GONZALES.
1971. Possibility of developing a sailfish fishery in the eas ern Pacific Ocean. *In* Sovetsko-Kubinskie Rybokhozi istvennye Issledovaniya, 3:104-110. [In Russ., Spa summ.] Pishchevaya Promyshlennost, Moscow.

Synopsis of Biological Data on the Mediterranean Spearfish, *Tetrapturus belone* Rafinesque[1]

DONALD P. de SYLVA[2]

1 IDENTITY

1.1 Nomenclature

1.11 Valid name

The valid scientific name of this taxon is *Tetrapturus belone* Rafinesque, 1810. The synonymy of this species has been recently treated by de Sylva (1973) and is reproduced here with slight changes.

1.12 Synonymy

Tetrapturus belone Rafinesque, 1810.

Tetrapturus belone Rafinesque, 1810, Caratteri, p. 54-55, pl. 1, Fig. 1 (original description; type locality: Sicily). Robins and de Sylva, 1963, Bull. Mar. Sci. Gulf Caribb. 13(1):84-122 (redescription; neotype based on a 1,268-mm specimen from Sicily, USNM 196527.

Skeponopodus typus Nardo, 1833, Isis (Oken) 26(4): 416-419 (type locality: Adriatic Sea).

Tetrapterurus belone Bonaparte, 1841, Icon. Fauna Ital. 3(1):19 (emended orthographic spelling of *Tetrapturus* Rafinesque).

Tetrapterus belone Agassiz, 1843, Recherches Poiss. Foss. 5:7, 89-90, table E (emended spelling).

Tetraplurus belone Vérany, 1847, Atti Ott. Riun. Sci. Ital. Genoa, p. 492-494 (Camogli, Ligurian Sea; misprint for *Tetrapturus*?).

Histiophorus belone Günther, 1860, Cat. Fishes 2:513 (new combination).

Scheponopodus prototypus Canestrini, 1872, Fauna Italia (3):112 (type locality: Italy; variation of spelling of *Skeponopodus* Nardo, 1833).

Histiophorus (*Tetrapturus*) *belone* Lütken, 1876, J. Zool. (Gervais) 5:60-63, pl. 3 (*Tetrapturus* a subgenus of *Histiophorus*).

Tetrapturus imperator Goode, 1883, Rep. U.S. Comm. Fish Fish. for 1880: 306-307 (*T. belone* erroneously placed in synonymy of *T. imperator* Schneider, a synonym of and based upon a drawing of *Xiphias gladius*, copied from Aldrovandi).

Makaira belone Tortonese, 1958, Atti Soc. Ital. Sci. Nat. 97(4):330 (new combination).

Tetrapturus beloni [sic] Briggs, 1958, State Mus. 2(8): 287 (erroneously listed from Florida; *non* Rafinesque but *pfluegeri*, as shown by Robins and de Sylva, 1963).

1.2 Taxonomy

The identity of this species has been very uncertain. For example, LaMonte and Marcy (1941:21-22) suggested that *belone* merely represented the young of some other billfish. Lack of specimens precluded determination of the identity of this taxon until Robins and de Sylva (1960), thanks largely to the late Al Pflueger, presented data on 23 western Atlantic specimens which they tentatively identified as *T. belone*. In 1960 and 1961 C. Richard Robins and the late John K. Howard visited Mediterranean countries to obtain specimens, especially from Sicily—the type locality of *T. belone*. Subsequently, Robins and de Sylva (1963) presented data on 35 specimens of *T. belone* from the Mediterranean, redescribed the species, and designated a neotype.

1.21 Affinities

The phylogenetic relations of this species have been reviewed by Robins and de Sylva (1960, 1963). This species clearly has affinities with other small spearfishes and is somewhat closely related to the sailfish genus *Istiophorus*, showing fewer affinities to the genus *Makaira* (*sensu stricto*). It is most closely related to the shortbill spearfish, *Tetrapturus angustirostris* Tanaka, of the Indo-Pacific and is more distantly related to the longbill spearfish, *T. pfluegeri* Robins and de Sylva, of the Atlantic. Its relationships with the roundscale spearfish, *T. georgei* Lowe, are discussed by Robins (1974a). Within the genus, its most distant relatives are the white marlin, *T. albidus* Poey, and the striped marlin, *T. audax* (Philippi).

The genus *Tetrapturus* is briefly characterized by having: the anterior lobe of the spinous dorsal fin higher than the body depth at the dorsal-fin origin, with the lobe often rounded; 12 precaudal plus 12 caudal vertebrae; flesh red; and the size usually small, usually less than 300 pounds (136.1 kg).

A synonymy of *Tetrapturus* was given by Robins

[1]Contribution No. 1834, University of Miami, Rosenstiel School of Marine and Atmospheric Science.

[2]Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, FL 33149.

and de Sylva (1963:385). Because Robins (1974b) repeats this synonymy for *Tetrapturus pfluegeri*, it is not duplicated here.

Artificial key to species of the adult stages (over 10 pounds or 4.5 kg) of *Tetrapturus*

A. Scales along sides of body between pectoral fin and anal fin flexible, broadly rounded anteriorly, with 2 or 3 flattened, soft projections posteriorly . . . . . roundscale spearfish, *Tetrapturus georgei* Lowe

A. Scales along body between pectoral fin and anal fin stiff, decidedly lanceolate, pointed anteriorly, with flattened, hard projections posteriorly . . . . . B

B. Spinous dorsal fin high throughout, the posterior rays about equal in height to the anterior lobe; anus anterior to anal-fin origin by a distance equal to or greater than the anal-fin height; dorsal profile, from dorsal-fin region to near tip of bill, straight; dorsal fin unspotted, body bands absent or faint; premandibular portion of bill shorter than distance from tip of mandible to eye C

B. Spinous dorsal fin low posteriorly, except in small juveniles (less than 10 pounds or 4.5 kg); anus closer to anal-fin origin, distance between them decidedly less than anal-fin height; dorsal profile strongly arched over and behind eyes except in very small juveniles; dorsal fin spotted, the spots usually fading after death but usually apparent in dorsal groove; body bands prominent; premandibular portion of bill longer than distance from tip of mandible to eye . . . . . D

C. Dorsal spines 39 to 46; Mediterranean Sea . . . . . .Mediterranean spearfish, *Tetrapturus belone* Rafinesque

C. Dorsal spines 45 to 53; Atlantic or Indo-Pacific . . . . . E

D. Tips of spinous dorsal, anal, and pectoral fins strongly rounded; height of first dorsal fin less than depth of body at first dorsal-fin origin: Atlantic . . . . . white marlin, *Tetrapturus albidus* Poey

D. Tips of spinous dorsal, anal, and pectoral fins pointed; height of first dorsal fin equal to depth of body at first dorsal-fin origin; Indo-Pacific . . . . . striped marlin, *Tetrapturus audax* (Philippi)

E. Bill short, only slightly exceeding length of lower jaw, premandibular portion of bill contained three times or more in preorbital distance; pectoral fin short, decidedly shorter than pelvic fin length; Indo-Pacific . shortbill spearfish, *Tetrapturus angustirostris* Tanaka

E. Bill long, decidedly longer than lower jaw, premandibular portion of bill contained less than three times in preorbital distance; pectoral fin long in adults, almost equal to pelvic fin in length; Atlantic . .longbill spearfi *Tetrapturus pfluegeri* Robins and de Sy

Tetrapturus belone Rafinesque

Type specimen: It is not known if a type specimen so designated in the generalized account given by finesque (1810:54-55, pl. 1, Fig. 1), or indeed if it ever preserved.

Neotype: USNM 196527, a male, body length 1, mm, collected 2 August 1961 (Robins and de Syl 1963:93, Fig. 4, p. 93).

Type locality: Sicily, 200 m off the coast of Punta Ranieri, Strait of Messina.

Diagnosis: The diagnosis of *T. belone* is that given Robins and de Sylva (1963:90-92) as follows:

> First dorsal elements 39-46 (usually 41-46), second do elements 5-7 (usually 6), first anal elements 11-15 (usually 14), second anal elements 6 or 7, pectoral rays 16-20 (usuall or 18); second anal elements usually equal to or one more th second dorsal elements. Vertebrae 24 (12 precaudal, 12 caud Anus far anterior to anal-fin origin, distance 7.8-11 per c body length (tip of lower jaw to fork of tail), the distance eq to or exceeding height of first anal fin. First dorsal fin unspot and with high profile, especially in juveniles, its 25th elem varying from 13 per cent of body length (at 1268 mm) to 5 c per cent at 1700 mm or longer. Pectoral fin small (10-13 per c body length throughout range studied). Dorsal profile straig from first dorsal-fin origin to in front of eyes. Bill short, the d tance from tip of upper jaw to anterior margin of eye 20-15 cent of body length (the lower figures characterizing the lar specimens so that the growth of bill shows negative allometr

The description of *T. belone*, based on specimens, is from Robins and de Sylva (1963:92-9 as follows:

> . . . All [specimens] had 12 caudal and 12 precaudal vertebr As the total number of 24 is not known to vary in t Istiophoridae, only caudal vertebrae were counted on specimens; all had 12.
>
> The general body form of *T. belone* changes with growt Specimens of moderate size (Fig. 4 [upper]; Robins and Sylva, 1961 [1960]: Fig. 3a) are not unlike *pfluegeri*, althou *belone* is heavier (Table 6). As they become larger they beco more robust forward (Fig. 5) and in this respect resemb *angustirostris*. The dorsal profile from the origin of the spino dorsal fin to the base of the bill is straight (Fig. 1 a-c), a featu enhanced by the usual slope of the partly folded dorsal fin. *belone* does not have a dorsal hump as in *T. albidus* (see Robi and de Sylva, 1961 [1960]: Fig. 3c). The dorsal and anal profil are subparallel except in the larger specimens, which are deep anteriorly. Both *T. belone* and *T. pfluegeri* are flat- or slab-sid but *belone* is thicker (in width) (see graph, Fig. 3) and propo tionately deep posteriorly (Fig. 4).
>
> The dorsal fin is lower throughout than in *T. pfluegeri* thou the form is much the same. Its anterior height varies from 13- per cent of body length (at 766 to 1500 mm body lengt gradually decreasing to 12-13 from 1500-1900 mm. This is gradual trend with considerable variation. The 25th spine is co respondingly reduced and as the posterior spines do not co

tinue to grow the reduction in percentage of body length is great, from 16 per cent (at 766 mm body length) to 5.1 per cent (at 1868 mm body length). The corresponding proportional reduction, though slight, in anterior height of the spinous dorsal, coupled with increasing depth at the origin of the spinous dorsal, means that in the 1000- to 1500-mm range the fin is higher than the corresponding depth, whereas the reverse is true in the larger specimens.

To show the overall height of the spinous dorsal fin, additional measurements were made in two instances. Med-22: spine 14 (97 mm), 20 (107), 25 (122), 30 (108), 35 (102). Med-26: spine 10 (99), 16 (107), 20 (122), 25 (117, slightly broken), 30 (26), 35 (81).

The first anal fin is low, its anterior height less than the distance from its base to the anus (in one specimen the two distances are equal). Proportionally, the first anal fin varies from 7.2 to 9.3 per cent of body length with no allometry in the size series studied.

A characteristic feature of *T. belone* is its short pectoral fin. It varies somewhat in length and in shape (see Figs. 4-5) but is always quite short (10-13 per cent of body length) and has the upper (or anterior) edge curved, the lower (or posterior) edge nearly straight. The arch of the lateral line ends between the mid-point and the tip of the pectoral fin.

The body is not banded (if bands are present they must not be prominent), but fresh material was not seen by the authors. Body color varied from dark slate above to dull whitish or grey below. The dorsal fin was unspotted in all specimens. While it may be argued that spots may fade, we always found spots in that portion of the dorsal fin below the edge of the groove in the dorsum in each species as white marlin and sailfish, even in very poorly treated specimens. Also, Col. Howard saw some specimens within a few hours of their capture, and noted that none had bands on the body or spots on the spinous dorsal fin. An unspotted dorsal characterizes *T. belone* as it does *pfluegeri* and probably *angustirostris*.

The flesh is pale in *belone* except for the tissue under the lateral line. The importance of this feature in istiophorid taxonomy is uncertain for it may reflect nothing more than feeding habits.

The gas bladder was examined in most specimens and its chambers were arranged in two rows, one layer deep, extending back to the level of the second anal fin. The species has a large pineal window nearly the size of a half dollar.

One juvenile (UMML 11056, Med-36) 766 mm in body length is available to us. Morphometric data are not included in Figure 3 but are available in the tables. At this size it has the characteristic features of the larger specimens.

Material examined:

Spec. Med-2: 1,340 mm body length, weight 25.1 pounds (11.4 kg), male. Zona Scaletta, south of Messina, Aug. 7, 1961 (pectoral girdle, UMML 11074).

Spec. Med-3: 1,290 mm, 26.4 pounds (12.0 kg), male, Torre Faro, Aug.-Sept. 1961.

Spec. Med-4: 1,545 mm, 40.7 pounds (18.5 kg), male, off Sicily, Aug.-Sept. 1961.

Spec. Med-5: 1,670 mm, 50.6 pounds (23.0 kg), male, Strait of Messina, Aug. 7, 1961.

Spec. Med-6: 1,730 mm, 51.7 pounds (23.5 kg), male, Strait of Messina, Aug.-Sept. 1961.

Spec. Med-7: 1,680 mm, 59.4 pounds (26.9 kg), male, Strait of Messina, Aug.-Sept. 1961.

Spec. Med-8: 1,430 mm, 24.2 pounds (11.0 kg), male, off coast of Messina, south side, Aug. 8, 1961.

Spec. Med-9: 1,500 mm, 29.5 pounds (13.4 kg), male(?), Scaletta, Aug. 1961.

Spec. Med-10: 1,522 mm, 28.5 pounds (12.9 kg), male, Tropea, Aug. 9, 1961.

Spec. Med-11: 1,715 mm, 46.2 pounds (21.0 kg), sex unknown, Strait of Messina, Aug. 7, 1961 (piece of skin, UMML 11075).

Spec. Med-12: 1,745 mm, 57.2 pounds (26.0 kg), male, coast of Calabria, region of Calabria, Aug. 7, 1961.

Spec. Med-13: 1,855 mm, 60.5 pounds (27.4 kg), female, Strait of Messina, Aug. 17, 1961.

Spec. Med-14: 1,565 mm, 41.8 pounds (19.0 kg), male, Scaletta, southern section of Straits of Messina.

Spec. Med-15: 1,450 mm, 34.1 pounds (15.5 kg), male, Strait of Messina, Aug. 10, 1961.

Spec. Med-16: 1,345 mm, 23.5 pounds (10.7 kg), male, Strait of Messina, Aug. 1961.

Spec. Med-17: 1,265 mm, 17.6 pounds (8.0 kg), male, Strait of Messina, Aug. 3, 1961.

Spec. Med-18: 1,350 mm, 24.2 pounds (11.0 kg), male, Sicily, Aug. 1961.

Spec. Med-19: 1,435 mm, 26.4 pounds (12.0 kg), sex unknown, Straits of Taormina, Aug. 7, 1961.

Spec. Med-20: 1,510 mm, 30.4 pounds (13.8 kg), female, Strait of Messina, Aug. 4, 1961.

Spec. Med-21: 1,465 mm, 32.8 pounds (14.9 kg), female(?), coast of Messina, Aug. 8, 1961 (pectoral girdle UMML 11073).

Spec. Med-22: 1,455 mm, 30.8 pounds (14.0 kg), male, Zona Scaletta, south of Messina, Aug. 8, 1961.

Spec. Med-23: 1,370 mm, 21.3 pounds (9.7 kg) (gutted), sex unknown, coast of Calabria.

Spec. Med-24: 1,410 mm, 23.1 pounds (10.5 kg), female, Strait of Messina, Aug. 3, 1961.

Spec. Med-25: 1,375 mm, 24.2 pounds (11.0 kg), female, Strait of Messina, Aug. 7, 1961.

Spec. Med-26: 1,460 mm, 29.7 pounds (13.5 kg), female, Strait of Messina, Aug. 3, 1961.

Spec. Med-27: 1,326 mm, 22 pounds (10.0 kg), female, Strait of Messina, Aug. 2-3, 1961.

Spec. Med-28: 1,307 mm, 20.9 pounds (9.5 kg), female, Strait of Messina, Aug. 1961.

Spec. Med-29: 1,390 mm, 26 pounds (11.8 kg), male, Strait of Messina, Aug. 2-3, 1961.

Spec. Med-30: 1,382 mm, 22 pounds (10.0 kg), male, Torre Faro, Aug.-Sept. 1961.

Spec. Med-31: Mutilated, 20.5 pounds (9.3 kg), male, Contesse, 500 m off coast, Aug.-Sept. 1961.

Spec. Med-32: 1,493 mm, 50.6 pounds (23.0 kg), sex unknown, Strait of Messina, Aug. 4, 1961.

Spec. Med-33: 1,640 mm, 44.4 pounds (20.1 kg), female, Strait of Messina, Aug. 4, 1961 (pectoral girdle UMML 11071).

Spec. Med-34: 1,360 mm, 26.4 pounds (12.0 kg), male, Strait of Messina, Aug.-Sept. 1961.

Spec. Med-35: 1,868 mm, 81.4 pounds (36.9 kg), sex unknown, Sicily, Aug.-Sept. 1961.

Spec. Med-36: UMML 11056, 766 mm, juvenile, vicinity of Malta, between Oct. 1 and 13, 1961, Joe Barbara.

In addition, Walter A. Starck, II, has measured and photographed three mounted specimens at the Museo Civico de Storia Naturale di Genova in September 1960, through the courtesy of Enrico Tortonese. All represent *T. belone*, however, and were so identified by Tortonese (1940:175).

Museo Civico Storia Naturale, 18388, 1,773 mm, Laigueglia, West Riviera, Bay of Genoa, Italy, May 18, 1924.

Museo Civico Storia Naturale, uncat., 1,522 mm, no data.

Museo Civico Storia Naturale, uncat., 1,605 mm, no data.

1.22 Taxonomic status

This species is clearly distinguishable from all other species of *Tetrapturus* in the juvenile and adult stage, based upon the research of Robins and de Sylva (1963). Meristic and morphometric data given in that paper clearly permit separation from all other species. *Tetrapturus georgei* Lowe is discussed by Robins (1974a), and is separable from *T. belone* based on characters discussed by Robins.

1.23 Subspecies

No subspecies are recognized.

1.24 Standard common names and vernacular names

In English-speaking countries this species is frequently referred to as the Mediterranean spearfish (Robins and de Sylva, 1963:89). In Italy and Monaco it is known as aguglia imperiale, as aguglia pelerana in Sicily, pastardella in Malta, iglan and iglokljun in Yugoslavia, auggia imbriale in Algeria, bu mkhiat in Morocco, and marlín in Spain. The Japanese have given the common name chichukai-furai, which they distinguish from all other istiophorids. Bini (1968:60, 62) gives regional variation of Italian spelling and pronunciation as the following: agugghia 'mpiriali (Messina and Catania, Sicily); aguggha imperiale, ugghia 'mpiriali (Reggio de Calabria and Crotone); uguglia imperiali (Manfredonia); auglia 'mperiale (Naples); acura 'mperiale (Gallipoli).

1.3 Morphology

The external morphology of the species is described by Rafinesque (1810) and Robins and de Sylva (1963:92-95), and given earlier in the present paper under Diagnosis. Valenciennes (1831:280-286, pls. 227-228) supplied detailed information on the gross morphology and skeletal structure. Characterization of the larval and juvenile stages is discussed here under the sections on "Reproduction" (3.1).

2 DISTRIBUTION

2.1 Total Area

The Mediterranean spearfish, *T. belone*, is known only from the Mediterranean Sea including the Ligurian, Adriatic, and Ionian seas (Robins and de Sylva, 1963:96-97). It is the most common istiophorid in the central basin around Italy.

Tetrapturus belone is recorded with reasonable certainty from Sicily (Rafinesque, 1810:54-55, Fig. 1), Messina (Cocco, 1884:373; Spartà, 1953, 1961, Calliere, 1962), the Gulf of Naples (Padoa, 1956: 513-51 Palermo (Gigliogli, 1880), Taranto (Costa, 1850: Canestrini, 1872:12), Venice (Ninni, 1912:271), Ligurian Sea (Parona, 1898:368; Tortonese and Tro 1949:134), Malta (Gulia, 1861, 1871), Mallorca (B celó y Combis, 1868:388), the Adriatic (Trois, 18 643-645), and Split, Yugoslavia (Kolombatvić, 18 Šoljan, 1948:376, 389).

Although Robins and de Sylva (1963:96) indicat that it has not been taken from the eastern Mediterranean, Ben-Tuvia (1953) reported upon a juven billfish (356 mm) from off Haifa, Israel, which identified as *Istiophorus gladius* (Broussonnett). T present writer subsequently examined the specim and identified it as the young of *T. belone*, which w later reported by Ben-Tuvia (1966:271-272). A seco juvenile specimen (597 mm SL) was collected by t Woods Hole Oceanographic Institution's vessel *Chain* on 2 October 1966. This specimen (de Sylva a Ueyanagi)[3] was taken by night light and dip net Lebanon, lat. 34°20.5′N, long. 34°41.0′E, and was o of two reported seen under the light (F. J. Mather Southeast Fisheries Center, National Mari Fisheries Service, NOAA, Miami, FL 33149, pe commun.) However, no adults have been report east of the Ionian Sea. Because of the difficulty identification of juvenile and adult billfish, it is poss ble that the species is more widely distributed in t Mediterranean than has been reported in t literature and that spearfish may have been identifi by sport and commercial fishermen as white marli *Tetrapturus albidus* Poey, which occur in the weste and central Mediterranean (Rodríguez-Roda a Howard, 1962; de Sylva, 1973).

References to *T. belone* outside of the Mediterranean Sea cannot be verified (Robins and de Sylv 1963:96-97) and are probably based upon *T. albid* Poey or *T. georgei* Lowe. For a discussion of the latte

[3]de Sylva, D. P., and S. Ueyanagi. Systematics, developme and distribution of the Atlantic species of the fam Istiophoridae. Manuscript in preparation. Rosenstiel School Marine and Atmospheric Science, University of Miami, Miami, 33149.

poorly known species, see Robins (1974b). Nakamura, wai, and Matsubara (1968) refer to spearfish being aken "from the northeast and north-central Atlanic." However, their illustrations (Fig. 18) do not ppear to be of *T. belone* or of *T. pfluegeri*; they may epresent *T. georgei* or even an undescribed species.

Tetrapturus belone belongs ecologically to the chthyofauna of the oceanic epipelagic (Parin, 1968). t evidently completes its entire life history in the pen sea. However, because of the steepness of the ontinental shelf and the proximity to shore of oceanic vaters in the Mediterranean, this spearfish could ome very close to land.

Vertical distribution is unknown. It is usually taken t the surface by harpoon and several types of nets. In he Gulf of Castellammare (Sicily) and near the towns f Torretta Granitola and Marinella, a few are caught n tuna traps (Robins and de Sylva, 1963:96). It is ocasionally caught by using flag lines and drifted andlines. Like other billfish, it probably feeds in the pper 200 m (epipelagic), generally above and within he thermocline.

2.2 Differential Distribution

2.21 Spawn, larvae, and juveniles

Spawning areas are unknown. Lo Bianco (1903, 909), Spartà (1953, 1961), Padoa (1956), and avaliere (1962) reported larval stages of *T. belone* rom the Strait of Messina.

The larvae reported by Lo Bianco (1903, 1909) cannot be identified with certainty, in spite of the statement by Robins and de Sylva (1963:95-96), which notes that one larva (illustrated by Padoa, 1956:514-16, pl. 36, fig. 7) "is clearly an istiophorid and, . . . robably . . . *T. belone*." It seems probable that the arvae depicted by Spartà (1961) may well be those of *T. belone* because the adults of other istiophorids are ot common near the Strait of Messina.

As mentioned earlier, two juveniles (356 mm and 597 mm) were taken from Haifa, Israel, and off Lebanon, respectively. These represent the only two early juvenile specimens of *T. belone* which can definitely be identified. Rafinesque's (1810) type specimen, a juvenile, was from Sicily but this does not ndicate that the specimen was spawned nearby. A uvenile of 37 mm was taken in the Strait of Messina along the beach at Ganzirri (Cavaliere, 1962:Fig. 2), which agrees with the 597-mm specimen collected rom Lebanon except for the lower dorsal fin and the ack of spots on the anterior rays. A juvenile billfish (BMNH) from Malta is figured by de Sylva and Jeyanagi (see footnote 3); however, is it unlikely that his is the young of *T. belone* because of the peculiar, extensive markings on the dorsal fin, which are imited to a few spots on the anterior rays in the larger uveniles of *T. belone* from Israel and Lebanon. Thus, nursery grounds for juveniles of *T. belone* are known nly from the eastern Mediterranean Sea.

2.22 Distribution of adults

Adults of *T. belone* are most common in the central Mediterranean about the Strait of Messina. Such inferences are based solely upon the catch of commercial fishermen, usually using harpoons, who are concentrated in this area. However, commercial fishing using the same method occurs throughout the Mediterranean for broadbill swordfish, *Xiphias gladius* Linnaeus, and it could be expected that *T. belone* would be collected elsewhere if it were common. There are no data on seasonal and annual variations in abundance or availability. Spartà (1961) noted that spearfish occurred in the Strait of Messina in August and September, and were rare in October and November, with some examples being taken during the winter. He noted that they preferred the upper waters of the Strait of Messina, which may well be associated with upwelling and the consequent concentration of food in these upper waters.

2.3 Determinants of Distribution Changes

As an inhabitant of the oceanic epipelagic, *T. belone* is a stenotopic species whose habitat at all stages of its life history is characterized by narrow geographical and seasonal variation in temperature, salinity, and other physicochemical parameters. Like other billfishes, it is typically a clear-water species requiring high-transparency waters for its feeding (which is largely visual. Robins and de Sylva (1963:97-98) postulated that the hypersaline waters and associated changes in temperature and dissolved oxygen of the eastern Mediterranean, especially the Levantine Basin, probably were influential in excluding *T. belone* from the eastern Mediterranean. This may be true for adults, although it does not apply to juveniles inasmuch as three are now reported herein from Israel and Lebanon.

Its food habits are poorly known. Probably it feeds upon clupeoids and sauries, whose distribution is closely determined by oceanographic conditions. Thus, changes, either natural or man-made, should be expected to affect the distribution of *T. belone*. Organisms concentrated at or just above the thermocline (de Sylva, 1962; de Sylva and Davis, 1963) should be expected to concentrate *T. belone* for feeding purposes (Hela and Laevastu, 1970).

Presumably, drastic changes in the habitat, such as from local freshwater runoff or pollutants, would affect the distribution of this stenotopic species at all stages of its life history.

2.4 Hybridization

No natural hybrids of this species are known. Popular speculation contends that all billfish hybridize, but there is no scientific evidence to substantiate this.

3 BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

As far as known, all billfishes, including *T. belone*, are dioecious. Fat deposits adhering to the ovaries of billfishes during their vegetative period are frequently mistaken for testes by fishermen, and hence it is popularly believed that some billfishes are hermaphroditic.

Sexual dimorphism is marked in the related istiophorid genus *Makaira* (*sensu stricto*), with the females growing to a much larger size (de Sylva, 1963a). Nakamura (1949) noted that the Pacific *T. angustirostris* did not exhibit sexual dimorphism, there being no difference in the mean weight of males and females.

Robins and de Sylva (1963:96) found that their data on the mean weight of males and females of *T. belone* were inadequate. No difference in mean size of the sexes was found in the related *T. albidus*, according to de Sylva and Davis (1963). There is no evidence that there are sexual differences in color or morphometric characteristics in *T. belone*.

3.12 Maturity

Nothing is known regarding the size or weight at which sexual maturity is reached in *T. belone*.

3.13 Mating

The spawning act has not been observed, nor is anything known about the location of spawning. Rafinesque (1810) commented that the fish usually travel in pairs, a male and a female together, and that very often both are captured together.

3.14 Fertilization

Reproductive behavior or fecundity have not been studied. Undoubtedly, fertilization is external. Spartà (1953, 1961) has described the pelagic eggs, from the plankton, of *T. belone*.

3.15 Gonads

Robins and de Sylva (1963:96) noted that the sex of 27 specimens of *T. belone* was difficult to determine. This may be because they were examined in August, which should represent a refractory period when the gonads are not well developed, if the hypothesis of Spartà (1953) is correct that spawning occurs from spring to summer. That Spartà (1961) found eggs in the plankton in May would further suggest that spawning is long complete by August.

The gonads of 27 fish examined from Sicily were asymmetrical, λ-shaped (the left side longer in one female, the right side longer in most others), and v orange (Robins and de Sylva, 1963:96).

3.16 Spawning

Spartà (1953) reported on eggs and larvae from Strait of Messina. He identified eggs from plank taken in May. The identity of the larvae, 4.88 and 5 mm, is questionable (Gehringer, 1956:169; Ueyana 1962:186; Robins and de Sylva, 1963:95). Subseque ly, Spartà (1961) indicated the spawning period spring to summer. The difficulty encountered in id tifying the sex of gonads from 27 specimens of *belone* collected in August from Sicily suggests t spawning did not occur during this period. The currence of juveniles in October (Spartà, 1961:20) a suggests a late-spring spawning period. However, earlier spawning season (i.e., winter months) can be precluded. A spawning season of Decem through February is postulated for *T. pfluegeri* in t western Atlantic by de Sylva and Breder,[4] and winter or spring spawning season might not be an u reasonable hypothesis for *T. belone*. Ueyana 1962:188; Ueyanagi (in Howard and Ueyana 1965:103) concluded that *T. angustirostris* spawn more intensively in winter in the Pacific.

Spawning has not been observed, but larval stag of *T. pfluegeri* have been reported from far offsh (de Sylva and Ueyanagi (see footnote 3)). Oth istiophorids (excluding the neritic *Istiophorus*) appe to spawn far from shoal water, and it is likely that *belone* similarly spawns offshore.

3.2 Pre-Adult Phase

3.21 Embryonic phase

The pelagic eggs of *T. belone* collected from t plankton are described in detail by Spartà (195 1961). The eggs averaged 1.48 mm in diameter and t oil globule was yellow-green. The eggs were incubat for several days and their development is described detail.

3.22 Larval phase

Two larval specimens of *T. belone* collected ne Capri, off Punta Campanella, were described by Bianco (1903:127, 166, 238). Comparisons of the specimens with other istiophorid larvae were made Lo Bianco (1909:755). Larvae from Messina report by Spartà (1953) do not seem to be those istiophorids (Gehringer, 1956:169; Ueyanag 1962:186; Robins and de Sylva, 1963:95-96). Larg specimens (29 and 54 mm) were reported from t coast of Faro, Strait of Messina, apparently throv ashore from the southeast wind (Sirocco). These t

[4]de Sylva, D. P., and P. R. Breder. Gonad histology and rep ductive cycle in the Atlantic species of the Istiophoridae. Ma script in preparation. Rosenstiel School of Marine and At spheric Science, University of Miami, Miami, FL 33149.

pecimens were an extreme rarity, other examples not aving been found in the past 50 yr in spite of diligent search (Spartà, 1961).

3.23 Adolescent phase

A juvenile specimen of *T. belone* (356 mm) was eported (as *Istiophorus gladius* (Broussonet) by Ben-uvia (1953:18), which was subsequently reidentified Ben-Tuvia, 1966) as *T. belone.* A second specimen 97 mm) was collected by the Woods Hole ceanographic Institution's RV *Chain* off Lebanon. he striking characteristic of these specimens is that ey generally resemble the adult except that the first rsal fin is high and saillike, almost as in the genus *tiophorus* (see de Sylva, 1963b:Fig. 2). A high dorsal n in the adolescent phase is common to all tiophorids, and accounts for much misidentification young stages in the literature.

The dorsal fin is unlike that of the young of *tiophorus* in that it lacks diffuse blotches (see ehringer, 1956, 1971; de Sylva, 1963b:Fig. 3). *Is-iophorus* has not been reported from the Mediterra-ean, nor has the blue marlin, *Makaira nigricans* acépède, but both could occur there and should be oked for (de Sylva, 1973). The dorsal fin of the oung of *Makaira* is distinctively convex, reaching its ighest point anteriorly. It has a distinct wavy pattern ot seen in other istiophorids. Young of *Tetrapturus lbidus* are known from North Carolina (de Sylva, 963b: Fig. 1) and off Havana (Anonymous, 1968). hey have five or six distinctive dark ocelli rimmed ith pale areas, near the base of the first dorsal fin. he young of *T. belone* is unique in that the dorsal fin ears several small, dark blotches at the bases of the nterior rays, and at the tips of the first few rays, he rest of the fin being dusky and otherwise un-narked.

Other features of development between the adoles-ent and adult stages are less distinctive. The body of he adolescent is typically thin and slab-sided, which ncreases in robustness in the adult. The bill, long and ender in the adolescent, is negatively allometric, be-g relatively short in the adult. Characteristic of all he spearfishes is the shape of the first dorsal fin in the dult, which has a slight lobe anteriorly, sloping bruptly behind the first several rays, then running ng and high, paralleling the dorsal profile (see obins and de Sylva, 1960:406; 1963:93, Fig. 4).

3.3 Adult Phase

3.31 Longevity

Nothing is known of the rate of growth or maximum ge of *T. belone.* Size-frequency analysis of adults, sed by de Sylva (1957) to estimate age and growth of *tiophorus americanus*, requires a great deal of size ata from individual specimens, which are presently navailable.

3.32 Competitors

Tetrapturus belone is ecologically quite similar to the broadbill swordfish and *T. albidus*, with which it undoubtedly competes for food. The blue shark, *Prionace glauca*, and the bluefin tuna, *Thunnus thynnus*, are other large predators which occur in the same ecological niche and which probably feed upon the same organisms. The broadbill swordfish and the bluefin tuna are routinely caught by commercial fishermen about Sicily using the same harpoon and trap-net techniques by which *T. belone* is taken.

3.33 Predators

Because of the relatively large size attained by *T. belone*, it is probably not preyed upon extensively in the adult phase except by sharks. The blue shark is occasionally common off Sicily and probably could be considered as an important predator, though the blue shark, being somewhat sluggish, would probably be more successful in capturing species not as swift as *T. belone.*

3.34 Parasites, diseases, injuries, and abnormalities

Nothing has been published on parasites and diseases of *T. belone.* A parasitic copepod (*Penella*), common on all istiophorids, has not been reported for *T. belone.* Abnormalities, such as deformed dorsal fins, broken or bent bills, and broken pectoral or caudal rays, are common among other istiophorids, but no such deformities are reported for *T. belone.*

3.4 Nutrition and Growth

3.41 Feeding

Feeding habits of *T. belone* have not been reported in the literature. Based on their food habits, however, they probably are visual feeders of the upper, well-lit layers. It is not known if they feed at night. Antonio Spartà reports (in Robins and de Sylva, 1963) that *T. belone* follows schools of Atlantic sauries, *Scomberesox saurus*, into the Strait of Messina and, hence, in this region they are feeders in the shallower strata of the epipelagic zone. Spartà (1961:20) wrote that *T. belone* preferred the upper waters of the Strait of Messina.

Probably because there is considerable upwelling of food and nutrients into the upper layers of the Strait of Messina, *T. belone* finds an ample food supply in the upper waters so that it does not have to search the deeper strata for food, such as occurs with many other istiophorids. About Messina, fishermen harpoon them during their fishing period from early in the morning (0530-0600) until about 1600. Since night fishing is not carried out here, therefore, specimens have not been studied to disclose if they feed at night. Cavaliere (1962:172) reported that, to the south of the

Strait of Messina, *T. belone* is caught at night with "palamidara," vertical nets used to catch albacore, *Thunnus alalunga*. Possibly *T. belone* is feeding at this time, although no reports on feeding or food habits based on night-caught specimens are forthcoming.

3.42 Food

The food of *T. belone* was discussed by Robins and de Sylva (1963:95), based on specimens caught by fishermen. Since specimens reported on from Sicily were not fresh, and many had been ultimately frozen, thawed, and refrozen, most contained only unrecognizable remains. Many had fish skeletons, which were evidently of Atlantic sauries and sardinelike vertebrae. A few contained the needlefish *Belone belone*. Rafinesque (1810) reported that in autumn off the Sicilian coast, *T. belone* pursues dolphins; pilotfish, *Naucrates*; and flyingfishes.

3.5 Behavior

3.51 Migrations and local movements

Seasonal movements of *T. belone* are based entirely upon their capture by commercial fishermen. Distinct migrations are not documented, nor is it known whether the appearances of *T. belone* in the commercial fishery results in longitudinal or vertical movements to the surface fishery where they can be captured by harpoons and shallow nets.

Adults occur in the Strait of Messina, the area of heavy fishing, in August and September, and occasionally in October and November. Some examples are rarely captured in winter (Spartà, 1953, 1961; Cavaliere, 1962; Bini, 1968).

3.52 Schooling

No mention is made in the literature as to the schooling of *T. belone*, but it seems unlikely because other istiophorids do not school in the strict sense. Rafinesque (1810) and Valenciennes (1831) reported that *T. belone* frequently travels in pairs, this being known to occur in other istiophorids, possibly being a behavioral mechanism for feeding.

4 POPULATION

4.1 Structure

Nothing is known of the sex ratio or age composition of the populations of *T. belone*. Robins and de Sylva (1963:96) reported that of 27 specimens taken off Sicily by commercial fishermen, 19 were males and 8 females.

The usual size composition of the commercial catch of *T. belone* is from 10 to 30 kg (Cavaliere, 1962:172), averaging about 2 m long (Spartà, 1961:20). Specimens of 4 to 5 kg are rarely caught. Maximu size reported is about 70 kg (Rafinesque, 1810).

5 EXPLOITATION

5.1 Fishing Equipment

The fishery for *T. belone* occurs incidentally to th carried out for the swordfish (Spartà, 1961; Cavalier 1962:174). In the Strait of Messina, the major gear the harpoon (fiocini), although *T. belone* is occasio ally captured in nets (ravastina) which are used Atlantic sauries, locally called "costardella," which *T. belone* feeds. To the south *T. belone* is o casionally caught in vertical nets (palamidare) whi are used principally for albacore.

5.2 Fishing Areas

5.21 General geographic distribution

See 5.22.

5.22 Geographic ranges

The fishing area in the Strait of Messina (bo Sicilian and Calabrian coasts) occurs from Bagnar Capo Rasocolmo in the north in the Tyrrhenian Se and from Capo Giardini-Melito di Porto Salvo to t south in the Ionian Sea. In addition to the harpo fishery, a few specimens of *T. belone* are caught traps set for tuna in the Gulf of Castellammare a near the towns of Torretta Granitola and Marinell Little is known specifically about the fishing groun in the remainder of the Mediterranean Sea, and quantitative data are available on the annual seasonal catch in various counties.

Francesca R. LaMonte kindly sent me an Englis translation of a paper by Mazzullo (1906) whi documents the swordfish fishery in the Strait Messina; because this fishery also captures *T. belon* the techniques and gear used in the fishery a noteworthy. Mazzullo gives a general summary of t composition of manpower in a typical fishing oper tion for swordfish off Sicily, including the number a kinds of personnel involved per vessel and the eve tual distribution of fish and profits.

6 PROTECTION AND MANAGEMENT

6.1 Regulatory (Legislative) Measures

No data.

6.2 Control or Alteration of Physical Features the Environment

See 6.3.

6.3 Control or Alteration of Chemical Features of the Environment

The relationship between the community population structure, though presently unknown, and the ecosystem will ultimately depend on man's effects upon the high-seas environment in which *T. belone* lives. Hypersaline conditions resulting from the reduced freshwater flow of the Nile River are increasing in the eastern Mediterranean, from which two juveniles of *T. belone* are reported, so that, if this were to constitute an important nursery ground for the juveniles, high salinity water would be expected to be unfavorable to this stenotypic species.

Man-made pollution, especially from oil and other petrochemicals, sewage pollution, and radioactivity, is potentially detrimental via the food chain, even to an epipelagic species such as *T. belone*. For example, Horn, Teal, and Backus (1970) found lumps of crude oil residue to be widely dispersed in the Mediterranean Sea and the eastern North Atlantic Ocean, stressing that toxic fractions of hydrocarbons have been isolated from these lumps. They found large amounts of tar in the stomachs of Atlantic sauries collected southwest of Sardinia, and concluded that this ingestion of the tar by sauries provides a direct introduction of a material known to be toxic into the oceanic food web." Because *T. belone* feeds mainly upon sauries, there is a possibility that an important part of the food web of *T. belone* could be affected if sauries are eventually killed by the toxic hydrocarbons. Further, concentrations of toxic hydrocarbons which were sublethal to the sauries could be theoretically concentrated within their bodies and eventually in *T. belone* and persons feeding upon them. Other man-made components, especially the chlorinated hydrocarbons, should be expected to occur in the tissues of *T. belone*, and could conveivably affect spawning or other metabolic processes, in addition to becoming concentrated in various tissues eaten by man.

6.4 Control or Alteration of the Biological Features of the Environment

See 6.3.

POND FISH CULTURE

Not applicable.

ACKNOWLEDGMENTS

I am especially grateful to Francesca R. LaMonte who gave me her personal library of billfish literature, including many valuable and accurate translations by her of the foreign literature on billfish.

My colleague, C. Richard Robins, kindly reviewed the manuscript and has always stimulated many discussions on billfish and other ancillary matters. The late John K. Howard and Al Pflueger were instrumental in supplying information, specimens, and contacts which permitted our study of spearfish; the many others who enabled this study are acknowledged by Robins and de Sylva (1960, 1963).

The research for this paper was supported in part by the National Science Foundation, grants NSF GB-893 and NSF G-23745/GN-2676; the U.S. Bureau of Sport Fisheries and Wildlife, contracts BSFW 14-16-0008-775/DI-14-16-0008-957; the Sport Fishing Institute, Washington, D.C.; the Port Aransas Rod and Reel Club; and the Maytag Chair of Ichthyology, University of Miami.

LITERATURE CITED

AGASSIZ, L.
1843. Poissons cycloides. Recherches sur les poissons fossiles Neuchâtel. Vol. 5, 160 p.

ANONYMOUS.
1968. Relentless sea relinquishes rarity to resourceful researchers. Fla. Conserv. News 3(11):5. [Photograph and article on a young white marlin.]

BARCELÓ Y C., F.
1868. Catálogo metodico de los peces que habitan ó frecuentan las costas de las Islas Baleares. Rev. Prog. Cienc. Madred 18(3/4):219-221, 224-234, 299-315, 385-395.

BEN-TUVIA, A.
1953. Mediterranean fishes of Israel. Bull. Isr. Sea Fish. Res. Stn. Haifa 8:1-40.
1966. Red Sea fishes recently found in the Mediterranean. Copeia 1966:254-275.

BINI, G.
1968. Atlante dei pesci delle coste Italiane. Vol. 6. Mondo Sommerso, Editrice, Milano, 177 p.

BONAPARTE, C. L. P.
1832-1841. Iconografia della fauna Italica per le quattro classi degli animali vertebrati. Vol. III. Roma, Salviucci, n.p.

BRIGGS, J. C.
1958. A list of Florida fishes and their distribution. Bull. Fla. State Mus., Biol. Sci. 2:223-318.

CANESTRINI, G.
1872. Pesci. *In* E. Cornalia, Fauna d'Italia, Part 3. Vallardi, Milano, 208 p.

CAVALIERE, A.
1962. Notizie su biologia e pesca de *Tetrapturus belone* Raf., cenni sull'adulto e descrizione di un suo rarissimo stadio giovanile. Boll. Pesca Piscic. Idrobiol., New Ser. 15(2):171-176.

COCCO, A.
1884. Indice ittiologico del mare di Messina. Nat. Sicil. 5:35-40.

COSTA, O.
1850. Fauna del regno di Napoli . . . Pesci. Naples, Azzolino, Part 1(49):10-11.

CUVIER, G., and A. VALENCIENNES.
1831. Histoire de naturelle poissons. Paris, 8:1-509.

DE SYLVA, D. P.
1957. Studies on the age and growth of the Atlantic sailfish, *Istiophorus americanus* (Cuvier), using length-frequency curves. Bull. Mar. Sci. Gulf Caribb. 7:1-20.
1962. Red-water blooms off northern Chile, April-May 1956, with reference to the ecology of the swordfish and the striped marlin. Pac. Sci. 16:271-279.
1963a. Preliminary report on the blue marlin sport fishery of Port Antonio, Jamaica. Univ. Miami, Inst. Mar. Sci., 15 p.
1963b. Postlarva of the white marlin, *Tetrapturus albidus*, from the Florida Current off the Carolinas. Bull. Mar. Sci. Gulf Caribb. 13:123-132.
1973. Istiophoridae. *In* J. D. Hureau and T. Monod (editors), Clofnam (Check-list of the fishes of the north-eastern Atlantic and of the Mediterranean) 1:477-481. UNESCO, Paris.

DE SYLVA, D. P., and W. P. DAVIS.
1963. White marlin, *Tetrapturus albidus*, in the middle Atlantic bight, with observations on the hydrography of the fishing grounds. Copeia 1963:81-99.

GEHRINGER, J. W.
1956. Observations on the development of the Atlantic sailfish, *Istiophorus americanus* (Cuvier), with notes on an unidentified species of istiophorid. U.S. Fish Wildl. Serv., Fish. Bull. 57:139-171.
1970. Young of the Atlantic sailfish, *Istiophorus platypterus*. U.S. Fish Wildl. Service., Fish. Bull. 68:177-189.

GIGLIOGLI, E. H.
1880. Elenco . . . dei pesci italiani. Pisces, Florence, p. 18-55.

GOODE, G. B.
1883. Materials for a history of the sword-fish. U.S. Comm. Fish and Fish. Rep. Comm. 1880, Part 8:289-394.

GULIA, G.
1861. Faunae Melitensis Quarta Classis Animalium Vertebratorum. Pisces. *In* Tentamen Ichthyologiae Melitensis, Malta, p. 9-34.
1871. L'aguglia imperiale. II Barth, Malta.

GÜNTHER, A.
1860. Catalogue of the acanthopterygian fishes in the collection of the British Museum. London 2, 548 p.

HELA, I., and T. LAEVASTU.
1970. Fisheries hydrography. How oceanography and meteorology can and do serve fisheries. Fish. News (Books) Ltd. Lond., 137 p.

HORN, M. H., J. M. TEAL, and R. H. BACKUS.
1970. Petroleum lumps on the surface of the sea. Science (Wash., D.C.) 168:245-246.

HOWARD, J. K., and S. UEYANAGI.
1965. Distribution and relative abundance of billfishes (*Istiophoridae*) of the Pacific Ocean. Univ. Miami, Inst. Mar. Sci., Stud. Trop. Oceanogr. 2, 134 p.

KOLOMBATOVIĆ, J.
1886. Imenik kraljesnka Dalmacije. Spalato, 12 p.

LAMONTE, F. R., and D. E. MARCY.
1941. Swordfish, sailfish, marlin and spearfish. Ichthyol. Contrib. Int. Game Fish Assoc. 1(2):1-24.

LO BIANCO, S.
1903. Le pesche abissali eseguite de F. A. Krupp col yacht Puritan nelle adiacenze di Capri ed in altre località del Mediterraneo. Mitt. Zool. Stn. Neapel 16:109-279.
1909. Notizie biologiche reguardanti specialmente il periodo di maturità sessuale degli animali del golfo di Napoli. [In Ital.] Mitt. Zool. Stn. Neapel 19:513, 760, 761.

LÜTKEN, C. G.
1876. Sur les histiophores à bec arrondi, en particulier l'*H*... *tiophorus orientalis*. J. Zool. (Gervais) 5:60-63.

MAZZULLO, L.
1906. Monograph on the [commercial] swordfishing in ... Straits of Messina. Chamber of Commerce and Arts ... Messina, Rosario Alico, Messina, 18 p. [Engl. translated ... F. R. LaMonte.]

NAKAMURA, H.
1949. The tunas and their fisheries. [In Jap.] Takeu... Shobō, Tokyo, 118 p. [Translated by W. G. Van Camp... 1952, U.S. Dep. Int., Fish Wildl. Serv., Spec. Sci. Rep. Fi... 82, 115 p.]

NAKAMURA, I., T. IWAI, and K. MATSUBARA.
1968. A review of the sailfish, spearfish, marlin, and swo... fish of the world. [In Jap.] Kyoto Univ., Misaki M... Biol. Inst., Spec. Rep. 4, 95 p.

NARDO, G. D.
1833. De skeponopodo novo piscium, genere el du Guebu... marcgravii specii, ill cognata. Isis (Oken) 26:415-419.

NINNI, E.
1912. Catalogo dei pesci del Mare Adriatico, Venice, 271...

PADOA, E.
1956. Monographia: Uova, larvae e stadi giovanili di Tel... stei. Divisione: Scombriformes (Eggs, larvae and juver... stages of the Scombriformes (in part) of the Gulf of Naple... [In Ital.] Fauna Flora Golfo Napoli 38:471-547. [Tra... lated by J. P. Wise and G. M. Ranallo, 1967, Transl. No.... of the Trop. Atl. Biol. Lab., 49 p.; avail. at Bur. Comm... Fish. (Natl. Mar. Serv.), Miami, Fla.]

PARIN, N. V.
1968. Ikhtiofauna okeanoskoi epipelagiali (Ichthyofauna ... the epipelagic zone). [In Russ.] Akad. Nauk. SSSR. ... Inst. Okeanol., Moscow, 186 p. [Translated by M. Rav... Israel Program Sci. Transl. 1970, 206 p.; available Natl. Te... Inf. Serv., Springfield, VA 22151 as TT 69-5902].

PARONA, C.
1898. La pesca marittima in Liguria. Atti Soc. Ligusti... Sci. Nat. Geogr., Genova 9:327-393.

RAFINESQUE, C. S.
1810. Caratteri di alcuni nuovi generi e nuove specie de a... mali e piante della Sicilia, con varie osservazioni sopr... medesimi. Palermo, 105 p.

ROBINS, C. R.
1974. The validity and status of the roundscale spearfis... *Tetrapturus georgei*. *In* R. S. Shomura and F. Willia... (editors), Proceedings of the International Billfish Symp... sium, Kailua-Kona, Hawaii, 9-12 August 1972. Part 2. R... view and Contributed Papers, p. 54-61. U.S. Dep. Co... mer., NOAA Tech. Rep. NMFS SSRF-675.
1975. Synopsis of biological data on the longbill spearfis... *Tetrapturus pfluegeri*, Robins and de Sylva. *In* R. S. Sh... mura and F. Williams (editors), Proceedings of the Intern... tional Billfish Symposium, Kailua-Kona, Hawaii, 9-... August 1972. Part 3. Species Synopses, p. 28-38. U.S. De... Commer., NOAA Tech. Rep. NMFS SSRF-675.

ROBINS, C. R., and D. P. DE SYLVA.
1960. Description and relationships of the longbill spearfis... *Tetrapturus belone*, based on western North Atlantic spe... mens. Bull. Mar. Sci. Gulf Caribb. 10:383-413.
1963. A new western Atlantic spearfish, *Tetrapturus pflu... geri*, with a redescription of the Mediterranean spearfi... *Tetrapturus belone*. Bull. Mar. Sci. Gulf Caribb. 13:8... 122.

RODRÍGUEZ-RODA, J., and J. K. HOWARD.
1962. Presence of Istiophoridae along the South Atlantic and Mediterranean coasts of Spain. Nature (Lond.) 196:495-496.
SOLJAN, T.
1948. Fauna i flora jadrana. 1. Ribe. Inst. Oceanogr. Ribarst. jugoslavia. Zagreb, Hrvatske, 437 p.
SPARTÀ, A.
1953. Uova e larve di *Tetrapturus belone* Raf. (aguglia imperiale). Boll. Pesca Piscic. Idrobiol. 8(1):58-62.
1961. Biologia e pesca di *Tetrapturus belone* Raf. e sue forme post-larvali. Boll. Pesca Piscic. Idrobiol. 15(1):20-24.
TORTONESE, E.
1940. Sugli scomberoidei mediterranei del genere *Tetrapturus*. Boll. Mus. Zool. Anat. Comp. Torino 48:173-178.
1958. Elenco dei leptocardi, ciclostomi, pesci cartilaginei ed ossei del Mare Mediterraneo. Atti Soc. Ital. Sci. Nat. 97: 309-345.
TORTONESE, E., and L. TROTTI.
1949. Catalogo dei pesci del mar Ligure. Atti Accad. Ligure 6:49-164.
TROIS, E. F.
1880. Annotazione sopra gli avanzi di un *Tetrapturus belone* preso nell'Adriatico. Atti Istit. Sci. Venezia, 1879-1880, Ser. 5, 6:643-645.
UEYANAGI, S.
1962. On the larvae of the shortnosed spearfish, *Tetrapturus angustirostris* Tanaka. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 16:173-189.
VÉRANY, J. B.
1847. Aggiunto al catalogo de' pesci della Liguria. Atti Ottava Riunione Sci. Ital. Genoa, 1846:492-494.

Synopsis of Biological Data on Striped Marlin, *Tetrapturus audax* (Philippi), 1887

SHOJI UEYANAGI[1] and PAUL G. WARES[2]

1 IDENTITY

1.1 Nomenclature

1.11 Valid Name

Tetrapturus audax (Philippi), 1887 is the name adopted by the most recent review of the family (Nakamura, Iwai, and Matsubara, 1968). The original combination was *Histiophorus audax* Philippi, 1887. Anal. Univ. Chile 71:34-39.

1.12 Objective synonymy

All synonyms are assumed to be subjective without consulting original papers and are listed under section 1.21.

1.2 Taxonomy

1.21 Affinities

Suprageneric

Phylum Chordata
Subphylum Vertebrata
Superclass Gnathostomata
Class Osteichthyes
Subclass Actinopterygii
Order Perciformes
Suborder Xiphioidei
Family Istiophoridae

Generic

Genus *Tetrapturus* Rafinesque, 1810.

Caratteri di alcuni nuovi generi e nuove specie di animali [principalmente di pesci] e piante della Sicilia, con varie osservazioni sopra i medisimi. Palermo, 105 p.

Type-species *Tetrapturus belone* Rafinesque, 1810.

Robins and de Sylva (1960, 1963) placed the striped marlin in *Tetrapturus*, following the works on hierarchical classification of the family by Hirasaka and Nakamura (1947) and Nakamura (1949). The placement of *audax* in *Tetrapturus* is supported by Ueyanagi (1963b), Howard and Ueyanagi (1965), and Nakamura et al. (1968).

We follow the generic concept of Nakamura et al. (1968), who described the genus as follows:

> The height of the dorsal fin is greater than the body depth. The ventral fin rays are rather long, the fin membrane not well developed. The body is compressed (flat) and except for the striped marlin (Makajiki) and the white marlin (Nishimakajiki), extends in a straight line from the pre-occular area to the base of the dorsal fin. The cranium is long and narrow. The neural and haemal spines of the central vertebrae form a parallelogram. There are 24 vertebrae (12 + 12 = 24). The lateral appophysis[3] is not well developed.

These authors include the following species in the genus: *T. angustirostris* Tanaka, 1914; *T. belone* Rafinesque, 1810; *T. pfluegeri* Robins and de Sylva, 1963; *T. albidus* Poey, 1861; *T. audax* (Philippi, 1887).

Specific

Identity of type specimen:

Species *T. audax* (Philippi, 1887).

Type specimen: Apparently one of the two deposited in the Museo Nacional de Historia Natural, Santiago, Chile by Rudolfo A. Philippi.

Type Locality: Iquique, Chile.

Diagnosis: Ventral fins and two caudal keels are present; snout cross section is nearly circular; first dorsal fin anteriorly is about same height as body depth or greater[4] and is not saillike but slopes abruptly

[1]Fisheries Agency of Japan, Far Seas Fisheries Research Laboratory, Shimizu 424, Japan.

[2]Southwest Fisheries Center, National Marine Fisheries Service, NOAA, Tiburon, CA 94920; present address: Northwest Fisheries Program, U.S. Fish and Wildlife Service, 495 Tyee Drive, Tumwater, WA 98502.

[3]See Ueyanagi and Watanabe (1965) for usage of this term.

[4]Due to allometric growth of the dorsal fin relative to body depth the fin height may be less than body depth in fish larger than 36 cm FL (Royce, 1957).

osteriorly, the middle rays being much shorter than ne anterior; snout is fairly long; vent is located very ose in front of first anal fin; pectoral fins are fairly road and long and fold against body; and the tips of ectorals, first dorsal, and first anal fins are pointed.

ubjective synonymy:

listiophorus audax	Philippi, 1887
stiophorus audax	Delfin, 1901
Tetrapterus mitsukurii	Jordan and Snyder, 1901
Tetrapturus ectenes	Jordan and Evermann, 1926
Makaira audax	Jordan and Evermann, 1926
Makaira grammatica	Jordan and Evermann, 1926
Makaira holei	Jordan and Evermann, 1926
Makaira mitsukurii	Jordan and Evermann, 1926
Makaira zelandica	Jordan and Evermann, 1926
Marlina mitsukurii	Grey, 1928
Marlina zelandica	Whitley, 1937
Kajikia mitsukurii	Hirasaka and Nakamura, 1947
Kajikia formosana	Hiraska and Nakamura, 1947
Tetrapturus tenuirostratus	Deraniyagala, 1951
Tetrapturus acutirostratus	Deraniyagala, 1952
Makaira formosana	Matsubara, 1955
Tetrapturus brevirostris	Munro, 1955
Marlina audax	Smith, 1956a, b
Marlina jauffreti	Smith, 1956b
Tetrapturus audax	Robins and de Sylva, 1960
Makaira jauffreti	Jones and Silas, 1964

rtificial key to genus (Nakamura et al., 1968):

A. Anterior fin rays of first dorsal fin fairly high, posterior rays about same height; vent situated decidedly anterior to origin of the first anal fin; second anal fin anterior to second dorsal fin.

B. Pectoral fin narrow and short

C. Snout very short. . . Shortbill spearfish (Furaikajiki) *T. angustirostris* Tanaka.

CC. Snout fairly long . . . . . Mediterranean spearfish (Chichukaifurai). *T. belone* Rafinesque.

BB. Pectoral fin wide and long . . . . . Longbill spearfish (Kuchinagufurai) *T. pfluegeri* Robins and de Sylva.

AA. Height of anterior portion of first dorsal fin about same as the body depth but gradually decreasing in height posteriorly; vent directly anterior to the origin of the first anal fin; second dorsal fin and second anal fin in parallel positions.

D. Pectoral fin wide and its tip rounded. The tip of the first dorsal fin and first anal fin rounded . . . . . White marlin (Nishimakajiki) *T. albidus* Poey.

DD. Pectoral fin narrow, and its tip pointed; the tips of the first dorsal fin and first anal fin pointed . . . . . Striped marlin (Makajiki) *T. audax* (Philippi).

1.22 Taxonomic status

The species is established on the basis of morphology without breeding data.

The species may be polytypic (see 1.31 below).

1.23 Subspecies

No subspecies are recognized.

1.24 Standard common names and vernacular names

Country	*Standard common name*	*Vernacular name(s)*
Sri Lanka		Seraman Koppara
Chile		Pez aguja
China		Chi zuo fo yii
Japan	Makajiki	Maka, Kajiki, Kajikimaguro, Nairagi, Nairage, Nairanbo
Kenya		Nduaro
Mexico	Marliń rayado	Marliń, agujon, pez puereo
New Zealand	Striped marlin	New Zealand marlin
Philippines		Spearfish
Russia		Polosatii marlin
Taiwan	Hung ju chi yii	Hung ju ting pan
United States	Striped marlin	Pacific striped marlin, barred marlin, Pacific marlin, striped swordfish, spearfish, spikefish
Vietnam		Ca co Mitsukurii

1.3 Morphology

1.31 External morphology (for description of spawn, larvae, and adolescents, see 3.17, 3.22, 3.23).

Generalized: Gregory and Conrad (1939) provide a scale diagram of the striped marlin outline based on modal body proportions of 30 specimens from New Zealand and Australia using standard length as the basic body measurement. Thirty-eight measurements were made on each specimen and are published in absolute and as percentage of standard length along with total body weight.

Nakamura et al. (1968) provide the following description:

External Characters. The first dorsal fin has 37-42 fin rays. The second dorsal fin has 6 soft rays. The first anal fin has 13-18 fin rays. The second anal fin has 5-6 soft rays. The pectoral fin has 18-22 soft rays. The ventral fin has one spine and 2 soft rays. The

body is elongated (the body length is about 5.9-7.3 times the body depth) and is rather thick. The snout is long (the head length is about 0.88-0.99 times the length of the maxillary), and its cross-section is almost round. The body is densely covered with scales; the tips of the scales are pointed. Scales from specimens less than 1 meter in length do not have this species characteristic. There are small file-shaped teeth on both jaws and on the palate. The lateral lines on the sides curve over the pectoral fin and then continue in a straight line to the area of the caudal fin. The head is large (the body length is 3.6-3.8 times the length of the head). The eyes are moderately large. There is a relatively conspicuous crest on the outer edge of the head between the pre-occular region and the origin of the first dorsal fin. There are 2 scutes on each side of the tail near the caudal peduncle; the tail is strong and deeply forked. The pectoral fin, has a pointed tip and is located rather low on the body; it is shorter than the head (the head length is about 1.14-1.99 times the length of the pectoral fin). The first dorsal fin begins above the posterior end of the pre-opercle bone; its first few rays are larger than the body depth, but as it progresses towards the back, it gradually becomes shorter, ending just in front of the origin of the second dorsal fin. The tip of the first anal fin is pointed, large and sickle-shaped. The second dorsal fin and the second anal fin are about the same size and shape. The latter is located a little further forward on the body than the former. In spite of the fact that the ventral fin is longer than the pectoral fin in smaller specimens, the opposite is true for larger specimens. The fin membrane of the first dorsal fin is dark blue. The back of the body is dark blue with splotches of black on it; towards the ventral side of the body, 10 or more rows of cobalt-colored stripes are clearly visible. The other fins are blackish brown, or sometimes a dark blue. The bases of the first and second anal fins are silvery-white.

Morrow (1952a) published morphometric data on 49 specimens from New Zealand and later (Morrow, 1957) published extensive morphometric data and anal ray counts on 39 fish from Peru. These data include standard length as the basic body length measurement. Ueyanagi (1957b) presented morphometric data on young specimens, 80- to 180-cm eye-fork length, from the western North Pacific. Royce (1957) reported extensive morphometric data of 25 specimens from the central Pacific using fork length as the basic measurement but also giving standard length and eye-fork lengths for some specimens. He also published more limited data on 30 specimens measured by the Hawaii Division of Fish and Game. Kamimura and Honma (1958) published morphometric data on five characters using eye-fork length as the basic body measure for 56 fish south of the equator and 124 fish north of the equator in the western Pacific. Williams (1967) presented dorsal and anal fin ray counts of 13 specimens from East Africa. Merrett (1971) gives fin measurements on about 23 other specimens.

Counts have been given by several authors and are shown in Table 1.

Georgraphic variation: Geographic variation appears to be considerable. Morrow (1957) concluded that striped marlin from Peru and northern New Zealand represented separate and distinct p ulations based on significant differences in morphometric and meristic characters as follo average absolute lengths of pelvic fins, counts spines and rays in the first anal fin, and regressions of the following measurements on st dard length: greatest body depth, length of bas second dorsal fin, length of base of first anal width of base of pectoral fin, snout tip to origin of f dorsal fin, snout tip to origin of second dorsal snout tip to origin of first anal fin, snout tip to p terior edge of operculum, snout tip to posterior of maxillary. By a character index (CI) in wh

$$CI = \frac{\text{pelvic length}}{10} - \frac{100}{SL} \text{ (Depth + Len}$$

of anal base + Width of pectoral base), Morr could separate correctly about 72% of the 69 spe mens from which the index was derived. The N Zealand specimens tended to have character indi of considerably lower numerical value than the P specimens.

In the western Pacific (west of long. 170° Kamimura and Honma (1958) found a remarka difference in the lengths of the pectoral fins betwe northern (lat. 30°-35°N) and southern (lat. 18°-25° striped marlin. Covariance analysis of regression pectoral fin on eye-fork length showed no significa difference in slope of regression but a highly signi cant (0.01) difference in adjusted means. Also signi cant differences (0.05) were found for both regressi coefficients and adjusted means for regressions of ey to insertion of second dorsal on eye-fork length. Fr these differences these authors concluded northe and southern populations in the western Pacific we extremely separated. In intermediate waters of t northern hemisphere (lat. 5°-25°N) all but three fi had pectoral lengths clustered about the regressi line for the northern population. The pectoral lengt of the other three fish, which were taken from lat. to 15°N, were close to the regression line for t southern population and were presumed to ha strayed from that population.

Honma and Kamimura (1958) supported t hypothesis of separate north and south populations the western Pacific with the following observation

a) a zone of low hook-rate along the equator sep rates the populations;
b) the main spawning grounds are widely separat and spawning seasons are a half-year apar
c) the maximum size attained is much larger in t southern population;
d) adaptations of the two populations to enviro mental circumstances do not coincide in d tails, differing with growth stages.

Howard and Ueyanagi (1965) extended t

Table 1.—Fin-ray counts of *Tetrapturus audax*.

Area and author	First dorsal fin	Second dorsal fin	First anal fin	Second anal fin	Pelvic fins	Pectoral fins
Chile						
Philippi, 1887	38	5	12	5	1	
Western Pacific						
Nakamura, 1938, 1949	37-43 (III+12–15+XXII–XXV)	6	14-15 (II+12–13[1])	6	1+2	
Western Pacific						
Ueyanagi, 1959	37-44					
New Zealand						
Morrow, 1957			12-16			
Peru						
Morrow, 1957			14-17			
East Africa						
Williams, 1967	35-41	5-7	11-15	5-6		
Peru						
Morrow (unpubl.) (after Williams, 1967)	37-44	5-7	14-17	5-7		
East Africa						
Morrow (unpubl.) (after Williams, 1967)	36-41	5-7	11-15	5-6		
East Africa						
Merrett, 1971	38-42	5-7	13-16	5-6		19-23
Eastern North Pacific						
Wares and Sakagawa, 1974		5-7		5-7		

[1]The figure of 25 which appears here in the translation of the 1949 paper is evidently a misprint.

ypothesis by including eastern Pacific fish in the outhern population and demonstrating that growth f the pectoral fin is allometric with an inflection at bout 185-cm eye-fork length. They pointed out that nixing of the two populations occurs in the tropics nd suggested that Mexican fish belong principally to he southern-eastern group and that southern Califor-ia fish are derived from both groups but dominated y the northern.

Merrett (1971) found striped marlin of the tropical vestern Indian Ocean to have relatively long pectoral ns indicating a closer relation to the southern-east-rn population than to the northern.

There has been little mention in the literature of eographical variation in color pattern, but the strik-ng almost zebralike bars on the freshly caught New ealand specimen in the photo published by Gregory nd Conrad (1939) appears more pronounced than in sh from other regions.

To date, the knowledge of subpopulations, if they ruly exist, is insufficient for morphological definition.

The morphological changes of larvae and adoles-ent phases are of course remarkable. See 3.22 and .23.

Morrow (1952a) found significant ($P < 0.001$) egative allometry in the dorsal and ventral lobes of ne caudal fin and a slight but not statistically signifi-ant ($P = 0.065$) negative allometry in the length of ne pectoral fin of 49 adult fish from Cape Brett, New ealand. The pelvic fins show extreme negative allometry appearing to cease growth after reaching a certain size (Morrow, 1957; Royce, 1957). Negative allometry was found in Peruvian specimens for body depth and snout tip to origin of first anal fin (Morrow, 1957). Ueyanagi (1957b) found extreme negative and positive allometry, respectively, in central dorsal rays relative to body depth and pectoral length relative to body depth in young specimens (85-175 cm) which proved the synonymy of *Kajikia formosana* and *T. audax*. Royce (1957) also found similar allometry.

Ontogenetic change in body form is shown in Figure 1. The morphological change of the snout (its growth relative to the body length) from the postlarval to adult stage of striped marlin is relatively small in comparison with those of other istiophorid species. However, the change of the dorsal fin shape during growth is remarkable in this species (Ueyanagi, 1963b).

In relation to functional morphology, Fierstine (1968) found an average aspect ratio (span/surface area of one side of fin) of 9.0 for the caudal fin of three striped marlin from the eastern Pacific. This high ratio, indicating relative efficency as a hydrofoil, was greater than that for seven other scombroid species but less than that for the sailfish, *Istiophorus platypterus*, and the white marlin, *T. albidus* (10.0 and 10.3, respectively).

Published weight-length regression constants are summarized in Table 2.



Figure 1.—Ontogenetic change in body form of striped marlin. A. 7.9 mm SL, B. 21.2 mm SL, C. 121.5 mm body length, D. 1,050 mm body length, E. 1,882 mm body length. (From Nakamura, 1968.)

1.32 Cytomorphology

No data available.

1.33 Protein specificity

No data available.

2 DISTRIBUTION

2.1 Total Area

In the eidogical classification of Parin (1968) the striped marlin is holoepipelagic, i.e., it inhabits the isothermic surface pelagic layer of the ocean at all stages of its life cycle. Such species are chiefly limited in distribution to the tropics, where a permanent th mocline exists, but penetrate higher latitudes in warm season. Atypical of the distribution of m scombroids, the striped marlin seems to prefer more temperate waters. In the Pacific the distribut resembles that of the albacore, *Thunnus alalun* and bluefin tuna, *T. thynnus*, in contrast to that the other billfishes and tunas (Howard and Ueyana 1965; Parin, 1968), however, in the Indian Ocean striped marlin distribution is centered in warm waters.

Striped marlin occur throughout the warmer wat of the Indian and Pacific oceans. The species rang eastward to the coast of the American continents a westward to the African coast. Off South Africa th are found a slight distance into Atlantic wat (Talbot and Penrith, 1962). Extreme poleward di tribution has been recorded to lat. 40°-45° in bo hemispheres. In the north this occurs in the Kurosh extension, primarily between long. 165°E and 18 but also at long. 150°W (Fisheries Agency of Japa Research Division, 1969-71). In the southe hemisphere this occurs in the Agulhas Curre (Talbot and Penrith, 1962; Fisheries Agency of Japa Research Division, 1969-72) and also rarely at lon 105°E (Fisheries Agency of Japan, Research Divisio 1969), which appears to be West Wind Drift water. (the eastern perimeter of the Pacific, Point Concepti (lat. 35°N) and Chañaral, Chile (lat. 29°S) appear be the northern and southern limits of distributio

The broad geographical distribution of this speci makes it difficult to generalize on the physical a biological characteristics of the areas inhabite Temperature, however, is one parameter which h been considered to influence total distribution. T 20° and 25°C isotherms tend generally to bound t total distribution at least in the western Pacif (Howard and Ueyanagi, 1965).

2.2 Differential Distribution

2.21 Spawn, larvae, and juveniles

Although information is lacking on the distributio of eggs, there are several reports (Ueyanagi, 195 1964; Jones and Kumaran, 1964; Nishikawa a Ueyanagi, 1969) pertaining to the distribution of la vae.

In the Pacific, larvae have been observed in th northwestern Pacific (west of long. 180°) between la 10° and 30°N; and in the South Pacific (west of lon 130°W) between lat. 10° and 30°S. The larvae a most abundant in early summer, with the peak o currence in the northwestern Pacific during Ma June, and in the South Pacific in Novembe December. The seasonal occurrence of mature femal coincides with that of the larvae (Ueyanagi, 1964 While the distribution of larvae is not known for th

Table 2.—Weight-length constants for *Tetrapturus audax* (log W = log a + b log L).

Location	Weight	Units of weight	Length	Units of length	Number	Approximate length range of specimens	a	b	s	Source
New Zealand and Australia	Whole	lb	FL[1]	cm	27	265-310	−6.515	3.624	0.045	Gregory and Conrad, 1939 (after Royce, 1957)
New Zealand	Whole	lb	FL[1]	cm	48	218-310	−5.024	3.011	0.056	Morrow, 1952a (after Royce, 1957)
Central Pacific	Whole	lb	FL	cm	13	142-304	−6.648	3.691	0.502	Royce, 1957
Hawaii	Whole[2]	lb	FL[3]	cm	30	166-253	−6.110	3.446	0.048	Hawaii Division of Fish and Game (after Royce, 1957)
East Africa	Whole	lb	FL	cm	98	170-270	−4.629	2.844	?	Williams, 1967
Fiji area	Gilled and gutted	kan[4]	Eye-fork[2]	cm	562	160-260	−6.737	3.504	?	Koga, 1967
Equatorial west Indian Ocean	Whole	lb	Eye-fork	cm	156	120-196	−4.782	[5]3.062	?	Merrett, 1968c
Eastern Pacific	Whole	kg	Eye-fork	cm	51	108-211	−5.255	3.089	?	Kume and Joseph, 1969b
Eastern Pacific	Whole	lb	Eye-fork	cm	1,982	110-215	−4.816	3.072		Wares and Sakagawa, 1974
Eastern Pacific	Whole	kg	Eye-fork	cm	1,982	110-215	−5.157	3.071		Wares and Sakagawa, 1974
Eastern Pacific	Whole	lb	FL	cm	535	153-271	−5.007	2.986		Wares and Sakagawa, 1974
Eastern Pacific	Whole	kg	FL	cm	535	153-271	−5.340	2.982		Wares and Sakagawa, 1974

[1]The original measurement was SL which apparently Royce converted to FL on the basis of regression for nine Hawaiian specimens and mislabeled "*total*" length in caption to Appendix Table 3-A.

[2]Presumed.

[3]The original measurement was naris to fork which Royce converted to FL on the basis of regression for 21 Hawaiian specimens.

[4]1 kan = 3.75 kg = 8.287 lb.

[5]±0.197.

eastern Pacific (east of long. 120°W), mature fish are reported to occur there between lat. 5° and 20°N, largely in May-June (Kume and Joseph, 1969b).

In the Indian Ocean, larvae have been reported to occur in the Banda and Timor seas during January to February (Ueyanagi, 1959), and in the western Indian Ocean during December to January between lat. 10°S and 18°S and in the eastern Indian Ocean during October to November between lat. 6°N and 6°S (Jones and Kumaran, 1964). Mature females are reported to occur in March-May in the Bay of Bengal although larval occurrence is not yet known there (Ueyanagi, 1964).

The lower temperature limit in the distribution of larvae is approximately 24°C both in the Indian and Pacific oceans. However, the distributions differ in that in the Pacific the larvae are scarcely found in equatorial waters. It is noted that striped marlin larvae are not likely to appear in the Kuroshio area, while sailfish larvae occur there exclusively (Ueyanagi, 1959).

Information is very sparse on the distribution of the juveniles. Nakamura (1968) reported on two juveniles (body length, 12.15 and 14.5 cm) found in stomachs of a yellowfin tuna, *Thunnas albacares*, and a dolphin, *Coryphaena hippurus*, taken by longline. One juvenile was found on 13 January 1955 at lat. 23°52′S, long. 175°49′W and the other on 21 December 1964 at lat. 17°57′S, long. 67°29′E. These two occurrences coincide with larval distributions in the South Pacific and Indian Ocean, respectively.

2.22 Adults

Major areas of high abundance are the western Arabian Sea, the central North Pacific (lat. 15°-30°N) and the eastern Pacific as shown in Figure 2. T figure shows distributions based on highest quar year hook rates by 5° squares for the years 196 (Fisheries Agency of Japan, Research Division, 19 72). Lesser areas of abundance occur off South Afr northern Madagascar, northern Sumatra, Sri Lan eastern and western Australia, central Pacific coas Japan, and the south central Pacific.

The high density in the Arabian Sea appears to seasonal, occurring in the second quarter of the ye However, fishing effort in other seasons has been v slight. Off East Africa (lat. 0°-12°S) where strip marlin are the most abundant marlin, the works Merrett (1968a, b) showed highest hook rates occur between lat. 2° and 4°S. The hook rate was six tim higher in the northeast monsoon (Nov.-Mar.) than the southeast monsoon (Apr.-Oct.).

The season of high abundance off western a eastern Australia is the fourth quarter (Fisher Agency of Japan, Research Division, 1969-72). Ko (1967) states, however, that good catches occur winter off western Australia. In the South Paci Koga states that it is a remarkable feature that t main fishing areas, which show good catches fr August to December, show very poor catches fr January to July.

In the northwestern Pacific, striped marlin abundant in Formosan waters, both in the Kurosh and in the South China Sea, during the wh northeast monsoon season with a peak in the mid of that season. Later in the spring, they move no into the waters of Japan where they appear at abo the same time as the albacore do (Nakamura, 194

In the Hawaiian area, striped marlin occur from through spring with the seasonal distribution be



Figure 2.—Areas of high apparent abundance of striped marlin, 1967-69. (After Fisheries Agency of Japan, Research Division, 1969-72.)

mplimentary to that of the blue marlin, *Makaira gricans*, which occurs primarily in the summer trasburg, 1970).

In the eastern Pacific, striped marlin are present roughout the year from lat. 30°N to 30°S. High undance is maintained throughout the year in the eas of the Revilla Gigedo Islands, Baja California, cuador, the Galapagos Islands, and in the high-sea ea bounded by long. 90°-110°W and lat. 10°-30°S Kume and Joseph, 1969b). Seasonal changes in ap-rent density are marked. The concentration tween Baja California and the Revilla Gigedo ands, which is restricted to a narrow band in the st quarter, expands southeast along the coast and award to long. 115°W during the second quarter, d then north to lat. 28°N, south to lat. 3°N and award to long. 125°W during the third quarter. Dur-g the fourth quarter the southern extension expands rther to long. 130°W. Another seasonal concentra-n develops in the second and third quarters in the shore area between long. 100° and 115°W centered about lat. 8°-13°N. The area of high density around e Galapagos Islands extends eastward to the coast Ecuador in the third and fourth quarters and then cedes again in the first.

The sport fisheries for striped marlin off Mexico d southern California are seasonal. In southern lifornia it is highly seasonal with almost all fish be-g taken between August and October. In Mexico me fish may be taken year round, but the best hing occurs from December through March at azatlán and April through August near the tip of ja California (Eldridge and Wares, 1974).

In addition to differential distribution in density, ere is also some differential distribution in size. In e eastern Pacific, fish on the southern spawning ounds are larger than those on the northern. The ngth frequency of the southern group has a single ode at 180-200 cm whereas that of the northern oup has two modes, one at 140 cm and one at 180 cm Kume and Joseph, 1969b). In the western Pacific titudinal stratification occurs. Honma and amimura (1958) show small marlin occurring in uatorial waters; these small fish are absent in the gion of lat. 5°-16°S. In mid-latitudes (15°-30°S) of e central South Pacific longitudinal stratification is parent; larger fish (> 180 cm) occur in the western cific (Koga, 1967). There may also be some vertical atification. Furukawa, Koto, and Kodama (1958) nd harpooned fish to be larger than longline-ught fish in the East China Sea. The harpooned fish re also fatter at a given length.

Off Formosa the smaller fish, which were long ought to be the separate species *Kajikia formosana*, cur with the shortbill spearfish, *Tetrapturus gustirostris*, offshore from the center of the uroshio, while the larger fish occur inshore Nakamura, 1949).

2.3 Determinants of Distribution Changes

Probably behavioristic factors related to feeding and reproduction are the primary determinants of changes in distribution. These in turn are affected by the seasonal cycle of warming of the surface waters, development of thermoclines and currents, and seasonal cycles in abundance of food organisms. The subject of the factors and relationships causing concentration of striped marlin has not received much discussion in the literature.

Nishimura and Abe (1971) have found a correlation between the position of the Kuroshio as indicated by the latitude at which the current crosses 139°30'E, and the catch made off Izu.

Kume and Joseph (1969b) noted that the appearance of the area of high hook rate centered at lat. 8°-13°N from long. 100° to 115°W is associated with the strong development of the Equatorial Countercurrent. They noted further that a diagonal band of high abundance extending from lat. 5°S, long. 120°W to lat. 8°S, long. 95°W was in the general region of the eastern extension of the South Equatorial Countercurrent.

In the area west of Australia, both the striped marlin and southern bluefin tuna grounds are centered at the boundary of currents which run along long. 115° E meridian in winter (Koga, 1967).

Manning (1957) states that off Chile, striped marlin are found in the green water that is normally found from shore to 10 to 25 miles offshore. Their occurrence in these waters was in common with bonito, sardines, and anchovies and in contrast to swordfish which occurred farther offshore in the blue and white waters.

Furukawa et al. (1958) show the fishing ground in the western Pacific associated with the surfacing of the 20°C isotherm over the edge of the continental shelf.

Nakamura (1938) states that surfacing of fish is associated with high waves generated by opposing wind and current as in the case of the Kuroshio and the northeast monsoon.

2.4 Hybridization

No record.

3 BIONOMICS AND LIFE HISTORY

3.1 Reproduction

3.11 Sexuality

Striped marlin are heterosexual with no reported intersexuality or hermaphroditism.

Sexual disorphism has not been reported in this species and the sexes are indistinguishable externally. Nakamura (1949) mentioned the sexual difference in body size is not great in the genus *Tetrapturus* in contrast to the case in *Makaira*. Differences in greatest

size or modal size between the sexes are small. Ueyanagi (1953) found, in the northwestern Pacific, that the modal size of males is about 10 cm smaller than that of females. Koga (1967) showed a length-frequency distribution by sex for 210 fish from the Fiji area in which the modal lengths for males and females were 195 and 205 cm, respectively. In a sample of 105 striped marlin taken by longline off East Africa, Williams (1967) found males did not exceed 240 cm fork length (equivalent to 182 cm eye-fork length from regression of Merrett, 1971). About 16 females were taken above this length to a size of about 270 cm (205 cm eye-fork length). Merrett (1971), however, found males up to 193 cm eye-fork length off East Africa. Modal size differences between the sexes were not found in Hawaiian fish (Strasburg, 1970) nor in the eastern Pacific (Kume and Joseph, 1969b).

3.12 Maturity

As has been found in other pelagic species such as albacore and dolphin, quantitative measure of maturity for males is difficult. There is only a small increase in testis size during the maturation cycle. Merrett (1971) found little correlation between relative testis size and maturity-stages based on microscopic examinations. In fact, Merrett (1970) suggests that there is continuous availability of spermatozoa in mature males based on differential maturation of the testicular lobules and the possession of a muscular seminal vesicle.

In the female, maturation is synchronous throughout the ovary and seasonal maturation is accompanied by marked increase in relative size of the gonads. Data from Kume and Joseph (1969b) showed a twentyfold increase. Moreover, there is good correlation of relative ovary weight and mean maximum egg diameter (Merrett, 1971; Eldridge and Wares, 1974).

Williams (1967) concluded from longline data in East Africa that first maturity was reached between 180 and 200 cm fork length (50-80 lb) which is equivalent to 141-157 cm eye-fork length (Merrett, 1971). Merrett reported similar results, 140-160 cm or 62-93 pounds. Ueyanagi (1957b) mentioned that 154 cm eye-fork length was the smallest size found in the spawning group of the northwestern Pacific. Kume and Joseph (1969b), using a gonad index, reported that individuals from the eastern Pacific do not regularly enter the spawning group until reaching about 160 cm eye-fork length but found one as small as 148 cm. Other data from the eastern Pacific (Eldridge and Wares, 1974) agree with these conclusions.

Since age at specific size is not known for striped marlin, age at maturity is also unknown. Koga (1967), however, stated "it is likely that growth rate of this species shows different values between the Indian Ocean and the Pacific Ocean and the groups in the Indian Ocean attain maturity earlier than those in Pacific."

3.13 Mating

Nothing has been published relating to mat habits of this species.

3.14 Fertilization

Fertilization is externally.

3.15 Gonads

Merrett (1970) has described the gonads of billf in detail. The following description is taken from work.

> The gonads are paired organs lying in the posterior half of body cavity, on each side of the stomach and intestine. They suspended from the lateral edges of the chambered air blad by mesenteries . . . the gonads are almost bilatera symmetrical and both terminate at their point of discharg the exterior in the urino-genital papilla . . . which lies poste to, and in a common groove with, the anus . . . The urinary genital systems are closely linked.
>
> The ovaries are elongate sausage-shaped organs, which ta at both ends, and joined only at their posterior ends . . . A str muscular sheath binds the ovaries to the urino-genital pap and the basal part of the intestine. They are invested in th layers of connective tissue which sometimes contain deposit fat; fat is also occasionally found in the mesovarium. Bene the connective tissue the ovaries are pale flesh-pink to wine in colour, depending upon the stage of maturity. Internally certain stages, a central lumen runs the length of the ovary the ovaries have been found to be unequal in length . . . eit ovary can be the longer of a pair.
>
> . . . Immediately posterior to . . . [the anal papilla] . . . is urino-genital papilla. In the female this carries only the urin duct. The point of discharge of the ovaries is situated betw the bases of the anal papilla and the urino-genital papilla.

This opening between the anal and urino-geni papillae which is absent in the males should serve an external characteristic in distinguishing the sex But in the experience of the junior author t difference is difficult to observe consistently in t species. It is more obvious in the sailfish.

Nakamura (1949) stated that the fecundity billfishes ranges from 1 to 1.2 million eggs, dependi on fish size and species. Morrow (1964) estimate million eggs for New Zealand marlins. These appe to be low estimates, however. Merrett (1971) report an estimated fecundity of 12 million for one Indi Ocean specimen of 182 cm eye-fork length, 126 poun with ovary weight of 1.53 kg, and mean maximum e diameter of 0.470 mm. Eldridge and Wares (197 reported fecundity estimates of three eastern Paci specimens which ranged from 11 to 29 million eg These fish ranged in size from 150 to 180 cm eye-fo length but the fecundities showed no relation to size the fish. Gosline and Brock (1960) estimated 1

illion eggs were contained in one ovary of a 154-
ound striped marlin landed in Honolulu. The other
ary of this fish was immature.

3.16 Spawning

Examination of size-frequency distributions of egg
ameters (Eldridge and Wares, 1974) indicates only
ne spawning per season.

In the western Indian Ocean it appears that the
gh catch rates during the northeast monsoon period
hich peak from December to February are associated
ith a postspawning feeding migration (Williams,
967; Merrett, 1971). Spawning must occur elsewhere
the Indian Ocean.

From larval occurrence, spawning was suggested to
ke place in the Banda and Timor seas during
nuary to February (Ueyanagi, 1959). On the basis of
rval occurrence, Jones and Kumaran (1964) stated
at striped marlin spawn in the western Indian
cean during December-January between lat. 10°
d 18°S and in the eastern Indian Ocean during Oc-
ber-November between lat. 6°N and 10°S. Further-
ore mature females are known to occur in March-
ay in the Bay of Bengal and in October-December
the waters south of the Lesser Sunda Islands
Jeyanagi, 1964).

In the western Pacific, mature females are found
om lat. 15° to 30° (north and south) in early
mmer, from May to June and October to January in
e northern and southern hemispheres, respectively.
arvae are also found in these areas (Ueyanagi, 1964).
akamura (1949) stated that in the Formosa area,
awning is thought to take place mainly in the South
hina Sea with its peak occurring from April to May.
oga (1967) reported that the spawning areas (lat.
°-30°S) in the western South Pacific is also the
ain fishing area and that the period of spawning cor-
sponds to the season of northward migration which
curs from September to November.

In the eastern Pacific, the spawning season also
pears to be the early summer in each hemisphere,
arters II and III in the northern and quarters IV and
n the southern. Highest frequencies of spawning fish
cur from May to June in the north and November to
ecember in the south (Kume and Joseph, 1969b).
vidence of spawning in the eastern Pacific is based
ly on relative gonad sizes of females. The northern
awning area appears to be isolated in a narrow band
om long. 107° to 114°W extending from about lat. 6°
19°N (Kume and Joseph, 1969b).

Ovaries of striped marlin caught in the Mexican
ort fishery undergo rapid development in June; ripe
h were never observed (Eldridge and Wares, 1974).
panese fishermen, however, have reported ripe and
nning ripe striped marlin in the waters around
corro Island from June to October (J. L. Squire, Jr.,
uthwest Fisheries Center, National Marine
sheries Service, NOAA, La Jolla, Calif., pers. com-
un.).

The southern spawning area appears fairly well confined to lat. 20°-25°S and long. 125°-130°W (Kume and Joseph, 1969b). Highest frequencies of the mature females occur in November and December.

Nakamura (1949) stated that sex ratios approached 1:1 at the peak of the spawning season. However, it was found that males dominate during the spawning season in the northwestern Pacific (Nakamura, Yabuta, and Ueyanagi, 1953). Kuma and Joseph (1969b) also found a high proportion of males in the spawning groups in the eastern Pacific. Male:female ratios ranged from 1.8 to 6.6 in spawning groups, whereas in nonspawning groups they tended to be less than 0.5 and to decrease with increased size of fish.

3.17 Spawn

There is little information pertaining to the eggs of this species. Nakamura (1949) mentioned that the external morphology of the eggs of striped marlin closely resembles that of sailfish eggs which are spherical, transparent, and buoyant, with a single oil globule and with no special structure on the egg membrane. Morrow (1964) reported that the ovarian eggs of striped marlin from New Zealand average about 0.85 mm in diameter. Size of the ovulated eggs of this species is presumed to exceed 1 mm in diameter considering that the mean diameter of eggs for shortbill spearfish is 1.442 mm and for sailfish is 1.304 mm as reported by Merrett (1970).

3.2 Pre-Adult Phase

3.21 Embryonic phase

No information available.

3.22 Larval phase

The postlarval stage of striped marlin is described in detail by Ueyanagi (1959). The study was based on 40 specimens ranging from 2.9 to 21.2 mm in standard length, collected from the northwestern Pacific, South Pacific, and the Indian Ocean. These specimens were captured by surface tows of the larvae net. The morphology of the striped marlin postlarvae is similar to that of other istiophorid species in the development and degeneration of head spination, fin formation, pigmentation, etc. Figure 3 from Ueyanagi (1959, 1963a) represents the postlarval stage of this species from an early stage with short snout to an advanced stage with elongated jaws. The snout begins to lengthen at around 7 mm standard length. Head spination becomes most conspicuous at this size. Fin rays of each fin reach their full complement at around 20 mm standard length. Pigmentation extends almost all over the body surface and on the dorsal fin membrane at this size. The dorsal fin begins to increase its height at around 10 mm and stands very high like a sail in larvae exceeding 20 mm. The key diagnostic character for the striped marlin larvae was reported as



Figure 3.—Larvae of striped marlin. From top to bottom: 2.9, 5.0, 12.0, and 21.2 mm SL. (From Ueyanagi, 1959, 1963a.)

llows: ". . . the profile of head, tip of snout and enter of eye are on a nearly equal level" (Ueyanagi, 963a). In addition, the shape (arrangement) of the terotic and preopercular spines is diagnostic for lar-al identification (Ueyanagi, 1974).

On the vertical distribution of billfish larvae, eyanagi (1964) indicated that larvae appear to be istributed most abundantly in the surface layer dur-ng the daytime and vertical, diurnal migration seems o occur in the upper 50 m of waters (Table 3.). illfish larvae appear from time to time in the tomach contents of the larvae and juveniles of ailfish and swordfish, *Xiphias gladius* (Arata, 1954; ehringer, 1956). It is assumed that striped marlin rvae are therefore preyed upon by many surface eding species, including the billfishes. Considering ne very large numbers of eggs spawned by the striped narlin (see 3.15), it appears that mortality at the lar-al stage is extremely high.

ime of first feeding: No direct information is vailable. There is an observation that five copepods ere seen in the stomach of a 3.9-mm sailfish larva Gehringer, 1956).

ype of feeding: There is no information pertaining o the larvae of striped marlin. Of the larvae of sailfish n the Atlantic, Gehringer (1956) noted that copepods constituted the food of specimens less than mm. long. At this size fish larvae also were eaten, nd no specimen exceeding 13 mm. had copepods in s stomach." Furthermore, it is also known that ailfish larvae have consumed istiophorid larvae half s long as their own body length. It is believed that ne larvae of striped marlin, like the sailfish, begin to eed on fish larvae after reaching a size of about 7 mm.

Table 3.—Comparison of numbers (and percentage) of tows from which istiophorid larvae were captured, showing the various depths and times of towing, in the western Pacific area. (From Ueyanagi, 1964.)

Depth (m)	32 day tows		31 night tows	
	Number	Percent	Number	Percent
0	22	68.8	12	38.7
ca. 20	8	25.0	12	38.7
40-50	2	6.2	10	32.3

3.23 Adolescent phase

Nakamura (1968) described the juveniles of striped marlin based on two specimens collected from the South Pacific and the western Indian Ocean (see 2.21). His drawing has been reproduced as Figure 4. In the juvenile stage the snout is very elongate and, in fact, is longest at this stage relative to body length. The shape of the first dorsal fin still differs from the adult configuration, being highest anteriorly and decreasing gradually in height posteriorly. The arrangement of the viscera is similar to that of the adult. Nakamura also mentioned that the juvenile of striped marlin is similar to that of the white marlin in many respects except that the former has no ocelli on the first dorsal fin.

It is believed that large, powerful pelagic fishes such as tunas, billfishes, dolphins, etc., are the principal predators of juvenile striped marlin.

Little is known about parasites in juvenile striped marlin.

Immature striped marlin (80-100 cm eye-fork



Figure 4.—Juvenile striped marlin (121.5 mm body length) collected from the southwestern Indian Ocean. (From Nakamura, 1968.)

length) are taken by longline. The shape of the first dorsal fin in these fish differs from the adults in that the posterior rays are still high (Ueyanagi, 1957b). Fish and squid are found in the stomachs, as in the adults.

Following the 1954 hydrogen bomb test in Bikini, it was reported that virtually all of the radioactively contaminated striped marlin found in the northwestern Pacific were fish under 130 cm long (Anon., 1955). It is not clear what the implications are with respect to ecological differences between young and adult fish.

3.3 Adult Phase (Mature Fish)

3.31 Longevity

The ability to determine age of individual striped marlin has not been developed and thus life tables cannot be developed and life expectancy and maximum age are unknown.

Koto (1963a), using size-frequency data from western Pacific catches, was able to discern six age classes (which he designates as n though $n + 5$) in fish greater than 100 cm eye-fork length. The Walford growth transformation of his data indicated an ultimate size of about 290 cm. Specimens of this size are apparently occasionally taken in the South Pacific (Honma and Kamimura, 1958). From the general pattern of growth indicated by Koto, fish of this size would be expected to be at least 10 yr old.

3.32 Hardiness

Very little is known. Because of the large size and activity of this species, physiological experimentation is difficult. The return rate for tagged fish is much higher for striped marlin than for sailfish (Squire, 1974), and this may indicate greater general hardiness for striped marlin. Healing of severely broken bills (Wisner, 1958) may also be an indication of hardiness. Sportsmen usually report that the striped marlin fights harder than the sailfish when hooked and tends not to die as easily as the sailfish.

3.33 Competitors

The other billfishes, particularly the smaller species (sailfish and probably the shortbill spearfish), swordfish, and the larger tunas are probably the closest competitors for food. Even the smaller scombrids share many forage species with the striped marlin. The dolphin and pelagic sharks such as *Prionace*, *Carcharhinus*, and *Isurus* utilize many of the same forage species (Parin, 1968). Striped marlin tend to feed more on epipelagic species and less on mesopelagic species than the oceanic tunas or the swordfish.

3.34 Predators

Predators of adults are probably extremely limit the only likely candidates being some of the la pelagic sharks and the toothed whales.

Bills of billfishes have been found in floating obje and other fish. Occasionally bills of striped marlin found to have been broken off, and the fish are kno sometimes to ram fishing boats when hooked, but i not certain that any of these occurrences have a relation to defense or aggressive action.

3.35 Parasites, diseases, injuries, and normalities

Parasites and diseases: The body surface usua harbors many caligoid copepods which frequen congregate on the ventral surface, particularly in area around the anal fin and on the head. Willia (1967) reports they may occur in the thousands. T skin surface in areas of concentration often appe red and irritated. Penellid copepods (*Penella fillo* are frequently found penetrating the skin a anchored in the muscle or sometimes in inter organs such as the gonads. Koga (1967) reports t percentage occurrence of caligoid copepods a penellid copepods on striped marlin in Fiji waters w 100% and 20%, respectively. Eldridge and Wa (1974) report the percent occurrence of penell above and below the lateral line on one side of t body as 26.2% and 22.8% with average infections 3.3 and 2.3 copepods per fish. Stalked barnac (*Conchoderma irgatum*) frequently are attached the penellid copepods and often to the marlin sk normally near the vent (Williams, 1967). Digene trematodes were reported found on the gills Williams (1967). Monogenetic trematodes of the fa ly Capsalidae are quite common on the surface of t skin (Eldridge and Wares, 1974).

Cestode worms resembling *Dibothriu manubriformes* have been found in the intestines a nematodes (*Contracaecum incurvum*) are very co mon in the stomach occurring in densities grea than 200 per stomach (Morrow, 1952b). The copepo *Philichthys xiphiae*, has been found in the muc canals of the preopercular and opercular bones, a capsalid trematodes are commonly found in the na capsule (Eldridge and Wares, 1974).

Injuries and abnormalities: Gastric ulcers we reported in 14% of 563 eastern Pacific striped mar (Evans and Wares, 1972). These may be associat with the presence of nematodes (R. T. B. Iverse Southwest Region, National Marine Fisheries S vice, NOAA, Honolulu, Hawaii, pers. commun.). T small squaloid shark (*Isistius brasiliensis*), t probable cause of crater wounds on many pela fishes including istiophorids (Jones, 1971), probab parasitizes striped marlin.

There is evidently little ability for regeneration

roken bills and pelvic fins are seen to heal over rather ıan regenerate. Striped marlin have recovered after sing almost all of the bill (Wisner, 1958).

3.4 Nutrition and Growth

3.41 Feeding

Most active feeding probably takes place in the ıorning. LaMonte (1955) reported that squid found ı the stomachs of striped marlin off Peru and Chile ere less digested in fish caught in the morning than ıose landed after noon. Kobayashi and Yamaguchi 1971), examining only fish caught after noon, found a ecline in feeding activity toward sunset. Williams 1967), however, suggests that East African fish feed t any time of the day or night.

Apparently the food is usually captured by grasping ith the mandibles rather than by spearing, slashing, r clubbing with the bill. Fish which have lost the bill mpletely survived well. Some food specimens are ccasionally found, however, which have been neatly peared (Wisner, 1958; Evans and Wares, 1972).

3.42 Food

Several authors have reported on food habits. Table gives an idea of the variety of food species which ave been found most important in different studies. is notable that, despite the large size and lack of gill akers in striped marlin, relatively small forage items re commonly taken (Nakamura, 1949).

Food habits do not appear to vary appreciably with ex or size over the range of sizes commonly caught. onsiderable variation in species composition of the iet occurs, however, with season and geographic cation (Evans and Wares, 1972). Such variations robably reflect variations in availability of the food rganisms in keeping with the generally accepted con-ept that these fish are broadly carnivorous, nonselec-ve feeders. This is true of epipelagic fishes generally Parin, 1968).

The average volume of food found in stomachs of riped marlin caught by the eastern Pacific sport shery ranged from 14 to 23 fluid ounces per stomach Evans and Wares, 1972). Yamaguchi (1969) reports iat empty stomachs were found in 66% of striped arlin caught by longline. This percentage tends to e larger in billfish than in tunas. Evans and Wares 972) found 19% empty stomachs.

Table 4.—Some reported major food species for striped marlin.

Area	Food species	Author
East Africa	*Fistularia* sp. *Auxis* sp. Squid	Williams, 1967
New Zealand	*Scomberesox saurus* *Arripis trutta* *Loligo* sp.	Morrow, 1952b
New Zealand	*Ommastrephes sloani* *Caranx lutescens* *Scomber japonicus*	Baker, 1966
Tasman Sea	*Alepisauridae* *Clupeidae*	Koga, 1968
Bonin Islands	*Gempylus* sp. Cephalopods *Pseudoscopelus* sp. *Alepisaurus* *Ostracion* Crustacea	Yabuta, 1953
California	*Cololabis saira* *Engraulis mordax* *Sardinops caerulea*	Hubbs and Wisner, 1953
California	*Engraulis mordax* *Trachurus symmetricus*	Evans and Wares, 1972 Craig, 1972
Mazatlán, Mexico	Squid *Etrumeus teres* *Fistularia* *Argonauta* sp.	Evans and Wares, 1972
Baja California, Mexico	Squid *Etrumeus teres* *Scomber japonicus* *Fistularia* sp.	Evans and Wares, 1972
Peru-Chile	Squid	LaMonte, 1955
Chile	*Engraulis ringens* *Trachurus symmetricus* Squid	de Sylva, 1962

3.43 Growth rate

Weight-frequency modal progressions in Hawaii nding data suggest an annual growth rate of about pounds (Royce, 1957). Merrett (1971) found agree-ent with Royce's data in size-frequency dis-ibutions from East Africa. Possible modal lengths of out 152, 167, 177, and 197 cm eye-fork length were und in the length-frequency distribution for the tal catch. Computed weights at modal lengths gave annual weight increments of 27, 21, and 23 pounds.

Koto (1963a) working with length-frequency data from the western North Pacific, found six modal groups (n though $n+5$). The monthly progression of these modes is shown in Figure 5. The designations of these modal groups are as follows:

Group	*Lengths* (cm)	*Increment* (cm)
n	100-120	?
$n+1$	120-160	35
$n+2$	160-185	27
$n+3$	185-205	19
$n+4$	205-220	16
$n+5$	220-233	13

There is a marked seasonal change in growth rate with rapid growth occurring from June to November and very little growth in the remainer of the year for age groups $n+1$, $n+2$, and $n+3$. Age group n appears



Figure 5.—Seasonal changes in modes derived from average year's length frequencies of longline-caught striped marlin. A. North Pacific Current area; B. East China Sea (long. 120°-130°E). (From Koto, 1963a.)

in the catch only after September, as fish larger than 100 cm and shows rapid growth right through the winter. The failure of the fishery to capture many fish below 100 cm makes it impossible to say what the absolute ages of the groups are.

The annual growths of the $n+1$ and $n+2$ age groups differ between areas and years. Koto (1963b) showed that in the North Pacific Current area the yearly differences in the length composition were caused by differences in the average length of the $n+1$ age group. In the North Equatorial Current area, however, these differences were probably caused by differences in age composition and the relative abundance of the n and $n+1$ age groups.

Koto (1963b) has shown growth rate to be affected by population density. The very close inverse relationship between growth of the $n+1$ age group and the total fish abundance, or especially the abundance of the n and $n+1$ groups is shown in Figure 6. The correlation coefficients between growth of the $n+1$ age group and total fish abundance were -0.958 (df = 6) for the North Equatorial Current area and -0.737 (df = 8) for the North Pacific Current areas.

Royce (1957) suggested the maximum size reac by striped marlin is less than 226.8 kg (500 poun Records were given as follows: 172.8 kg (381 pour off New Zealand; 142.4 kg (314 pounds) from Pac equator; a "questionable" record of 196.9 kg (pounds) from the Hawaiian market, where norm only occasional specimens approach 136.1 kg (pounds); a somewhat questionable record becaus possible confusion with the blue marlin of 219.1 (483 pounds) from Chile; the world's record, taken California, was 692 pounds, which was based o misidentified blue marlin.

The theoretical maximum length of about 290 mentioned in section 3.31 is equivalent to 259.5 (572 pounds). A fish of 290 cm was taken by longl in the South Pacific (Honma and Kamimura, 19

Average semimonthly condition factors (K) co puted

$$K = \frac{W \times 10^5}{L^3}$$

where W is whole fish weight in kg
L is eye-fork length in cm

for eastern Pacific fish ranged approximately fr 0.80 to 1.26 (Eldridge and Wares, 1974).

In the East China Sea, condition factor

$$K = \frac{W}{L^3}$$

where W = gilled and gutted weight in kan (= 3 kg)
L = eye-fork length in m

was found to increase with body length from abou at 110 cm to a peak of about 10 at 180 cm and then decline to about 9 with larger fish up to 230



Figure 6.—Relation between total fish abundance and growth the $n+1$ age group. A North Equatorial Current area; B. No Pacific Current area. ____ growth per year; .....hook ra (From Koto, 1963b.)

Furukawa et al., 1958). The condition factor also showed a seasonal drop from about 10.8 in December to about 8.7 in June for fish of body length 166-195 cm. The drop in condition factor before June was much more abrupt than the increase after June. A similar seasonal cycle was also apparent for fish of 136-165 cm.

3.44 Metabolism

There are no data on metabolic rates for this species. Lindsey (1968) found body muscle temperatures as much as 2.6°C higher than the surrounding seawater. The highest temperature recorded occurred near the center of the epaxial muscle mass with a lesser maximum slightly below the hypaxial muscle mass center. Red muscle was found to be lower than white muscle at comparable depths beneath the skin. Temperatures of the viscera exceeded seawater temperatures by 0.7-1.3°C.

Barrett and Williams (1965) report the mean hemoglobin content in 16 striped marlin was 11.3 ± 0.75 g Hb/100 ml and ranged from 5.8 to 16.8.

3.5 Behavior

For feeding behavior, see 3.41; for reproductive behavior see 3.13, 3.21.

3.51 Migrations and local movements

See also 2.22 and 2.3.

The migration pattern of striped marlin appears to be principally a simple latitudinal movement between spawning areas and productive feeding areas (Parin, 1968). The movement is toward higher latitudes in the summer of each hemisphere and back toward equatorial waters in winter. In the northern hemisphere the peak of the northward migration is August-September and the southward migration begins in October and continues through February. The northward movement is also accompanied by an eastward expansion in the eastern Pacific (Howard and Ueyanagi, 1965; Fisheries Agency of Japan, Research Division, 1969-72).

In the central North Pacific small fish of about 13.6 kg (30 pounds) appear in Hawaii in winter, grow to 22.7-27.2 kg (50-60 pounds) by May or June, then migrate north for several months, and return to Hawaii as larger fish the next year. A similar pattern of migration is common to areas west of long. 180° (Howard and Ueyanagi, 1965).

In the southern hemisphere west of long. 150°W fish migrate north from south of lat. 30°S from August through November and form concentrations which are exploited in the area lat. 18°-19°S. The period of this migration corresponds to the spawning season. After November the fish appear to migrate south (Koga, 1967).

In the eastern South Pacific a high density area, which occurs in the area lat. 10°-17°S, long. 90°-115°W during the second and third quarters, appears to move southwest to the region of lat. 20°-28°S, long. 100°-110°W in the fourth and first quarters (Kume and Joseph, 1969a, b).

In the eastern North Pacific the seasonal north-south movements are apparent but less pronounced. Striped marlin do occur in their extreme northern range (southern California) during late summer and fall when surface temperatures reach a peak, but it is not clear whether these fish have come from the south or from the west (Howard and Ueyanagi, 1965). Data from fish tagged in the sport fisheries of southern California and Mexico provide evidence that striped marlin are capable of fairly long migratory movements up to 3,000 miles (Fig. 7). Some fish do move from California southward to the tip of Baja California and further, but there is no evidence of migration from Mexican waters to southern California (Squire, 1974).

Howard and Ueyanagi (1965) suggested on the basis of the appearance of an unusually small size group in California in 1958 and subsequent appearance 2 yr later of a small group in New Zealand, which was of a size expected for fish 2 yr older than the former, that there may be transpacific interchange between these remote areas.

Following the 1954 hydrogen bomb test in Bikini, contaminated fish were found only from the North Pacific, suggesting the possibility of separate populations in the two hemispheres (Nakamura, 1969).

In addition to the primary migratory trend, there are also lesser local movements reported. The area of high density off the central Mexican coast generally tends to expand westward seasonally reaching its maximum westward extent at about long. 130°W during the fourth quarter of the year (Kume and Joseph, 1969a, b). The region of high density around the Galapagos Islands during the second and third quarters expands eastward to the coast of Ecuador during the fourth and first quarters (Kume and Joseph, 1969a, b; Kume and Schaefer, 1966).

Nakamura (1949) mentions dense schools move from south to north along the coast of Vietnam in March and April.

In the western Pacific, Furukawa et al. (1958) report a gradual westward migration of fish from the vicinity of the Bonin Islands to the East China Sea in July and August where they stay until November after which a southward emigration takes place.

Migratory patterns in the Indian Ocean are unknown. The seasonal increase in density off East Africa during the northeast monsoon is believed to be a postspawning feeding migration (Williams, 1967; Merrett, 1971). The north-south type of seasonal movements that are typical in the Pacific are most evident off South Africa. A northward movement in the springtime (second quarter) is evident in the





Figure 7.—Movements of tagged strip marlin in the eastern Pacific Ocean

western Arabian Sea (Fisheries Agency of Japan, Research Division, 1969-72).

3.52 Schooling

Striped marlin, like the other istiophorids, do not form dense schools like the tuna, and the individuals are usually dispersed at wide intervals (Nakamura, 1949). Frequently, however, several fish are seen together; sometimes following one another, especially during the spawning season.

Surfacing is apparently more common when the wind and waves are high. When wind and current are moving in the same direction, the surface is calm and few fish are seen at the surface. When the wind runs counter to the current, high waves result, and fish are most often seen at the surface and are usually swimming in the direction of the wind, at least in Formosan waters (Nakamura, 1938).

When surfaced, the striped marlin usually is swimming very slowly with the upper caudal lobe above the surface and the dorsal fin retracted and not showing. This characteristic reportedly distinguishes them from swordfish which are unable to depress the dorsal fin and show both the dorsal and caudal fins when surfaced. Striped marlin swim faster and are less easily approached when surfaced than the swordfi (Philippi, 1887).

Little is known about how far the fish move v tically. Saito et al. (1972) report that striped mar have been caught at 150- to 290-m depths by verti longline experiments in Fiji waters.

For composition of stocks by size see also secti 2.22. As mentioned previously, there is considerab variation in size composition between various regio and particularly between the northern and southe groups in the Pacific (see, for example, Howard a Ueyanagi, 1965; Koga, 1967; and Kume and Josep 1969b). As mentioned earlier, southern fish tend to larger than northern fish throughout the Pacific. Si composition of striped marlin in the Indian Oce resembles the North Pacific size distribution mo than the South Pacific.

Striped marlin "schools" occur in waters in co mon with the schools of most of the Indo-Paci scombroids, particularly albacore, yellowfin tuna, a bigeye tuna, *Thunnus obesus*.

3.53 Responses to stimuli

Very little is known of this subject as it relates striped marlin. The possible response to temperatu

as been mentioned previously. It may be pertinent to ote here that when billfishes are caught by hook or arpoon, they first make several leaps into the air and hen swim wildly in broad circles near the surface. unas, on the other hand, try to escape in a vertical irection by diving deep (Nakamura, 1949).

POPULATION

4.1 Structure

4.11 Sex ratio

Sex ratio of the population as a whole is unknown; ut from the data of Williams (1967), Kume and oseph (1969b), and Merrett (1971), females usually redominate in longline catches. The notable excepion to this is on the spawning grounds, where males end to predominate (Kume and Joseph, 1969b). The ercentage of females tends to increase with size of ish (Kume and Joseph, 1969b). There is a tendency or spawning grounds to be dominated by larger fish han in nonspawning areas.

4.12 Age composition

Since age determination of individual fish is not ossible at present, age composition of the population as not been studied. See 3.43 on growth rate.

4.13 Size composition

Size composition varies greatly between stocks and with seasons (see 3.52).

Size at first capture by tuna longline gear is about 80 cm eye-fork length.

Size at first maturity is between 140 and 160 cm eye-fork length.

Maximum size is probably about 290 cm eye-fork length or 258.6 kg (570 pounds).

See 1.31 for length-weight relationships.

4.2 Abundance and Density (of Population)

4.21 Average abundance

No data.

4.22 Changes in abundance

Relative abundance in terms of average CPUE (catch per unit of effort) for major ocean areas is shown in Table 5 for the years 1962-70. These data show no apparent trend for the eastern, northern or the Pacific Ocean at large, however, there is a general decline noted for the South Pacific and a slight decline in the Indian Ocean, the latter since 1966 (Fisheries Agency of Japan, Research Division, 1965-72).

Year-to-year changes in CPUE for the various

Table 5.—Catch statistics by major ocean areas of Japanese longline fishery for striped marlin for 1962-1970 (Fisheries Agency of Japan, Research Division, 1965-72).

	1962	1963	1964	1965	1966	1967	1968	1969	1970
Indian Ocean									
E[1]	68.4	57.3	68.9	79.9	89.6	126	119	101	78
C[2]	48	34	38	81	106	114	63	59	45
CPUE[3]	0.07	0.06	0.06	0.10	0.12	0.09	0.05	0.06	0.06
Whole Pacific									
E	290	337	283	288	301	306	286	306	282
C	287	338	508	421	351	406	506	323	450
CPUE	0.10	0.10	0.18	0.14	0.12	0.13	0.18	0.11	0.16
North Pacific									
E	156	154	140	153	150	193	165	169	163
C	144	123	210	156	98	159	154	101	242
CPUE	0.09	0.08	0.15	0.10	0.06	0.08	0.09	0.06	0.15
South Pacific									
E	109	131	81.2	92.2	104	70.6	71.0	70.1	67.7
C	59	49	28	29	30	17	14	14	32
CPUE	0.05	0.04	0.03	0.03	0.03	0.02	0.02	0.02	0.05
Eastern Pacific (E of 130°W)									
E	24.7	52.1	62.0	43.6	47.5	42.4	50.4	67.2	52.1
C	84	166	270	236	223	230	338	208	177
CPUE	0.34	0.32	0.44	0.54	0.47	0.54	0.67	0.31	0.34

[1]E = Effort in hooks $\times 10^6$.
[2]C = Catch $\times 10^3$.
[3]CPUE = Catch/100 hooks.

regions of the Pacific and Indian oceans are examined in section 5.41.

Seasonal variations in available stock are marked. See 2.22 on differential distribution. Kume and Joseph (1969a) have shown that there is a threefold seasonal fluctuation in CPUE for the various regions in the eastern Pacific.

4.23 Average density

No data.

4.24 Changes in density

No data.

4.3 Natality and Recruitment

4.31 Reproduction rates

Annual egg production rates have not been estimated. Little is known of fecundity relationship with fish size. See 3.15 for some estimates.

Nothing is known of survival rates of eggs and larvae because they are so rarely collected.

4.32 Factors affecting reproduction

No data.

4.33 Recruitment

There is little information pertaining to the variation in annual recruitment (see 3.43 and 4.24).

4.4 Mortality and Morbidity

No work has been done on mortality rates or causes of mortality.

4.5 Dynamics of Population (as a Whole)

No work has been done.

4.6 The Population in the Community and Ecosystem

There is no specific information on this sub available; some general information on distribu and life history is presented in sections 2 and 3.

5 EXPLOITATION

5.1 Fishing Equipment

5.11 Gear

Virtually all of the commercial catch of stri marlin is by longlining. The harpoon fishery billfishes is responsible for less than 5% of the t catch.

The longline gear aims largely at tunas billfishes which are distributed at depths of aro 100-150 m. The gear consists of mainline, float li branch lines, hooks, and buoys. The constructio the longline gear differs according to the species of sought, but Morita (1969) presents an example "standard" gear (Table 6). Several hundred of th units (each unit is referred to as a "basket") are joi in a series to make up a set. The gear is retrieved w a longline hauler.

Suda and Schaefer (1965) gave examples of various types of longline gear presently in use (Ta 7), and indicated that types 1 through 5 are the m typical ones in use. They also included the estima hook depths for the different types of gear.

Billfishes are generally found closer to the sea s face than tunas. Therefore, in fishing primarily billfishes, the longline gear is modified by shorten the float line and the branch lines and also by add another buoy in the middle of the mainline secti the latter bringing the hooks closer to the surfa Furukawa et al. (1957) reported that for fish billfishes in the East China Sea, the combined len

Table 6.—General form of tuna longline. (From Morita, 1969.)

Name of part	Material	Length	Number u for 1 bas
Mainline	Cremona (20S, 55×3×3)	250 m[1]	1
Branch line	Cremona (20S, 55×3×3)	11 m	4
Sekiyama	Steel wire (27 # 3×3) and hemp yarn coiled with thread No. 5)	5.5 m	4
Kanayama	Steel wire (27 #, 3×3, Type M)	3 m	4
Hook	Steel	3.8 sun[2]	4
Float Line	Cremona (20S, 55×3×3)	22 m	1
Flag buoy	Flag, bamboo, float (glass ball or synthetic resin ball)	—	1
Radio buoy			2 or 3 (fo basket

[1]Length per one basket.
[2]1 sun = 3.03 cm.

Table 7.—Some examples of Japanese tuna longline gear. (From Suda and Schaefer, 1965.)

Type no.	1	2	3	4	5	6	7	8
A. The structure of one basket of the line (unit of length: meters)								
Length of mainline	300	300	300	300	300	245	360	350
Length of branch line								
Cotton or vinyl rope	13.5	10.5	12.0	11.5	12.0	12.0	14.5	12.0
Ganged wire leader	6.0	6.0	8.0	6.0	7.0	8.0	7.0	7.0
Wire leader	1.5	2.0	2.5	1.5	2.0	2.0	3.5	2.5
Length of float line	18.5	19.5	20.0	16.5	22.0	18.0	22.5	25.0
Number of hooks	5	5	5	5	5	4	4	6
B. The estimated maximum depth of each hook of the lines shown above								
No. 1 hook	86.5	85.0	89.5	82.5	90.0	84.5	113.5	92.5
2	127.5	126.0	130.5	123.5	131.0	123.0	169.5	136.5
3	148.5	147.0	151.5	144.5	152.0	123.0	169.5	168.5
4	127.5	126.0	130.5	123.5	131.0	84.5	113.5	168.5
5	86.5	85.0	89.5	82.5	90.0			136.5
6								92.5

f the float line and branch line should not exceed 30 n.

As for the amount of gear fished per day, Yoshida 1966) reported that "Vessels of 39 to 99 gross tons fish 10 to 355 baskets; vessels 100 to 190 gross tons fish 55 to 400 baskets; and vessels 200 to 500 tons fish 400 o 450 baskets." Suda and Schaefer (1965) reported hat Japanese vessels in the eastern Pacific fished an verage of 2,000 hooks (about 400 baskets) per set.

The basic construction of the longline gear has emained unchanged over the years. However, due to nanpower problems some effort has been directed owards developing laborsaving devices in longlining. Two examples are the reel-type and tub-type of onglining.

In the reel-type, the mainline is continuously reeled nto a drum, while in the tub-type, the retrieved line s coiled into a large tub. Important advances in addi- ion to the line hauler, include the "slow-conveyor" for line setting and line hauling), the "guide stand" nd "guide roller" (for leading line), gear-transport- ng conveyor system, line-winder system, etc. In using hese methods, the branch lines are joined to the nainline by the use of snaps (Katsuo-Maguro Venkan, 1969).

The principal bait used in longlining is frozen Pacific saury, *Cololabis saira.* Squid is also commonly sed. Mackerel, *Scomber* sp., as well as mackerel cad, *Decapterus* sp., have been used as alternate ait. In addition, experiments are underway to utilize ilver carp, *Hypophthalmichthys molitrix*, as well as rtifical preparations (e.g., infused with extracts of aury) as longline bait.

Depending on the location, certain baits have been eported to have advantages in catching billfishes. For nstance, in the East China Sea fishing grounds, live nackerel were believed effective and were used exten- ively. However, Furukawa et al. (1957) reported that sing live or dead mackerel did not significantly affect the catches of white marlin (= black marlin), striped marlin, and broadbill.

The main piece of equipment in the harpoon fishery is the harpoon itself. The harpoon pole of oak is about 4 m long and at its tip is a three-pronged iron piece about 7 mm in diameter. The detachable harpoon of steel about 10 cm in length, connected to about 100 m of line, is placed over this iron tip. Recently the electric harpoon has been used in order to kill the fish quickly. When the harpoon enters the fish, a wire distributed along the harpoon line is charged with electricity.

5.12 Boats

The longline vessels fishing in the Indian and Pacific oceans for striped marlin are largely those from Japan. Other vessels are from Taiwan and Korea.

The details on the construction of longline vessels are given by Kanasashi (1960) and by Yoshida (1966). There are two types of longline vessels: those that use longline exclusively and those that use both longline and pole and line. The holds on longline vessels are not divided into small compartments to carry live bait. Thus, the hold space in the longliners is 20% to 40% greater than in the combination vessels.

Longline vessels are constructed of wood or steel; those larger than 100 gross tons are usually constructed of steel. Most of the longliners are 250 to 350 gross tons; at this stage of the fishery they appear to be the most economical and efficient size to operate. The specifications of typical longliners of this size class and those of some typical combination vessels are given in Yoshida (1966).

Other than the independently operating vessels, there is the mother ship operation in which several catcher boats are transported on the deck of a mother ship to the fishing grounds.

Small mother ships of 400 to 800 gross tons are able to carry only one portable catcher boat, but the larger 2,000- to 3,000-ton mother ships carry six catcher boats. The portable catcher boats measure about 15 m in length (Yoshida, 1966). To decrease the weight of the portable boats, recent constructions have been of fiber-reinforced plastic (FRP). The weight of a FRP boat is about 13 tons, or approximately two-thirds the weight of former catcher boats (Kazama, 1967).

Significant advances have also been made in preserving the catch of high-quality sashimi fish. Many vessels are equipped with refrigeration equipment capable of preserving the catch at very low temperatures of −40° to −45°C. In this way, the vessels are able to deliver fish in excellent condition and as a result, billfish prices have increased greatly in the Japanese market.

In addition, there have been advances in automating ship operations, fishing gear, and other equipment. Almost all longliners are equipped with fish detectors. Living conditions on the ships have also been improved considerably, and many vessels are now equipped with air conditioning for the crew's comfort (Katsuo-Maguro Nenkan, 1969).

Harpoon vessels are constructed with an extended prow where the harpooner is stationed. These vessels are constructed of wood, and range in size from about 10 to 40 tons. Billfishes taken by these vessels are kept in ice for delivery to the market.

5.2 Fishing Areas

5.21 General geographic distribution

The longline fishery now virtually covers the entire distribution of the species (see 2.1 and 2.22). The major sport fishing areas are southern California, Mexico, Panama, Ecuador, Peru, Chile, Hawaii, Tahiti, Fiji, New Zealand, Australia, and East Africa (not listed in order of importance).

5.22 Geographic ranges

See also 5.41.

Longlining is carried out across the high seas to within 5 miles of coastlines in places. The sport fishery is generally restricted to within about 75 miles of coastlines with the bulk of the fishing much closer.

The greatest fishing pressure is exerted in the North Equatorial Current in the Pacific and western Indian oceans, in the Kuroshio and Kuroshio extension, in the North Pacific Gyral northeast of the Hawaiian Islands, in the South Equatorial Current from about long. 90° to 140°W, and also off Mexico and Ecuador.

Regarding the development of the Japanese longline fishery, Suda and Schaefer (1965) report that prior to about 1952 the fishery was confined to the western and central Pacific. After this date it expanded into the Indian Ocean extending west of long. 80°E in 1954 and throughout the Indian Ocean by the end of 1955. In the Pacific, the fishery expanded eastw between lat. 10°N and lat. 10°S reaching 130°W late 1956 and long. 85°W by 1961. After 1963 fishery in the eastern Pacific expanded rapi poleward, the northward expansion being prima for striped marlin (Kume and Joseph, 1969a).

The fishing grounds for the Japanese harp fishery are located in the waters of Sanriku northeast of Honshu), around Izu, and East Ch Sea.

5.23 Depth range

Little has been written regarding fishing effort depth range. Merrett (1968a) reports almost half the striped marlin caught during experimen longlining off East Africa were caught over less th 1,000 fathoms even though most of the effort was pended beyond this depth contour.

5.24 Conditions of the grounds

See sections 2.1, 2.2, and 2.3.

5.3 Fishing Seasons

For sections 5.31, 5.32, and 5.33 see section 2.2

The fishing season for the Japanese harpoon fish in the waters of the Sanriku fishing ground exter from June to November with the peak occurring fr July through September; in the Izu area fr December to August with its peak from Febru through April; and in the East China Sea fr December to February.

5.4 Fishing Operations and Results

5.41 Effort and intensity

Type of unit effort: Detailed data on fishing eff and catch in the Japanese tuna longline fishery a published in the "Annual report of effort and cat statistics by area on Japanese tuna longline fisher by the Research Division, Fisheries Agency of Japa Fishing effort is reported in terms of number operations and number of hooks fished; catch reported in terms of number of fish. The statistics a reported on a monthly basis by 5° units.

Since 1967, Taiwan has also begun to publish da from their tuna longline fishery, following the sa format as the Japanese publication. The Taiwan da are published annually in "Report on survey production and marketing of Taiwan's tuna longli fishery" by the Taiwan Fisheries Bureau. Publicati of effort and catch statistics of the Korean longli fishery started in 1970 in "Yearbook of catch and e fort statistics on Korean tuna longline fishery" issue by the Office of Fisheries, Korea.

Landings per unit of fishing effort: As noted abov the catch statistics are reported in terms of numbe

f fish taken in the various unit areas. Landings by eight can be estimated from the data along with verage body weight data.

atches per unit of fishing effort: The catch per unit f effort (CPUE) can be obtained in terms of catch in umbers per 100 or 1,000 hooks fished. Strasburg 1970) studied year-to-year changes in CPUE of illfishes in the Pacific by analyzing the 1953-63 apanese longline data for quadrangles measuring 20° f latitude and longitude. He noted a progressive ecline over the years in the CPUE for the striped arlin of western and central South Pacific (lat. 20°-0°S) areas. Strasburg (1970) concluded that while me workers have attributed the decline in CPUE to eavy fishing, "It is impossible to determine its real ause without more information on various biological atures related to migration, reproduction, age, and ear classes."

Honma and Suzuki (1969) studied the apparent bundance of striped marlin in the principal fishing rounds in the Pacific (northwestern Pacific, eastern acific, and waters east of Australia) for the years 960-66. They noted no apparent trends in CPUE in ie northwestern Pacific and eastern Pacific grounds. n the other hand, for waters east of Australia, they eported a definite decreasing trend; the CPUE has een at a low level of around 0.1 fish per 100 hooks nce 1964.

Of the striped marlin in the Indian Ocean, Kikawa al. (1969) examined the annual changes in CPUE umber of fish per 1,000 hooks) based on data for the ears 1962-67. They reported CPUE of about 0.6 fish er 1,000 hooks between 1962 and 1964, followed by an crease to about 1.0 in 1965-67. The authors concluded that "An increasing trend in the CPUE for striped arlin in this period may probably represent the increase in effectiveness in catching fish."

shing effort per unit area: Figure 8 shows the distribution of fishing effort of the Japanese tuna ngline vessels in 1970. The effort, in terms of numbers of hooks fished, is shown by 5° quadrangles.

If we examine the data along with catch per unit area data of striped marlin (Fig. 9) it is apparent that the relatively large fishing effort in the eastern Pacific off Mexico and Ecuador is principally related to pursuit of striped marlin in those areas.

Total fishing intensity: The total fishing effort of the Japanese longline vessels in recent years (1965-69) for the Pacific and Indian oceans is estimated at about 400 million hooks fished per year. The efforts by areas are given in Table 5 (see 4.23).

Since 1963 there has been a significant increase in fishing effort in the eastern Pacific region. Correspondingly the fishing grounds for striped marlin also has increased during this period.

In the South Pacific region, however, effort has decreased from about 100 millin hooks in 1965-66 to around 70 million hooks beginning in 1967.

5.42 Selectivity

Small striped marlin under 80 cm eye-fork length are virtually never taken by longline.

Furukawa et al. (1958) have reported that striped marlin taken by harpooning in the East China Sea fishing grounds are relatively heavier (higher fatness index) than fish taken by longline.

5.43 Catches

Total annual yields: The FAO "Yearbook of fishery statistics" reports the total annual yields of striped marlin; Table 8 summarizes the 1970 catch (FAO, 1971). The recent total annual landings of striped marlin have been around 25,000 tons from the Pacific and Indian oceans.

Total annual yields from different fishing grounds: As seen in Table 8, the striped marlin catches are high in the eastern Pacific; the catch from this area comprises about one-half of the total Pacific

Table 8.—Catches of striped marlin by fishing areas for 1964-70. (From FAO, 1971.)

Area	1964	1965	1966	1967	1968	1969	1970
	In thousands of metric tons						
Indian Ocean	(2.4)	(3.8)	(3.8)	(6.2)	(3.8)	(4.3)	(3.1)
Western	1.2	2.1	3.2	4.9	2.2	2.5	2.1
Eastern	1.2	1.7	0.6	1.3	1.6	1.8	1.0
Pacific Ocean	(25.6)	(22.5)	(20.5)	(19.5)	(21.6)	(20.5)	(22.1)
Northwest	8.6	8.8	7.0	6.9	7.2	8.5	8.4
Eastern central	14.0	11.3	9.1	10.4	11.0	8.7	10.9
Southwest	1.4	1.3	2.3	0.9	1.7	1.6	1.6
Southeast	1.6	1.1	2.1	1.3	1.7	1.7	1.2
Total	28.0	26.3	24.3	25.7	25.4	24.8	25.2



Figure 8.—Distribution of estimated total fishing effort, in numbers of hooks per unit area (1970).



Figure 9.—Distribution of the catch of striped marlin in numbers per unit area (1970). Catch data in Atlantic Ocean area pertains to white marlin, *Tetrapturus albidus*.

landings. The Indian Ocean landings of striped marlin amount to approximately 20% of the Pacific landings.

Maximum equilibrium yield: Honma and Suzuki (1969) made some preliminary determinations of the maximum sustainable yield on the basis of 1960-66 data. They suggested that striped marlin catches in the Pacific can probably be increased about 60% over the recent average annual landings.

However, judging by the fact that catches have leveled off in recent years after fishing effort has virtually covered all of the known areas of striped marlin distribution, it appears that the present catches may be close to the maximum equilibrium yield level.

Obviously, further research is needed on this subject.

6 PROTECTION AND MANAGEMENT

6.1 Regulatory (Legislative) Measures

6.11 Limitation on reduction of total catch

The Japanese tuna longline fishery is regulated in terms of fleet size by the vessel licensing system. Licensing is reviewed at 5-yr intervals and fleet size is governed on the basis of the condition of the tuna resources. No increase in fleet size has been permitted since 1963. However, other countries such as Korea and Taiwan have increased the size of their longline fleet.

7 POND FISH CULTURE

Not applicable.

LITERATURE CITED

ANONYMOUS.
1955. Fish contamination by radioactivity. *In* Research report of the effects and influences of the radioactivity caused by the H-bomb tests at Bikini in 1954, p. 101-121.

ARATA, G. F., JR.
1954. A contribution to the life history of the swordfish, *Xiphias gladius* Linnaeus, from the South Atlantic coast of the United States and the Gulf of Mexico. Bull. Mar. Sci. Gulf Caribb. 4:183-243.

BAKER, A. N.
1966. Food of marlins from New Zealand waters. Copeia 1966:818-822.

BARRETT, I., and A. A. WILLIAMS.
1965. Hemoglobin content of the blood of fifteen species of marine fishes. Calif. Fish Game 51:216-218.

CRAIG, W. L.
1972. Game fish bulletin. Calif. Res. Agency, Dep. Fish Game, Long Beach, Calif., 3 p. [Processed.]

DELFIN, F. T.
1901. Catalogo de los pesces de Chile. Rev. Chil. Hist. Nat. 3 and 4:1-133.

DERANIYAGALA, P. E. P.
1951. The Istiophoridae and Xiphiidae of Ceylon. Spe Zeylan. Bull. Natl. Mus. Ceylon 26:137-142.
1952. A colored atlas of some vertebrates from Ceylon. 1. Fishes. Ceylon Natl. Mus. Publ., 149 p.

DE SYLVA, D. P.
1962. Red-water blooms off northern Chile, April-May 19 with reference to the ecology of the swordfish and the stri marlin. Pac. Sci. 16:271-279.

ELDRIDGE, M. B., and P. G. WARES.
1974. Some biological observations of billfishes taken in eastern Pacific Ocean, 1967-1970. *In* R. S. Shomura and Williams (editors), Proceedings of the International Billf Symposium, Kailua-Kona, Hawaii, 9-12 August 1972, P 2. Review and Contributed Papers, p. 89-101. U.S. D Commer., NOAA Tech. Rep. NMFS SSRF-675.

EVANS, D. H., and P. G. WARES.
1972. Food habits of striped marlin and sailfish off Mex and southern California. U.S. Fish Wildl. Serv., Res. R 76, 10 p.

FIERSTINE, H. L., and V. WALTERS.
1968. Studies in locomotion and anatomy of scombroid fish Mem. South Calif. Acad. Sci. 6, 31 p.

FISHERIES AGENCY OF JAPAN, RESEARCH DIVISION.
1965. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1962. 183 p.
1966. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1963. 322 p.
1967a. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1964. 379 p.
1967b. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1965. 375 p.
1968. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1966. 299 p.
1969. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1967. 293 p.
1970. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1968. 283 p.
1971. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1969. 299 p.
1972. Annual report of effort and catch statistics by area Japanese tuna longline fishery, 1970. 326 p.

FAO (FOOD AND AGRICULTURAL ORGANIZATION O THE UNITED NATIONS).
1971. Yearbook of fishery statistics. Catches and landing 1970. FAO, Vol. 30, 476 p.

FURUKAWA, I., N. ANRAKU, Y. KUROHIJI, T. KOTO, and K KODAMA.
1957. Study on the tuna long line fishery in the East Chi Sea - I. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Re Lab. 6:46-76.

FURUKAWA, I., T. KOTO, and K. KODAMA.
1958. Study on the tuna long line fishery in the East Chi Sea - II. Ecology of the striped marlin. [In Jap., En synop.] Rep. Nankai Reg. Fish. Res. Lab. 8:49-60.

GEHRINGER, J. W.
1956. Observations on the development of the Atlantic sa fish *Istiophorus americanus* (Cuvier) with notes on an u identified species of istiophorid. U.S. Fish Wildl. Ser Bull. 57:139-171.

GOSLINE, W. A., and V. E. BROCK.
1960. Handbook of Hawaiian fishes. Univ. Hawaii Pres Honolulu, 372 p.

GREGORY, W. K., and G. M. CONRAD.
1939. Body-forms of the black marlin (*Makaira nigrica marlina*) and striped marlin (*Makaira mitsukurii*) of Ne

Zealand and Australia. Bull. Am. Mus. Nat. Hist. 76:443-456.

GREY, Z.
1928. Big game fishing in New Zealand seas. Nat. Hist. 28: 46-52.

HIRASAKA, K., and H. NAKAMURA.
1947. On the Formosan spearfishes. Bull. Oceanogr. Inst. Taiwan 3:9-24.

HONMA, M., and T. KAMIMURA.
1958. A population study on the so-called makajiki (striped marlin) of both northern and southern hemispheres of the Pacific. II. Fishing conditions in the southern hemisphere. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 8:12-21.

HONMA, M., and J. SUZUKI.
1969. Distribution and catches of striped marlin caught by long-line in the Pacific Ocean 1960-1966. [In Jap.] Tuna Fish. (82 old series no.):9-22.

HOWARD, J. K., and S. UEYANAGI.
1965. Distribution and relative abundance of billfishes (*Istiophoridae*) of the Pacific Ocean. Univ. Miami Inst. Mar. Sci., Stud. Trop. Oceanogr. 2, 134 p.

HUBBS, C. L., and R. L. WISNER.
1953. Food of marlin in 1951 off San Diego, California. Calif. Fish Game 39(1):127-131.

JONES, E. C.
1971. *Isistius brasiliensis*, a squaloid shark, the probable cause of crater wounds on fishes and cetaceans. Fish. Bull., U.S. 69:791-798.

JONES, S., and M. KUMARAN.
1964. Distribution of larval billfishes (Xiphiidae and Istiophoridae) in the Indo-Pacific with special reference to the collections made by the Danish Dana Expedition. Proc. Symp. Scombroid Fishes, Part 1, p. 483-498. Mar. Biol. Assoc. India, Mandapam Camp.

JONES, S., and E. G. SILAS.
1964. A systematic review of the scombroid fishes of India. Proc. Symp. Scombroid Fishes, Part 1, p. 1-105. Mar. Biol. Assoc. India, Mandapam Camp.

JORDAN, D. S., and B. W. EVERMANN.
1926. A review of the giant mackerel-like fishes, tunnies, spearfishes and swordfish. Calif. Acad. Sci., Occas. Pap. 12:1-113.

JORDAN, D. S., and J. O. SNYDER.
1901. Descriptions of nine new species of fishes contained in museums of Japan. J. Coll. Sci., Imp. Univ. Tokyo 15:301-311.

KAMIMURA, T., and M. HONMA.
1958. A population study of the so-called makajiki (striped marlin) of both northern and southern hemispheres of the Pacific. I. Comparison of external characters. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 8:1-11.

KANASASHI, Y.
1960. Longline fishing: Deck design and equipment. *In* J.-O. Traung (editor), Fishing boats of the world, 2:73-83. Fish. News (Books) Ltd., Lond.

KATSUO-MAGURO NENKAN.
1969. Katsuo-Maguro Nenkan (Skipjack-Tuna Yearbook). [In Jap.] Suisan Shinchosha, Tokyo, 950 p.

KAZAMA, J.
1967. Recent developments in Japanese tuna longliners. *In* J.-O. Traung (editor), Fishing boats of the world, 3:550-557. Fish. News (Books) Ltd., Lond.

KIKAWA, S., T. KOTO, C. SHINGU, and Y. NISHIKAWA.
1969. Status of tuna fisheries in the Indian Ocean as of 1968. Far Seas Fish. Res. Lab., S Ser. (2), 28 p.

KOBAYASHI, H., and Y. YAMAGUCHI.
1971. Feeding ecology and the hooking tendency of tunas and marlins in the eastern equatorial Pacific. [In Jap., Engl. abstr.] Bull. Jap. Soc. Sci. Fish. 37:83-89.

KOGA, S.
1967. Studies on the fishery biology of the tuna and marlin in the Indian Ocean and the South Pacific Ocean. [In Jap., Engl. summ.] J. Shimonoseki Univ. Fish. 15(2):1-208.
1968. A study of the fishing conditions of the tuna and marlin in the Tasman Sea. [In Jap., Engl. abstr.] J. Shimonoseki Univ. Fish. 16(2):1-20.

KOTO, T.
1963a. Some considerations on the growth of marlins, using size frequencies in commercial catches. III. Attempts to estimate the growth of striped marlin, *Tetrapturus audax* (Philippi) in the western North Pacific Ocean. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 17:63-85.
1963b. Annual variations in the growth of striped marlin, *Tetrapturus audax* (Philippi) in the western North Pacific Ocean. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 17:87-92.

KUME, S., and J. JOSEPH.
1969a. The Japanese longline fishery for tunas and billfishes in the eastern Pacific Ocean east of 130°W, 1964-1966. [In Engl. and Span.] Bull. Inter-Am. Trop. Tuna Comm. 13:277-418.
1969b. Size composition and sexual maturity of billfish caught by the Japanese longline fishery in the Pacific Ocean east of 130°W. [In Engl., Jap. summ.] Bull. Far Seas Fish. Res. Lab. (Shimizu) 2:115-162.

KUME, S., and M. B. SCHAEFER.
1966. Studies on the Japanese long-line fishery for tuna and marlin in the eastern tropical Pacific Ocean during 1963. [In Span. and Engl.] Bull. Inter-Am. Trop. Tuna Comm. 11:103-170.

LAMONTE, F. R.
1955. A review and revision of the marlins, genus *Makaira*. Bull. Am. Mus. Nat. Hist. 107:323-358.

LINDSEY, C. C.
1968. Temperatures of red and white muscle in recently caught marlin and other large tropical fish. J. Fish. Res. Board Can. 25:1987-1992.

MANNING, J. A.
1957. Summary investigations of the pelagic fish survey of Chilean waters with special reference to the swordfish, marlins and tunas. Q. Rep. Fish. Mar. Lab., Fla. Board Conserv., Univ. Miami Publ. 57-4, 12 p.

MATSUBARA, K.
1955. Fish morphology and hierarchy. Part I. [In Jap.] Ishizaki-Shoten, Tokyo, 789 p.

MERRETT, N. R.
1968a. Tuna longline survey in the equatorial western Indian Ocean. East Afr. Agric. For. J. 34:17-65.
1968b. Tuna fishing research in the Indian Ocean. Africana 3(7):20-23.
1968c. Weight-length relationships for certain scombroid fishes from the equatorial western Indian Ocean. East Afr. Agric. For. J. 34:165-169.
1970. Gonad development in billfish (Istiophoridae) from the Indian Ocean. J. Zool. (Lond.) 160:355-370.
1971. Aspects of the biology of billfish (Istiophoridae) from the equatorial western Indian Ocean. J. Zool. (Lond.) 163: 351-395.

MORITA, T.
1969. Studies on the fishing gear of tuna long-line. [In Jap., Engl. abstr.] Mem. Fac. Fish., Kagoshima Univ. 18:145-215.

MORROW, J. E.
1952a. Allometric growth in the striped marlin, *Makaira mitsukurii*, from New Zealand. Pac. Sci. 6:53-58.
1952b. Food of the striped marlin, *Makaira mitsukurii*, from New Zealand. Copeia 1952:143-145.
1957. Races of the striped marlin, *Makaira mitsukurii*, in the Pacific. Bull. Bingham Oceanogr. Collect., Yale Univ. 16(2):72-87.
1964. Marlins, sailfish and spearfish of the Indian Ocean. Proc. Symp. Scombroid Fishes, Part 1, p. 429-440. Mar. Biol. Assoc. India, Mandapam Camp.
MUNRO, I. S. R.
1955. The marine and fresh water fishes of Ceylon. Dep. External Affairs, Canberra, 351 p.
NAKAMURA, H.
1938. Report of an investigation of the spearfishes of Formosan waters. [In Jap.] Reports of the Taiwan Government-General Fishery Experiment Station, No. 10, p. 1-34. (Translated by W. G. Van Campen, 1955, U.S. Dep. Int., Fish Wildl. Serv., Spec. Sci. Rep. Fish. 153, 46 p.)
1949. The tunas and their fisheries. [In Jap.] Takeuchi Shōbo, Tokyo, 118 p. (Translated by W. G. Van Campen, 1952, U.S. Dep. Int., Fish Wildl. Serv., Spec. Sci. Rep. Fish. 82, 115 p.)
1969. Tuna distribution and migration. Fish. News (Books), Ltd., Lond., 76 p.
NAKAMURA, H., Y. YABUTA, and S. UEYANAGI.
1953. Relation between the spawning season and the sex-ratio of some fishes of the family Istiophoridae. [In Jap., Engl. abstr.] Contrib. Nankai Reg. Fish. Res. Lab. 1(13), 8 p.
NAKAMURA, I.
1968. Juveniles of the striped marlin, *Tetrapturus audax* (Phillipi). Mem. Coll. Agric., Kyoto Univ. 94:17-29.
NAKAMURA, I., T. IWAI, and K. MATSUBARA.
1968. A review of the sailfish, spearfish, marlin and swordfish of the world. [In Jap.] Kyoto Univ., Misaki Mar. Biol. Inst., Spec. Rep. 4, 95 p.
NISHIKAWA, Y., and S. UEYANAGI.
1969. Spawning areas of the billfishes. [In Jap.] Tuna Fish. 3:13-16.
NISHIMURA, K., and N. ABE.
1971. A study on fishing conditions of the striped marlin around the Izu Island. [In Jap., Engl. abstr.] Bull. Jap. Soc. Sci. Fish. 37:1062-1066.
PARIN, N. V.
1968. Ikhtiofauna okeanoskoi epipelagiali (Ichthyofauna of the epipelagic zone). Izdatel'stov "NAUKA" Moscow. (Translated by M. Raveh, Israel Program Sci. Transl., 1970, 206 p.; available U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA, as TT 69-59020.)
PHILIPPI, R. A.
1887. Sobre los tiburones y algunos otros pesces de Chile. Anal. Univ. Chile 71:34-39.
POEY, F.
1851. Memorias sobre la historia natural de la isla de Cuba, acompañadas de sumarios latinos y extractos en frances. Imprenta de la Viuda de Barcina, Habana, Vol. 1, 463 p.
1856-58. Memorias sobra la historia natural de la isla de Cuba, acompañadas de sumarios latinos y extractos en frances. Imprenta de la Viuda de Barcina, Habana, Vol. 2, 442 p.
RAFINESQUE, C. S.
1810. Caratteri di alcuni nuovi generi e nuove specie di animali e piante della Sicilia, con varie osservazioni sopra i medesimi. Palermo, 105 p.
ROBINS, C. R., and D. P. DE SYLVA.
1960. Description and relationships of the longbill spearfish, *Tetrapturus belone*, based on Western North Atlantic specimens. Bull. Mar. Sci. Gulf Caribb. 10:383-413.
1963. A new western Atlantic spearfish, *Tetrapturus pfluegeri*, with a redescription of the Mediterranean spearfish *Tetrapturus belone*. Bull. Mar. Sci. Gulf Caribb. 13:84-122.
ROYCE, W. F.
1957. Observations on the spearfishes of the central Pacific. U.S. Fish Wildl. Serv., Fish. Bull. 57:497-554.
SAITO, S., G. KOBAYASHI, H. YASUMA, and S. SASAKI.
1972. Swimming depths of large sized albacore in the South Pacific Ocean - II. Depths of capture of albacore by improved vertical longline gear. [In Jap.] Proceedings, 1971 Tuna Research Conference, Shimizu, Japan, 2-4 February 1971. Fish. Agency, Far Seas Fish. Res. Lab., p. 144-147.
SMITH, J. L. B.
1956a. The striped marlin (*Makaira audax* Phillipi) in South Africa. Nature (Lond.) 177(4512):758.
1956b. Swordfish, marlin and sailfish in South and East Africa. Rhodes Univ., Dep. Ichthyol., Ichthyol. Bull. 25-34.
SQUIRE, J. L., JR.
1974. Migration patterns of Istiophoridae in the Pacific Ocean as determined by cooperative tagging programs. *In* R. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972, Part 2. Review and Contributed Papers, p. 22-237. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-675.
STRASBURG, D. W.
1970. A report on the billfishes of the central Pacific Ocean. Bull. Mar. Sci. 20:575-604.
SUDA, A., and M. B. SCHAEFER.
1965. General review of the Japanese tuna long-line fishery in the eastern tropical Pacific Ocean 1956-1962. [In Span. and Engl.] Bull. Inter-Am. Trop. Tuna Comm. 9:307-462.
TALBOT, F. H., and M. J. PENRITH.
1962. Tunnies and marlins of South Africa. Nature (Lond.) 193:558-559.
TANAKA, S.
1914. Figures and descriptions of the fishes of Japan, including Riukiu Islands, Bonin Islands, Formosa, Kurile Islands, Korea, and South Sakhalin. [In Jap. and Engl.] Dai Nihon Suisan Shoin 18:295-318. Tokyo.
UEYANAGI, S.
1953. The sexual difference of the sizes of marlins. [In Jap., Engl. abstr.] Contrib. Nankai Reg. Fish. Res. Lab. 1(14), 5 p.
1957a. Young of the black marlin, *Eumakaira nigra* Nakamura. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 6:91-102.
1957b. On *Kajikia formosana* (Hirasaka et Nakamura). [In Jap., Engl. abstr.] Rep. Nankai Reg. Fish. Res. Lab. 6:103-112.
1959. Larvae of the striped marlin, *Makaira mitsukurii* (Jordan et Snyder). [In Jap., Engl. summ.] Rep. Nankai Reg. Fish. Res. Lab. 11:130-146. (Translated by W. G. Van Campen, 1961, 20 p.; available at Southwest Fisheries Center, National Marine Fisheries Service, NOAA, Honolulu, Hawaii.)
1963a. Methods for identification and discrimination of the larvae of five istiophorid species distributing in the Indo-Pacific. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 17:137-150.
1963b. A study of the relationships of the Indo-Pacific istiophorids. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 17:151-165.

1964. Description and distribution of larvae of five istiophorid species in the Indo-Pacific. Proc. Symp. Scombroid Fishes, Part 1, p. 499-528. Mar. Biol. Assoc. India, Mandapam Camp.

1974. On an additional diagnostic character for the identification of billfish larvae with some notes on the variations in pigmentation. *In* R. S. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972, Part 2. Review and Contributed Papers, p. 73-78. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-675.

EYANAGI, S., and H. WATANABE.

1965. On certain specific differences in the vertebrae of the istiophorids. [In Jap., Engl. synop.] Rep. Nankai Reg. Fish. Res. Lab. 22:1-8.

ARES, P. G., and G. T. SAKAGAWA.

1974. Some morphometrics of billfishes from the eastern Pacific Ocean. *In* R. S. Shomura and F. Williams (editors), Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972, Part 2. Review and Contributed Papers, p. 107-120. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-675.

WHITLEY, G. P.

1937. Studies in ichthyology. No. 10. Rec. Aust. Mus. 20(1): 3-24.

WILLIAMS, F.

1967. Longline fishing for tuna off the coast of east Africa 1958-1960. Indian J. Fish., Sect. A, 10:233-390.

WISNER, F. L.

1958. Is the spear of istiophorid fishes used in feeding? Pac. Sci. 12:60-70.

YABUTA, Y.

1953. On the stomach contents of tuna and marlin from the adjacent seas of Bonin Islands. [In Jap.] Contrib. Nankai Reg. Fish. Res. Lab. 1(15), 6 p.

YAMAGUCHI, Y.

1969. Studies on the feeding habits of tunas. [In Jap., Engl. summ.] J. Fac. Fish. Prefect. Univ. Mie 8:1-15.

YOSHIDA, H. O.

1966. Tuna fishing vessels, gear, and techniques in the Pacific Ocean. *In* T. A. Manar (editor), Proc. Gov. Conf. Cent. Pac. Fish. Resour., p. 67-89. State of Hawaii.

☆ U. S. GOVERNMENT PRINTING OFFICE: 1975—698-822 /36 REGION 10

. Weight loss of pond-raised channel catfish (*Ictalurus punctatus*) during holding in cessing plant vats. By Donald C. Greenland and Robert L. Gill. December 1971, iii + 7 , 3 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government nting Office, Washington, D.C. 20402.

. Distribution of forage of skipjack tuna (*Euthynnus pelamis*) in the eastern tropical ific. By Maurice Blackburn and Michael Laurs. January 1972, iii + 16 pp., 7 figs., 3 les. For sale by the Superintendent of Documents, U.S. Government Printing Office, shington, D.C. 20402.

. Effects of some antioxidants and EDTA on the development of rancidity in Spanish ckerel (*Scomberomorus maculatus*) during frozen storage. By Robert N. Farragut. ruary 1972, iv + 12 pp., 6 figs., 12 tables. For sale by the Superintendent of cuments, U.S. Government Printing Office, Washington, D.C. 20402.

. The effect of premortem stress, holding temperatures, and freezing on the chemistry and quality of skipjack tuna. By Ladell Crawford. April 1972, iii + 23 pp., 3 ., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing ice, Washington, D.C. 20402.

. The use of electricity in conjunction with a 12.5-meter (Headrope) Gulf-of-Mexico imp trawl in Lake Michigan. By James E. Ellis. March 1972, iv + 10 pp., 11 figs., 4 les. For sale by the Superintendent of Documents, U.S. Government Printing Office, shington, D.C. 20402.

. An electric detector system for recovering internally tagged menhaden, genus voortia. By R. O. Parker, Jr. February 1972, iii + 7 pp., 3 figs., 1 appendix table. For e by the Superintendent of Documents, U.S. Government Printing Office, Washington, C. 20402.

. Immobilization of fingerling salmon and trout by decompression. By Doyle F. herland. March 1972, iii + 7 pp., 3 figs., 2 tables. For sale by the Superintendent of cuments, U.S. Government Printing Office, Washington, D.C. 20402.

. The calico scallop, *Argopecten gibbus*. By Donald M. Allen and T. J. Costello. May 2, iii + 19 pp., 9 figs., 1 table. For sale by the Superintendent of Documents, U.S. vernment Printing Office, Washington, D.C. 20402.

. Making fish protein concentrates by enzymatic hydrolysis. A status report on earch and some processes and products studied by NMFS. By Malcolm B. Hale. vember 1972, v + 32 pp., 15 figs., 17 tables, 1 appendix table. For sale by the perintendent of Documents, U.S. Government Printing Office, Washington, D.C. 02.

. List of fishes of Alaska and adjacent waters with a guide to some of their literature. Jay C. Quast and Elizabeth L. Hall. July 1972, iv + 47 pp. For sale by the Superinten- t of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

. The Southeast Fisheries Center bionumeric code. Part I: Fishes. By Harvey R. lis, Jr., Richard B. Roe, and Judith C. Gatlin. July 1972, xl + 95 pp., 2 figs. For sale by Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 02.

. A freshwater fish electro-motivator (FFEM)-its characteristics and operation. By nes E. Ellis and Charles C. Hoopes. November 1972, iii + 11 pp., 9 figs.

. A review of the literature on the development of skipjack tuna fisheries in the cen- and western Pacific Ocean. By Frank J. Hester and Tamio Otsu. January 1973, iii + pp., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Of- , Washington, D.C. 20402.

662. Seasonal distribution of tunas and billfishes in the Atlantic. By John P. Wise and Charles W. Davis. January 1973, iv + 24 pp., 13 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

663. Fish larvae collected from the northeastern Pacific Ocean and Puget Sound during April and May 1967. By Kenneth D. Waldron. December 1972, iii + 16 pp., 2 figs., 1 table, 4 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

664. Tagging and tag-recovery experiments with Atlantic menhaden, *Brevoortia tyrannus*. By Richard L. Kroger and Robert L. Dryfoos. December 1972, iv + 11 pp., 4 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

665. Larval fish survey of Humbolt Bay, California. By Maxwell B. Eldridge and Charles F. Bryan. December 1972, iii + 8 pp., 8 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

666. Distribution and relative abundance of fishes in Newport River, North Carolina. By William R. Turner and George N. Johnson. September 1973, iv + 23 pp., 1 fig., 13 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

667. An analysis of the commercial lobster (*Homarus americanus*) fishery along the coast of Maine, August 1966 through December 1970. By James C. Thomas. June 1973, v + 57 pp., 18 figs., 11 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

668. An annotated bibliography of the cunner, *Tautogolabrus adspersus* (Walbaum). By Fredric M. Serchuk and David W. Frame. May 1973, ii + 43 pp. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

669. Subpoint prediction for direct readout meteorological satellites. By L. E. Eber. August 1973, iii + 7 pp., 2 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

670. Unharvested fishes in the U.S. commercial fishery of western Lake Erie in 1969. By Harry D. Van Meter. July 1973, iii + 11 pp., 6 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

671. Coastal upwelling indices, west coast of North America, 1946-71. By Andrew Bakun. June 1973, iv + 103 pp., 6 figs., 3 tables, 45 appendix figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

672. Seasonal occurrence of young Gulf menhaden and other fishes in a northwestern Florida estuary. By Marlin E. Tagatz and E. Peter H. Wilkins. August 1973, iii + 14 pp., 1 fig., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

673. Abundance and distribution of inshore benthic fauna off southwestern Long Island, N.Y. By Frank W. Steimle, Jr. and Richard B. Stone. December 1973, iii + 50 pp., 2 figs., 5 appendix tables.

674. Lake Erie bottom trawl explorations, 1962-66. By Edgar W. Bowman. January 1974, iv + 21 pp., 9 figs., 1 table, 7 appendix tables.

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS STAFF
ROOM 450
1107 N.E. 45TH ST.
SEATTLE, WA 98105

OFFICIAL BUSINESS

FOURTH CLASS
book

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM-210